Supplementary Information

Ecological assets behind improving accounts: a boundary-matched test of composition masking

Amin Abaee
Independent Researcher, Tehran, Iran

S1 National account coverage and availability criteria

The source workbook is the National Ecological Footprint and Biocapacity Accounts, Edition 2026, results and metadata version 1.0. The `national_data` sheet contains 13652 records, 51 columns and 236 country and territory identifiers. The primary interval comprises nine annual observations, 2015–2023 inclusive. A country identifier satisfies the account availability criteria if:

1. every year in the interval is present with a unique country–year key;
2. each publication flag is `+++`, `++` or `+`;
3. all five biocapacity components, their total, forest area, cropland area and population are finite and nonnegative, with positive population;
4. the absolute difference between the five-component sum and the reported total does not exceed 3 gha.

The 3 gha bound follows from rounding five components and one total to the nearest hectare equivalent: the maximum discrepancy is $5(0.5) + 0.5 = 3$ gha. The largest observed discrepancy is 2 gha. Publication flags differ in trade and consumption detail; the three included flag classes all release the biocapacity and production fields used here. The workbook's actual field is named `flag`, while its definition sheet also uses the term `quality_flag`.

There are 183 identifiers satisfying these criteria and 53 not satisfying them. The complete identifier-level table records missing years, publication scope, required-value availability and component consistency. The selected countries in Table S1 all satisfy the account criteria. Their physical boundary and measurement compatibility are not determined by this result.

Table S1: Countries selected for evaluation of physical observation compatibility. Account availability refers to the complete 2015–2023 interval under the criteria in S1.

Country	ISO identifier	Account availability
Brazil	BRA	Complete
Indonesia	IDN	Complete
Democratic Republic of the Congo	COD	Complete
Cameroon	CMR	Complete
Canada	CAN	Complete
Finland	FIN	Complete
France	FRA	Complete
Australia	AUS	Complete

The selection is purposive rather than probabilistic. The availability result cannot be extrapolated to the prevalence of account–asset agreement or disagreement. The 2024–2025 workbook records do not form part of the primary interval. Account areas remain inputs to the account observation system rather than independent validation measurements of forest extent.

S2 Biomass products, units and spatial support

The source is ESA CCI Biomass v7.0, DOI https://doi.org/10.5285/6429d1aafe1e43b9b414e4a5a7f8b903. Metadata inspection used the 50 km aggregate and the 2010–2020 change product. Their exact names, byte sizes, hashes and retrieval URLs are supplied in the source register. Only coordinate variables, time values, dimensions and metadata were read; biomass values and change signs were not evaluated.

Table S2: Observation definitions in the inspected CCI products.

Attribute	Definition or interpretation
Grid	720 longitude by 360 latitude cells, with 0.5° spacing in WGS84 coordinates. The nominal 50 km designation is not an equal-area definition.
Epochs	2005–2012 and 2015–2024, comprising 18 annual epochs. No observed 2013 or 2014 map is supplied in this series.
Biomass density	Above-ground living woody biomass in $\mathrm{Mg\,ha^{-1}}$. The aggregate is normalized by total grid-cell area, not forest area.
Aggregate uncertainty	The guide describes uncertainty in the aggregated mean using variance and spatial covariance contributions. The variable name `agb_sd` alone does not establish an independent-pixel error model.
Change example	Difference, standard deviation and quality-class layers for 2010–2020. This example does not validate change estimates for 2015–2023.
Temporal bounds	The change file's time-bounds attribute names `time_bnds1`; no variable with that name is stored in the file. The documented epoch pair defines the interval considered here.
Observation coverage	The guide specifies a native product domain between 80°N and 60°S, subject to remote-sensing availability. The global coordinate rectangle is not a valid-observation mask.

Country–product intersections require a territorial definition, including overseas areas where relevant. Native-resolution or appropriately aggregated forest observations must distinguish no-data cells, observed zero values, nonforest cover and converted land. An estimated density of zero is not, by itself, an independent land-cover observation. Density averaging, mask application and area normalization must refer to the same support. Coarse aggregates do not identify fine-resolution forest-to-cropland transitions.

S3 Change-quality classes and uncertainty

The inspected provider implementation derives quality classes from endpoint biomass estimates and their uncertainty intervals, followed by a growth-limit condition and spatial filtering. The class definitions are summarized in Table S3. These labels describe the provider decision rule; they are not independent observations of ecological change.

Table S3: Provider change-quality classes. Descriptions refer to the processing rule rather than independently validated confidence levels.

Code	Interpretation
0	Both endpoint biomass estimates are zero
1	Strong loss classification
2	Possible loss classification
3	No resolved change under the processing rule
4	Possible gain classification
5	Strong gain classification

The example implementation computes change standard deviation as $\sqrt{\sigma_0^2 + \sigma_1^2}$ and uses one-standard-deviation endpoint intervals in its classification operations. In general,

$$\operatorname{Var}(\widehat{d}_1 - \widehat{d}_0) = \sigma_1^2 + \sigma_0^2 - 2\operatorname{Cov}(\widehat{d}_1, \widehat{d}_0). \tag{S1}$$

The variance sum corresponds to zero temporal covariance. A one-standard-deviation interval rule combined with growth and filtering operations does not define a calibrated 95% test. Temporal covariance, systematic biases and the decision rule's operating characteristics require separate evaluation for the specified observation domain.

For an area-weighted total $\widehat{Q} = \sum_i a_i \widehat{d}_i$ on a fixed support, the measurement variance is $\boldsymbol{a}^{\mathsf{T}}\Sigma\boldsymbol{a}$, where Σ is the joint error covariance matrix. Spatial independence reduces this to a sum of diagonal terms, but that simplification requires evidence. Uncertain support, missing or converted cells, and forest classification error introduce additional terms not represented by a density-only covariance matrix. Product-quality classes cannot provide independent validation when they are constructed from the target maps themselves.

S4 Geographic area calculation

For a rectangle bounded by longitude λ_w, λ_e and latitude ϕ_s, ϕ_n, all expressed in radians, the WGS84 area in hectares is

$$A = \frac{a^2(1-e^2)}{10^4}(\lambda_e - \lambda_w)\left[F(\sin\phi_n) - F(\sin\phi_s)\right], \tag{S2}$$

where $a = 6378137\,\mathrm{m}$, $f = 1/298.257223563$, $e^2 = f(2-f)$ and

$$F(s) = \frac{s}{2(1-e^2s^2)} + \frac{\operatorname{atanh}(es)}{2e}.$$

The formula integrates the ellipsoidal area element between meridians and parallels. Longitude intervals crossing the dateline are split before integration. Geographic cell area varies with latitude even when angular spacing is constant.

For the full ellipsoid, the rectangle integral reduces to

$$A_{\mathrm{WGS84}} = \frac{2\pi a^2}{10^4}\left[1 + \frac{1-e^2}{e}\operatorname{atanh}(e)\right].$$

Areas of adjacent latitude bands add exactly at the analytic level because their boundary terms telescope. These identities specify geometric consistency independently of ecological observation accuracy.

S5 Extent, density and spatial compensation

For positive forest areas, expanding the symmetric terms gives

$$\begin{aligned}\frac{(A_1 - A_0)(D_1 + D_0) + (D_1 - D_0)(A_1 + A_0)}{2} \\ = A_1D_1 - A_0D_0 = Q_1 - Q_0.\end{aligned} \tag{S3}$$

The identity assigns a pure area change entirely to the extent term and a pure mean-density change entirely to the density term. Mean forest density is undefined at a zero-area endpoint; appearing or disappearing strata therefore require a separately specified allocation convention.

For local net stock changes, $G - L = \sum_i \Delta Q_i$ and $G + L = \sum_i |\Delta Q_i|$. Consequently, $0 \le K \le 1$ when $G + L > 0$. A one-direction change gives $K = 0$, while equal positive and negative sums

give $K = 1$. The offset mass is $\min(G, L)$ and has biomass-mass units. When every local change is zero, K is undefined.

These quantities distinguish a national endpoint from its spatial composition, but measurement error can produce the same pattern. If true change is zero and observed local changes are errors e_i, positive and negative errors may yield a large $\sum_i |e_i|$ while $\sum_i e_i$ is close to zero. The resulting high cancellation fraction does not establish ecological redistribution. Interpretation therefore requires an error-model reference at the effective observational support. Resampling to a finer grid does not provide additional independent physical information.

The sums represent local net changes. Gross growth, recruitment, harvest and mortality require additional observations or a separately identified demographic model.

S6 Validation, information dependence and inferential scope

Account availability, physical observation compatibility and validation are distinct requirements. A physical comparison requires stable territorial boundaries, a justified forest support, matched observation dates and a change-error model. Transition validation must distinguish the target forest-to-cropland class from other disturbances and quantify uncertainty using an appropriate reference design. Biomass-level validation does not automatically establish biomass-change precision.

Practical-change and classification thresholds require calibration independent of held-out evaluation outcomes. The calibration design must evaluate the complete reporting rule under relevant covariance, systematic-bias and missingness scenarios; individual feature tests do not determine the error rate of the combined rule. Comparisons of reporting information require the same cases and weights. Full-population coverage and pairwise overlap must also be reported so that a score change is not attributed to a change in the assessed population. Input lineage must distinguish independent observations from target-derived features and shared calibration/reference data. Algebraically recoverable features do not constitute additional independent measurements.

The results supplied here are account availability counts, product definitions and computational verification. No country-level physical stock changes, conversion associations, ecological change-resolution probabilities or reporting-rule accuracy estimates are supplied. These quantities cannot be inferred from account publication flags or product access alone.

Source attribution and availability

National accounts: Lo, K., et al. (2026), National Ecological Footprint and Biocapacity Accounts, Edition 2026, results and metadata v1.0; https://footprint.info.yorku.ca/data/. Derived account tables and figures are distributed under CC BY-SA 4.0.

Biomass: Santoro, M., and Cartus, O. (2026), ESA CCI Biomass v7.0; https://doi.org/10.5285/6429d1aafe1e43b9b414e4a5a7f8b903. The ESA Climate Change Initiative, Biomass CCI project and data providers are acknowledged. Product definitions follow the Product User Guide v7.0, 19 March 2026; https://climate.esa.int/en/projects/biomass/key-documents/.

Source hashes, metadata, derived account-availability tables and computational methods are available at https://github.com/MIKEAA2020/general-sustainability/tree/main/batch%209/agent%202/ecomod_empirical_followup/F03. Raw data, the guide and provider code are not redistributed.