

The Fundamental Higher Consciousness Premise

The consolidation turns assembler: the parameter-map verdict corrected from “does not exist” to “no unified map, published fragments”; five load-bearing sources read from the supplied PDFs; and the concrete seven-phase Level-2 decombination protocol — coupling and criticality, anesthesia sweeps, receptor parameterization, patient-level inversion, the joint S1-S4 demonstration, and preregistered adjudication — every phase with named methods, open data, null models, and failure conditions.

Audit and Consolidation, Seventh Edition · Z.ai

Newly-read sources: the Virtual Epileptic Patient (Jirsa et al. 2017, NeuroImage); hierarchical whole-brain modeling of critical synchronization (Myrov et al., bioRxiv); serotonin receptor maps explaining the non-linear effects of LSD (Deco et al. 2018, Current Biology); ketanserin blockade attributing LSD to the 5-HT_{2A} receptor (Preller et al. 2018, eLife); metabolic connectivity mapping (Riedl et al., PNAS)

Method: systematic web search with fetch-verified links (48 URLs, 12 papers resolved), line-level reading of the supplied sources, assembly of the five program areas, and the protocol drafted against the decombination specification's four requirements

Scope: the neural parameter map assembled as far as the literature permits; the phase-parameters gap stated; the conventional-level ceiling maintained under the equivalence doctrine