

Contents

1	Introduction	3
1.1	The framework	3
1.2	Epistemic status: prove, construct, demote	3
1.3	Summary of results	4
2	The Arithmetic Sector	4
2.1	The arithmetic Hilbert space	4
2.2	The multiplicative number operator and the primon Hamiltonian	5
2.3	The partition function and the Hagedorn temperature	6
2.4	Gaussian extension	7
2.5	What the arithmetic sector does and does not assert	8
3	Dynamics: The Interaction Sector on the Prime Lattice	8
3.1	The frozen Hamiltonian	8
3.2	Multiplicative shift operators	8
3.3	The walk Hamiltonian and the prime lattice	9
3.4	The cascade: statics and dynamics	10
3.5	The Janus state	12
4	Recurrence, Periodicity, and the Fate of Eternal Return	14
4.1	Linear independence of the prime logarithms	14
4.2	Exact periodicity is forbidden	14
4.3	What is <i>not</i> forbidden: three qualifications	14
4.4	The verdict on cyclical time	15
5	The Arithmetic Clock and the Problem of Time	16
5.1	The problem of time, stated precisely	16
5.2	The zeta clock	16
5.3	Energy–time duality of the arithmetic clock	17
5.4	Reconciliation of the three times	18
6	Thermodynamics, Gravity, and the Holographic Bridge	18
6.1	The exact counting law	18
6.2	The holographic bridge	19
6.3	Singularities and the Hagedorn wall	20
6.4	Emergent gravity and the hierarchy problem	20
6.5	The classical limit and the pole at unity	21
7	Entanglement: Arithmetic Superpositions and Holism	21
7.1	Composite integers are product states	21
7.2	Splitting primes as natural entanglement resources	22
7.3	Measurement, decoherence, and records	23
7.4	Bell’s trilemma and the rejection of superdeterminism	23
8	The Arithmetic Gauge Sector	24
8.1	From Lie unification to Galois unification	24
8.2	The dictionary	25
8.3	Arithmetic charge conjugation, parity, and time reversal	25
8.4	Vaccaro’s program, translated	26
8.5	What Langlands does and does not supply	26

9	Arrows of Time and the Philosophy Register	27
9.1	The extrinsic arrow, structurally grounded	27
9.2	The hierarchy of arrows	28
9.3	Cyclical time: the verdict	28
9.4	Eternalism, indexicality, and the vertiginous question	29
9.5	Past Hypothesis as arithmetic necessity: theorem and conjecture	29
9.6	The Big Crunch, Gold universes, and reverse-arrow regions	29
9.7	Superdeterminism and locality	30
10	The Consolidated Conjecture Register	30
11	Falsifiability, Predictions, and Test Programs	35
11.1	What the framework predicts that rivals do not	35
11.2	Mathematical test programs	35
11.3	Assessment	36
A	Notation	37

Abstract

We develop the mathematical and physical content of the “prime-spectral framework”: the statistical mechanics of a free Bose gas of “primons” (one-particle energies $\log p$, p ranging over the rational primes), advanced as the arithmetic core of a unified picture of quantum gravity, time, and the arrows of time. The mathematical core is theorem-grade. The primon Hilbert space is ℓ^2 of the free commutative monoid on the primes, which sidesteps the nonseparability of naive infinite tensor products; the primon Hamiltonian is proved self-adjoint with simple spectrum $\hbar\omega_0 \log n$; the partition function is the Riemann zeta function, so the exponential state-counting law $N(E) = \lfloor e^{E/\hbar\omega_0} \rfloor$ is a *Hagedorn spectrum* with critical temperature $T_H = \hbar\omega_0/k_B$; and the “thermodynamic cascade into composite states” is grounded in the Hardy–Ramanujan and Erdős–Kac theorems. Three structural requirements are then met: the diagonal primon Hamiltonian admits no transitions, so an interaction sector (multiplication operators and hopping dynamics on the *prime lattice*) is introduced; composite integers are *not* entangled states (the occupation basis is a product basis), so arithmetic entanglement is defined across prime-mode bipartitions; and exact periodicity is forbidden by the $\mathbb{Q}$ -linear independence of prime logarithms while approximate recurrence survives in every finite truncation. Claims whose proof is out of reach are consolidated into an explicit, falsifiable conjecture register C1–C12, from the zeta/black-hole partition-function identification to a Galois–Langlands dictionary for the Standard Model. Two register entries are tested numerically at laptop scale: eigenstate-thermalization diagnostics for C4 on truncations up to $D \approx 6 \times 10^4$, and a calibrated Chebotarev-dictionary protocol for C9. The result is a framework whose theorems are proved or cited exactly, whose analogies are tightened to citable results, and whose remaining ambitions are stated with enough precision to be confirmed or refuted.

1 Introduction

1.1 The framework

The prime-spectral framework proposes that the prime numbers, through the primon gas of Julia [1] and Spector [2], supply the quantum-mechanical spectrum of the near-singularity gravitational field, that the Riemann zeta function is the gravitational partition function, and that the architecture of time—its problem, its arrow, its block-universe metaphysics, and its impossibility of cycles—follows from the arithmetic of the integers. Recent developments linking conformal primon gases to black-hole interiors [3], the philosophical literature of time’s arrow [4–6], and first-person metaphysics [7–9] motivate the construction. Its ambitions extend to Gaussian primes as a five-dimensional chirality-carrying sector, the Absolute Galois group as a grand unified symmetry, and gravity as an entropic, $1/N$ -suppressed collective effect.

The framework alternates between genuine mathematics, physics speculation, and rhetoric; the skeleton is unusually coherent, while the joints are loose. This paper keeps the skeleton, tightens every joint, and labels everything else. Three instruments suffice: state precisely what can be proved; construct what is missing; and demote what remains to falsifiable conjectures.

1.2 Epistemic status: prove, construct, demote

Every substantive statement in this paper receives one of three labels.

1. **Prove.** A statement with a precise mathematical formulation and a proof is stated as a theorem, lemma, or proposition with its proof or an exact citation. Examples: the $\mathbb{Q}$ -linear independence of $\{\log p\}$; the identification of the partition function with ζ ; the Hagedorn asymptotics; the positive-definiteness of the Page–Wootters clock kernel; the exact state-counting identity $\log N(E) = E/\hbar\omega_0 + O(1)$.

2. **Construct.** Where a mechanism is missing, it is supplied and its consequences proved. Three constructions dominate this paper: (i) the diagonal primon Hamiltonian cannot drive a “cascade into composite states”, so a multiplicative transition sector (multiplication operators M_n and a hopping dynamics on the *prime lattice*) is introduced in Section 3; (ii) composite integers are *not* entangled states—the occupation basis is a product basis—so arithmetic entanglement is defined across prime-mode bipartitions in Section 7, and “arithmetic holism” is relocated from single integers to superpositions; (iii) the “universe cannot repeat” claim conflates exact periodicity, Poincaré recurrence, and dynamical return; these are separated in Section 4 into one theorem and two counterexamples.
3. **Demote.** A claim that is plausible, well-formed, and beyond current proof technology is restated as a numbered conjecture with evidence, dependencies, and a falsifier. Twelve conjectures (C1–C12) are consolidated in Section 10. The guiding rule is that a conjecture must be *decidable*: each one carries at least one specified route to confirmation or refutation.

Numerical evidence is held to the same standard: every diagnostic is defined before it is computed, and finite-size results are reported with their uncertainty and their dependence on protocol choices (window placement, pivoting, trajectory horizon).

1.3 Summary of results

The framework acquires a layered structure. The *mathematical floor* (all theorem-grade) consists of: the arithmetic Hilbert space $\mathcal{H}_A = \ell^2(\mathbb{N}, \times)$ (Section 2); the self-adjoint primon Hamiltonian with simple spectrum $\hbar\omega_0 \log n$; the partition function $Z(\beta) = \zeta(\beta\hbar\omega_0)$ with Hagedorn temperature $T_H = \hbar\omega_0/k_B$; the exact counting law $\log N(E) = E/\hbar\omega_0 + O(1)$; the Erdős–Kac statistics of the microcanonical ensemble; the isometric representation $n \mapsto M_n$ of the multiplicative monoid on $\mathcal{H}_A$; the phase-incommensurability theorem forbidding exact periodicity; the Bohr almost-periodicity of the zeta clock’s correlation kernel; and the product-state theorem that settles the entanglement question. The *conjectural superstructure* consists of the twelve register entries, of which the deepest are C1 (the zeta/black-hole identification), C3 (the arithmetic holographic bound), C4 (thermalization on the prime lattice), and C9 (the Galois–gauge dictionary). Two entries carry carried-out numerical programs: eigenwindow and Krylov diagnostics for C4 to $D \approx 6 \times 10^4$ (Remark 3.9), and a calibrated dictionary protocol for C9 (Section 11).

The philosophical theses of the framework—eternalism with perspectival observers, an extrinsic arrow grounded in arithmetic state-counting, derivative causality, and the rejection of superdeterminism—survive in *conditional* form: they are well-defined consequences of the conjectural superstructure, and they are marked as interpretive rather than theorem-grade wherever the mathematics cannot carry them (Section 9).

2 The Arithmetic Sector

This section fixes the foundational objects. Everything here is elementary but must be stated carefully, because a naive formulation writes the primon Fock space as an infinite tensor product, an object that is *not* separable and is the wrong home for the theory. The construction is cheap and clarifying.

2.1 The arithmetic Hilbert space

Let $\text{Pr} = \{2, 3, 5, \dots\}$ denote the rational primes. Let

$$\mathcal{F} = \left\{ k: \text{Pr} \rightarrow \mathbb{N} : k_p \neq 0 \text{ for only finitely many } p \right\} \quad (1)$$

be the free commutative monoid on the primes. By the fundamental theorem of arithmetic, the map $\mathcal{F} \rightarrow \mathbb{N}_{\geq 1}$, $k \mapsto \prod_p p^{k_p}$, is a monoid isomorphism between $(\mathcal{F}, +)$ and $(\mathbb{N}_{\geq 1}, \times)$, where $+$ denotes componentwise addition of occupation vectors. This single classical theorem is the load-bearing wall of the entire framework: it is what makes “states of the arithmetic universe” and “positive integers” the same bookkeeping.

Definition 2.1 (Arithmetic Hilbert space). The *arithmetic Hilbert space* is

$$\mathcal{H}_{\mathcal{A}} = \ell^2(\mathcal{F}) \cong \ell^2(\mathbb{N}_{\geq 1}), \quad (2)$$

with orthonormal basis $\{|k\rangle\}_{k \in \mathcal{F}}$, written $|n\rangle$ for $n = \prod_p p^{k_p}$ when no confusion arises. Each prime p defines a *mode* with occupation counter

$$\hat{n}_p |k\rangle = k_p |k\rangle, \quad (3)$$

a self-adjoint operator with spectrum $\mathbb{N}$.

Remark 2.2 (Why not an infinite tensor product?). A naive formulation writes $\mathcal{F}_{\text{Pr}} = \bigotimes_{p \in \text{Pr}} \mathcal{F}_p$ with one bosonic oscillator per prime. The literal von Neumann infinite tensor product of the $\ell^2(\mathbb{N})$ over a countably infinite index set is nonseparable, and its physically relevant separable subspaces depend on a choice of reference vector. Definition 2.1 achieves the intended object directly: ℓ^2 of the countable basis $\mathcal{F}$ is separable, and the tensor factorization over any *finite* set of modes is available whenever needed (Proposition 7.1). This is the first, and easiest, instance of the general principle of this paper: informal infinite constructions are replaced by the countable objects they were meant to describe.

For each prime p the standard bosonic ladder operators act on the p -mode: on basis vectors

$$a_p^\dagger |k\rangle = \sqrt{k_p + 1} |k + e_p\rangle, \quad a_p |k\rangle = \sqrt{k_p} |k - e_p\rangle, \quad (4)$$

where e_p is the unit occupation vector, and $[a_p, a_q^\dagger] = \delta_{pq}$ holds on the dense span of basis vectors. The total particle number is $\hat{N}_{\text{tot}} = \sum_p \hat{n}_p$, self-adjoint on $\{\psi : \sum_p k_p^2 |\psi_k|^2 < \infty\}$.

2.2 The multiplicative number operator and the primon Hamiltonian

The central operator of the framework is the *multiplicative number operator*, whose eigenvalue on $|k\rangle$ is the integer $n = \prod_p p^{k_p}$ itself. We define it together with the Hamiltonian it generates.

Definition 2.3 (Multiplicative number operator; primon Hamiltonian). The *multiplicative number operator* $\hat{N}$ is the diagonal operator

$$\hat{N} |k\rangle = \left(\prod_p p^{k_p} \right) |k\rangle, \quad (5)$$

self-adjoint on its maximal domain. The *primon Hamiltonian* is

$$H_P = \hbar \omega_0 \log \hat{N} = \hbar \omega_0 \sum_{p \in \text{Pr}} (\log p) \hat{n}_p, \quad \omega_0 > 0, \quad (6)$$

where the equality of the two expressions is pointwise on basis vectors and extends by closure.

Equation (6) exhibits H_P as a *free Bose gas* whose one-particle spectrum is $\{\log p : p \in \text{Pr}\}$: each prime is one species of elementary excitation (a *primon*), and a many-body state $|n\rangle$ has energy equal to the logarithm of the integer it represents. This is Julia’s primon gas [1], cleaned of its informal packaging.

Theorem 2.4 (Self-adjointness, spectrum, and simplicity). H_P is self-adjoint and nonnegative. Its spectrum is the pure point set

$$\text{spec}(H_P) = \overline{\{\hbar\omega_0 \log n : n \in \mathbb{N}_{\geq 1}\}}, \quad (7)$$

a discrete set whose only accumulation point is $+\infty$; it has no finite accumulation point. Every eigenvalue $\hbar\omega_0 \log n$ is simple: $\log m = \log n$ if and only if $m = n$. The ground state is the vacuum $|1\rangle$ with eigenvalue 0, and it is unique.

Proof. Under the unitary $U: \mathcal{H}_A \rightarrow \ell^2(\mathbb{N}_{\geq 1})$, $U|k\rangle = |n\rangle$ (unitary by the fundamental theorem of arithmetic), H_P becomes multiplication by the real sequence $a_n = \hbar\omega_0 \log n$. Multiplication by a real-valued sequence is self-adjoint on its maximal domain $\{c \in \ell^2 : \sum_n a_n^2 |c_n|^2 < \infty\}$, and its spectrum is the closure of $\{a_n\}$. Since $a_n \rightarrow \infty$ monotonically, the closure adds no finite points. Simplicity: if $a_m = a_n$ with $m \neq n$, then $\log m = \log n$ forces $m = n$, a contradiction. The ground state: $a_n \geq 0$ with equality only at $n = 1$. $\square$

Two consequences deserve emphasis. First, the *unique zero-entropy ground state* $|1\rangle$ is the multiplicative identity; this is the arithmetic floor underlying the Janus point developed in Sections 6 and 9. Second, the *simplicity* of every level means the primon spectrum has no degeneracy whatsoever: the integers label states one-to-one. Any degeneracy in a putative physical realization must therefore come from sectors *beyond* H_P (for instance the Gaussian sector of Section 8).

2.3 The partition function and the Hagedorn temperature

Proposition 2.5 (Partition function). For inverse temperature $\beta > 0$,

$$Z(\beta) = \text{Tr } e^{-\beta H_P} = \sum_{n=1}^{\infty} e^{-\beta \hbar\omega_0 \log n} = \sum_{n=1}^{\infty} n^{-\beta \hbar\omega_0} = \zeta(\beta \hbar\omega_0), \quad (8)$$

and the series converges if and only if $\beta \hbar\omega_0 > 1$, i.e. $T < T_H$, where

$$T_H = \frac{\hbar\omega_0}{k_B} \quad (9)$$

is the Hagedorn temperature of the primon gas. On the half-line $\sigma = \beta \hbar\omega_0 > 1$, the Euler product

$$\zeta(\sigma) = \prod_{p \in \text{Pr}} \frac{1}{1 - p^{-\sigma}} = \prod_{p \in \text{Pr}} \frac{1}{1 - e^{-\beta \varepsilon_p}}, \quad \varepsilon_p := \hbar\omega_0 \log p, \quad (10)$$

is precisely the grand-canonical factorization of a free Bose gas into independent prime modes, with each mode contributing the standard Bose factor. The mean occupation of mode p is

$$\langle \hat{n}_p \rangle_\beta = \frac{1}{e^{\beta \varepsilon_p} - 1} = \frac{1}{p^{\beta \hbar\omega_0} - 1}. \quad (11)$$

Proof. The trace formula is the spectral decomposition of Theorem 2.4; convergence at $\sigma > 1$ and the Euler product are Euler's theorems; the occupation formula is the standard free-boson computation on each mode. $\square$

The identification $Z = \zeta$ carries a consequence that gives it physical teeth: the primon gas is a *Hagedorn system* [10]. Its density of states grows exponentially in the energy, and the partition function diverges at a finite critical temperature. In string theory this is the famous signal that, near T_H , energy input goes into multiplying the number of light degrees of freedom rather than into heating. The same reading applies here with total exactness, because the counting is exactly solvable.

Theorem 2.6 (Hagedorn asymptotics). *Let $k_B = 1$ and write $T_H = \hbar\omega_0$. As $T \uparrow T_H$ (i.e. $\sigma = \beta\hbar\omega_0 \downarrow 1$):*

$$\zeta(\sigma) = \frac{1}{\sigma - 1} + \gamma + O(\sigma - 1), \quad (12)$$

$$U(T) := -\frac{\partial}{\partial\beta} \log Z(\beta) = \frac{\hbar\omega_0}{\sigma - 1} + O(\hbar\omega_0) = \frac{T T_H}{T_H - T} + O(T_H), \quad (13)$$

$$S(T) := \beta U + \log Z = \frac{1}{\sigma - 1} - \log(\sigma - 1) + O(1), \quad (14)$$

$$C(T) := \frac{dU}{dT} = \frac{T_H^2}{(T_H - T)^2} (1 + O((T_H - T)/T)). \quad (15)$$

In particular the specific heat diverges as $T \uparrow T_H$, while the entropy diverges only as $T/(T_H - T)$; energy is absorbed at nearly constant temperature.

Proof. The Laurent expansion $\zeta(\sigma) = 1/(\sigma - 1) + \gamma + O(\sigma - 1)$ is classical [11]. Then $-\zeta'/\zeta(\sigma) = [1/(\sigma - 1)^2 + O(1)]/[1/(\sigma - 1) + O(1)] = 1/(\sigma - 1) + O(1)$, giving $U = \hbar\omega_0(-\zeta'/\zeta)(\sigma)$. Substituting $\sigma = T_H/T$ gives $\sigma - 1 = (T_H - T)/T$ and the stated form. The entropy follows from $S = \beta U + \log Z$ with $\log Z = -\log(\sigma - 1) + O(1)$. Differentiating $U = T T_H/(T_H - T)$ in T gives $C = T_H^2/(T_H - T)^2$ up to the stated relative error. $\square$

Remark 2.7 (Microcanonical reading). Section 6 proves the exact counting law $\log N(E) = E/\hbar\omega_0 + O(1)$, which says that the *microcanonical* temperature $(dS/dE)^{-1}$ equals T_H identically, at every energy scale: the primon gas is an exactly critical system. It never heats; it only multiplies states. This is the sharpened form of the source's statement that “the state space expands exponentially”: the expansion rate is not merely large, it is *uniform* in E , and that uniformity is what the pole of ζ at $s = 1$ encodes. The classical smooth term “ x ” of the prime number theorem is the same statement read on the counting function $\psi(x)$, and the oscillatory remainder $-\sum_\rho x^\rho/\rho$ is the fluctuation sector; the paper returns to this duality in Sections 5 and 8.

2.4 Gaussian extension

The arithmetic sector extends to the Gaussian integers $\mathbb{Z}[i]$ for the putative five-dimensional (and chirality-carrying) sector, and here the mathematics is genuinely supportive, because the extension is an instance of the standard theory of number fields.

Definition 2.8 (Gaussian primon gas). Let $\mathcal{P}_{\mathbb{Z}[i]}$ denote the Gaussian primes modulo units $\{\pm 1, \pm i\}$, with norm $N(\pi) = |\pi|^2$. The *Gaussian arithmetic Hilbert space* is $\mathcal{H}_{\mathbb{Z}[i]} = \ell^2(\mathcal{F}_{\mathbb{Z}[i]})$ over the free commutative monoid on $\mathcal{P}_{\mathbb{Z}[i]}$, and the Gaussian primon Hamiltonian is

$$H_{\mathbb{Z}[i]} = \hbar\omega_0 \sum_{\pi \in \mathcal{P}_{\mathbb{Z}[i]}} \log N(\pi) \hat{n}_\pi, \quad Z_{\mathbb{Z}[i]}(\beta) = \prod_{\pi} \frac{1}{1 - N(\pi)^{-\beta\hbar\omega_0}}. \quad (16)$$

Theorem 2.9 (Dedekind factorization). $Z_{\mathbb{Z}[i]}(\beta) = \zeta_{\mathbb{Q}(i)}(\beta\hbar\omega_0)$, the Dedekind zeta function of the Gaussian rationals, and

$$\zeta_{\mathbb{Q}(i)}(s) = \zeta(s) L(s, \chi_{-4}), \quad (17)$$

where $L(s, \chi_{-4}) = \sum_m (-1)^{m-1} (2m-1)^{-s}$ is the Dirichlet L -function of the nontrivial character mod 4. Both factors converge for $\sigma > 1$; the product is entire on the common half-plane and $L(1, \chi_{-4}) = \pi/4 \neq 0$.

Proof. Unique factorization of ideals in $\mathbb{Z}[i]$ (it has class number 1) gives the Euler product over Gaussian prime ideals; identifying prime ideals with Gaussian primes modulo units gives the stated product, and the splitting classification of rational primes in $\mathbb{Z}[i]$ ($p \equiv 1 \pmod{4}$ splits, $p \equiv 3 \pmod{4}$ is inert, 2 ramifies) yields $\zeta_{\mathbb{Q}(i)} = \zeta L(\chi_{-4})$ [12]. The nonvanishing $L(1, \chi_{-4}) = \pi/4$ is Leibniz's series. $\square$

Equation (17) exhibits the Gaussian sector as the rational sector *times* a chiral factor: $L(s, \chi_{-4})$ is precisely the asymmetry between the splitting class $p \equiv 1 \pmod{4}$ and the inert class $p \equiv 3 \pmod{4}$. This is developed into a precise (and appropriately demoted) arithmetic model of charge conjugation and CP violation in Section 8.

2.5 What the arithmetic sector does and does not assert

It is worth fixing the epistemic boundary before dynamics is introduced. Everything in this section is unconditional mathematics: the primon gas exists as a quantum system, its spectrum is exactly the logarithms of the integers, its thermodynamics is exactly the analytic theory of ζ , and its critical behavior is exactly Hagedorn behavior. *No claim has yet been made that any physical system is a primon gas.* The identification $Z_{\text{grav}}^{\text{ren}}(s) = \zeta(s)$ for near-singularity gravity is Conjecture C1 of the register, the reported 2025–2026 developments [3] being its current physical evidence. The discipline of this paper is that such identifications live in the register, not in the theorems.

3 Dynamics: The Interaction Sector on the Prime Lattice

3.1 The frozen Hamiltonian

The dynamical claims of the framework—transitions into composite integer states, and factorization of the evolving state into highly composite integers—require a Hamiltonian with off-diagonal matrix elements in the occupation basis. The primon Hamiltonian has none:

Proposition 3.1 (Frozen spectrum). $[H_P, \hat{N}] = 0$ and $[H_P, \hat{n}_p] = 0$ for every p . Consequently every basis vector $|n\rangle$ is a stationary state: $e^{-i\tau H_P/\hbar}|n\rangle = e^{-i\omega_0\tau \log n}|n\rangle$ never leaves the ray $\mathbb{C}|n\rangle$. In particular the vacuum $|1\rangle$ is an eigenstate with eigenvalue 0 and evolves trivially, $e^{-i\tau H_P/\hbar}|1\rangle = |1\rangle$ for all τ .

Proof. Both operators are diagonal in the same basis (Theorem 2.4). $\square$

A diagonal Hamiltonian cannot create primons, destroy them, or entangle anything. The “arithmetic cascade,” the entanglement growth, and the emergence of a thermodynamic arrow all require dynamics *off* the diagonal of $\hat{N}$. The interaction sector must be supplied. Meeting this requirement produces, as a byproduct, the objects on which the rest of the framework rests: the multiplicative shift operators, the prime lattice, and a well-posed thermalization problem.

3.2 Multiplicative shift operators

Definition 3.2 (Multiplication operator). For $n \in \mathbb{N}_{\geq 1}$ define $M_n: \mathcal{H}_{\mathcal{A}} \rightarrow \mathcal{H}_{\mathcal{A}}$ on basis vectors by

$$M_n|k\rangle = |nk\rangle, \quad (18)$$

extended linearly, with adjoint $M_n^\dagger|k\rangle = |k/n\rangle$ if $n \mid k$ (i.e. $k_p \geq a_p$ for all p with $n = \prod p^{a_p}$) and $M_n^\dagger|k\rangle = 0$ otherwise.

Proposition 3.3 (Monoid representation). *The assignment $n \mapsto M_n$ is an isometric representation of the multiplicative monoid $(\mathbb{N}_{\geq 1}, \times)$ on $\mathcal{H}_{\mathcal{A}}$:*

1. $M_m M_n = M_{mn} = M_n M_m$ (the representation is abelian);
2. $M_n^\dagger M_n = I$ (each M_n is an isometry);
3. $M_n M_n^\dagger = I - P(n \mid k)$, where $P(n \mid k)$ is the projection onto $\overline{\text{span}}\{|k\rangle : n \mid k\}$, a product over the modes $p^{a_p} \parallel n$ of the projections onto $\{\hat{n}_p \geq a_p\}$;

4. $\|M_n\| = \|M_n^\dagger\| = 1$, and $M_1 = I$.

Moreover, for each prime p ,

$$M_p = a_p^\dagger (\hat{n}_p + 1)^{-1/2}, \quad (19)$$

so M_p is the unilateral shift of the p -mode tensored with the identity on all other modes.

Proof. (1) $M_m M_n |k\rangle = |mnk\rangle$ by associativity and commutativity of multiplication in $\mathbb{N}$. (2) M_n maps the orthonormal basis $\{|k\rangle\}$ to the orthonormal family $\{|nk\rangle\}$ (distinct k give distinct nk by cancellation in $\mathbb{N}$), hence is an isometry. (3) The initial space of $M_n^\dagger$ is the range of M_n , i.e. the closed span of $|k\rangle$ with $n \mid k$; this condition factorizes across modes because divisibility does. (4) is immediate. For (19): on $|k\rangle$, $a_p^\dagger (\hat{n}_p + 1)^{-1/2} |k\rangle = \sqrt{k_p + 1} (k_p + 1)^{-1/2} |k + e_p\rangle = |k + e_p\rangle = M_p |k\rangle$. $\square$

The operators M_n are the arithmetic analogues of the Weyl shifts, and Proposition 3.3 is the exact statement behind the phrase that composites are “correlated multi-prime states”: multiplication *creates* occupation, division *annihilates* it, and the two are adjoints. Note also the co-isometric defect $M_n M_n^\dagger = I - P(n \mid k)$: multiplication by n is invertible precisely on states already divisible by n . The failure of unitarity of each individual M_n is the operator-theoretic fingerprint of the irreversibility that arithmetic carries: the semigroup $(\mathbb{N}, \times)$ is free, hence has cancellation but no inverses, and the quantum representation inherits exactly that structure.

3.3 The walk Hamiltonian and the prime lattice

Define, for a finite set $P_f \subset \text{Pr}$ of active modes and real couplings $g = (g_p)_{p \in P_f}$, the *multiplicative walk Hamiltonian*

$$H_W(P_f, g) = \sum_{p \in P_f} g_p (M_p + M_p^\dagger), \quad (20)$$

and the *hopping Hamiltonian*

$$H_{\text{hop}}(P_f, h) = \sum_{p \neq q \in P_f} h_{pq} (a_p^\dagger a_q + a_q^\dagger a_p). \quad (21)$$

Proposition 3.4 (Well-posed interacting dynamics). *For finite P_f , H_W and H_{hop} are bounded and self-adjoint. The full Hamiltonian*

$$H_{\text{tot}} = H_P + H_W + H_{\text{hop}} \quad (22)$$

is self-adjoint on the domain of H_P (Kato–Rellich), and $[H_{\text{tot}}, \hat{N}] \neq 0$: the dynamics generates genuine transitions

$$\langle m | H_W | k \rangle \neq 0 \iff m/k \in \{p, p^{-1}\} \text{ for some } p \in P_f, \quad (23)$$

i.e. precisely the edges of the divisibility graph of $\mathbb{N}$ restricted to P_f .

Proof. Each M_p is bounded with norm 1 (Proposition 3.3); a finite sum of bounded self-adjoint operators is bounded self-adjoint. $a_p^\dagger a_q$ is bounded (particle-number preserving hop). H_{tot} is a bounded perturbation of the self-adjoint H_P , hence self-adjoint on $D(H_P)$. Non-commutation: M_p changes the eigenvalue of $\hat{N}$ from n to np , and a direct computation gives

$$[H_W, \hat{N}] |n\rangle = \sum_p g_p n \left(-(p-1) |np\rangle + \left(1 - \frac{1}{p}\right) \mathbf{1}_{p|n} |n/p\rangle \right) \neq 0 \quad (24)$$

for $n \geq 1$ and nonzero g : the two coefficients are different in both sign and magnitude ($-(p-1)$ versus $1 - 1/p$), so the terms cannot cancel. $\square$

Three remarks interpret these objects.

Remark 3.5 (The prime lattice). H_{hop} is a Bose–Hubbard-type tight-binding Hamiltonian on a lattice whose *sites* are the primes, with on-site energies $\varepsilon_p = \hbar\omega_0 \log p$ and hopping h_{pq} . We call this graph the *prime lattice*. It is the natural home for every interaction claim in the framework: interactions among prime modes are hops along the lattice; the chaotic sector conjectured in Conjecture C5 is a spectral-statistics statement about H_{tot} on the prime lattice; and the “chemistry of the primes” becomes the statement that the Standard Model’s force sectors correspond to structured sub-lattices (Conjecture C9).

Remark 3.6 (Conservation structure). The three sectors carry a strict conservation hierarchy. H_P is diagonal in the occupation basis and conserves everything. H_{hop} moves single primons between modes ($a_p^\dagger a_q$: one occupation up, one down), so it conserves the *total primon number* $\hat{N}_{\text{tot}} = \sum_p \hat{n}_p$ exactly, but not $\hat{N}$ (it rescales n by q/p). H_W conserves *neither*: each M_p raises the occupation of the p -mode by one and hence raises $\hat{N}_{\text{tot}}$ by one, while each $M_p^\dagger$ lowers it—the walk is the nucleation sector, the engine of entropy growth in the precise quantitative sense of Proposition 3.10 ($d\langle \hat{N}_{\text{tot}} \rangle / d\tau > 0$ from [1] whenever $H_W \neq 0$). The full theory therefore has a graded conservation hierarchy: H_P conserves $\hat{N}$ and $\hat{N}_{\text{tot}}$; H_{hop} conserves $\hat{N}_{\text{tot}}$ only; H_W conserves neither and is the only source of net primon creation. The additive and multiplicative structures of $\mathbb{N}$ thereby acquire distinct dynamical roles—spectral order (H_P), transport (H_{hop}), creation (H_W)—which separates “spectral order” from “arithmetic cascade.” Register note: the CP ratchet of C10 lives entirely in this non-conserving H_W sector.

Remark 3.7 (Relation to the reported physics). The conformal primon gas reported near black-hole singularities [3] corresponds in this language to the claim that *some* sector of near-singularity gravity is governed by H_P (or its Gaussian analogue) with the conformal $\mathfrak{sl}(2, \mathbb{R})$ symmetry of the axioms acting on the ω_0 scale. The interacting extension (22) is the framework’s addition; its necessity follows from the arrow-of-time claims, which (by Proposition 3.1) are impossible under H_P alone.

3.4 The cascade: statics and dynamics

The “thermodynamic cascade” is the framework’s central image from sparse prime states into dense composite states. As dynamics under H_P this is impossible (Proposition 3.1); as a statement about the *microcanonical ensemble*, however, it is a classical theorem. The dichotomy therefore splits the cascade into a theorem (statics) and a conjecture (dynamics).

Theorem 3.8 (Normal order and Erdős–Kac for the primon ensemble). *Fix $E > 0$ and let $x = e^{E/\hbar\omega_0}$. Equip $\{n \leq x\}$ with the uniform measure, i.e. the microcanonical ensemble of H_P at energy E . Let $\omega(n)$ be the number of distinct prime factors of n . Then:*

1. (Hardy–Ramanujan [13]; Turán [14].) $\omega(n)$ has normal order $\log \log n$: for every $\varepsilon > 0$ the proportion of $n \leq x$ with $|\omega(n) - \log \log x| > \varepsilon \log \log x$ tends to 0 as $x \rightarrow \infty$.
2. (Erdős–Kac [15].) The distribution of $(\omega(n) - \log \log x) / \sqrt{\log \log x}$ converges to the standard Gaussian: for all t ,

$$\lim_{x \rightarrow \infty} \frac{1}{x} \# \left\{ n \leq x : \frac{\omega(n) - \log \log x}{\sqrt{\log \log x}} \leq t \right\} = \Phi(t). \quad (25)$$

The interpretation is exact. The number of states at energy E is $\lfloor e^{E/\hbar\omega_0} \rfloor$ (Section 6), and Theorem 3.8 describes the *typical* member of that ensemble: a composite integer with about $\log \log x$ distinct prime factors, $\log \log x = \log(E/\hbar\omega_0)$ growing only logarithmically in the energy. The entropy $\log N(E) = E/\hbar\omega_0 + O(1)$ is astronomically larger than the typical complexity $\log \log N(E)$: the ensemble is dominated by integers that are composite but *far from*

maximally composite (the maximally composite integers, the powers of two and their smooth kin, are exponentially atypical). Thus a “cascade into composite states” is correct as a description of where the measure lives, and the Erdős–Kac Gaussian is the precise fluctuation law of the cascade. What remains open is the dynamical version:

Conjecture 3.1 (Cascade/thermalization on the prime lattice, interacting completion). *Let*

$$H_{\text{tot}}^{(U,V)} = H_P + H_W + H_{\text{hop}} + U \sum_{p < q} \hat{n}_p \hat{n}_q + V \sum_{p < q} (\hat{n}_p + \hat{n}_q) (a_p^\dagger a_q + a_q^\dagger a_p) \quad (26)$$

on $\mathcal{H}_A$ with generic couplings and $U \neq 0$ or $V \neq 0$ (diagonal or kinetic interacting completion; both are quartic in the ladder operators, and the kinetic one is off-diagonal and $\hat{N}_{\text{tot}}$ -preserving, with matrix elements $V(k_p + k_q)\sqrt{(k_p + 1)k_q}$ on the hops of the prime lattice), restricted to the energy window $[E, E + \Delta]$. Then along any truncation sequence with truncation dimension $D \rightarrow \infty$ at fixed effective dose: (i) eigenstate expectation values of local occupation observables are smooth in E with within-window fluctuations $\sigma_{\text{ETH}}(D) \rightarrow 0$; (ii) diagonal-ensemble values converge to the microcanonical values of Theorem 3.8; and (iii) the relaxation time scale is polynomial in E . The bare H_{tot} ($U = V = 0$) is excluded: it is a quasi-quadratic object (see below) to which the thermalization question does not apply.

Conjecture 3.1 (register entry C4) is the engine for every arrow-of-time claim in the framework: entropy growth, decoherence, record formation, and the emergence of the second law are all downstream of it. It is a well-posed, hard, and conventional-sounding problem of quantum statistical mechanics, which is exactly the point: the framework’s most sweeping physical claims reduce to questions that experts know how to test. The two completions in (26) are not equivalent and the difference matters: the U term is a diagonal potential on the exponent lattice (a Wannier–Stark-type renderer, as the data confirm), while the V term is the Bose–Hubbard-type *density-assisted hopping* known from the interacting-boson literature—quartic yet kinetic. The two are distinguished by computation:

Remark 3.9 (Numerical status of the cascade conjecture). Conjecture 3.1 has been tested by dense diagonalization at $D \leq 4096$; by shift-invert Lanczos eigen-windows ($k = 350$ interior eigenpairs placed at the median of the density of states, diagonal pivoting in the factorization, every window certified by eigenpair residuals $\leq 6 \times 10^{-8}$) up to $D = 59319$ in $d = 3$ and $D = 20736$ in $d = 4$, beyond which the LU fill exceeds the 4 GB address-space guard; and by restarted-Lanczos Krylov equilibration to $D \approx 10^5$, with restart steps sized by a Lanczos-estimated spectral radius (validated against exact eigendecomposition propagation to 10^{-12} , state-vector norm drift $\leq 10^{-13}$). The kinetic family is examined at matched effective dose $VK^2 \approx 36$, the occupation-weighted coupling, so that the dose is fixed while the truncation widens; code and data accompany the paper. The evidence: (a) *Level statistics*: generic couplings give GOE ratio statistics $\langle r \rangle = 0.45\text{--}0.58$ at every scale computed from $D = 625$ to 5.9×10^4 (the upper end of the band is the $d = 4$, $K = 11$, $D = 20736$ window; the sub-GOE values sit at the smallest matched-dose grids, $K \leq 14$, where the fixed effective dose leaves spectral mixing incomplete); the weak-coupling control collapses the participation ratio to $\text{PR}/D \approx 10^{-4}$ with σ_{ETH} inflated by an order of magnitude, and its $\langle r \rangle$ is *window-placement-dependent in a way the generic regime is not*: a σ -quantile sweep of the same weak configuration ($d = 3$, $K = 28$, $D = 24389$) at quantiles $\{0.02, 0.10, 0.15, 0.25, 0.5, 0.75, 0.9, 0.98\}$ gives $0.42\text{--}0.56$, non-monotone (the dip at the 15% quantile, 0.436 , is reproduced exactly in an independent repetition—density-of-states cluster structure, not solver noise), while the generic control is flat ($0.505\text{--}0.514$ at quantiles $0.05/0.5/0.95$); the weak median value is additionally direction-dependent (0.53 at $d = 3$ versus 0.44 at $d = 4$, $D = 20736$). Any protocol claim about weak-coupling statistics must therefore pin the quantile and the direction; the robust weak-coupling diagnostic is PR/D , not $\langle r \rangle$. (b) *ETH fails for the bare H_{tot} structurally*: H_P is a separable ramp, H_W is a per-axis adjacency (Wannier–Stark chains; propagation along the

p -axis is Stark-suppressed when $g_p \lesssim \log p$), and H_{hop} is quadratic—no quartic term exists anywhere in H_{tot} , so the system is essentially single-particle on the exponent lattice $\mathbb{Z}^d$. The scaled data make this quantitative: the normalized eigenstate fluctuation $\sigma_{\text{ETH}}/\text{std}(a_j)$ of local occupations sits at 0.94–0.99—fluctuations as large as the observable’s full spread—flat in D at every coupling and every fixed dose (item (e) below); and the time-averaged occupation retains 6–24% of the initial mode’s occupation ($|\text{diag} - \text{micro}|/K$) at every scale from $D = 10^4$ to 10^5 , with no downward trend. (c) *The diagonal quartic completion does not rescue thermalization at any accessible scale*: the dose-response is monotone localization ($\langle r \rangle$: 0.514 at $U = 0$, 0.502 at $U = 0.5$, 0.387 at $U = 2$, all at $D \approx 2 \times 10^4$; PR/D : $0.18 \rightarrow 0.03 \rightarrow 0.001$; frozen-mode memory $|\text{diag} - \text{micro}|/K \approx 0.8$). (d) *The kinetic (density-assisted hopping) completion*—a one-term addition to the Hamiltonian, Hermitian and exactly $\hat{N}_{\text{tot}}$ -preserving, validated against a first-principles dense assembly to 10^{-15} —is the only completion that relaxes. At matched dose $VK^2 \approx 36$ on the $d = 4$, $K = 11$, $D = 20736$ platform, the occupation $\hat{n}_d$ falls from its initial value 11 to ≈ 3.5 within $\tau \approx 20$ and then drifts on a slow secular tail; the eigenstates stay delocalized with GOE statistics ($\text{PR}/D = 0.14$, $\langle r \rangle = 0.51$) where the quartic completion collapses to 0.001–0.03. The equilibration is a dose phenomenon, not a monotone onset: at $V = 0.1$ the plateau sits *away* from microcanonical ($|\text{diag} - \text{micro}|/K = 0.18$, the bare value; the short-horizon averages at the same platform are 0.10 bare and 0.81 quartic), and strong dose localizes, as the occupation-weighted matrix elements predict (effective coupling VK^2 : at $V = 1$, $\langle r \rangle \rightarrow 0.41$ and $\text{PR}/D \rightarrow 0.05$; at fixed $V = 0.3$ on the $d = 3$, $K = 28$ grid, where $VK^2 \approx 235$, $\langle r \rangle = 0.36$ with $\text{PR}/D = 0.016$). (e) *Scaling at matched dose; the diagonal ensemble*. Two legs decide the conjecture. The eigenstate fluctuation ratio $\sigma_{\text{ETH}}/\text{std}(a_j)$ is flat: 0.95–0.98 along the $d = 3$ ladder $D = 3375 \rightarrow 59319$ (a factor 17.6 in D) and 0.95–0.97 along the $d = 4$ ladder $D = 4096 \rightarrow 20736$, with exact full-spectrum values 0.68–0.83 at $D \leq 10^4$; leg (i) has no positive evidence at accessible scales. The diagonal ensemble $\langle a \rangle_{\text{diag}} = \sum_n |\langle n | \psi_0 \rangle|^2 \langle n | a \rangle$ is computed exactly at $D \leq 10^4$: there the finite-time plateau equals it to $|\text{plateau} - \text{diag}|/K \leq 0.005$ (the dephasing identity $\lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T \langle a(t) \rangle dt = \langle a \rangle_{\text{diag}}$ holds with no ETH input, and nested late-time window averages supply the convergence diagnostic). But at matched dose the diagonal ensemble is not microcanonical: $|\langle a \rangle_{\text{diag}} - \langle a \rangle_{\text{micro}}|/K = 0.040 \rightarrow 0.061 \rightarrow 0.092$ along $d = 4$ ($D = 4096 \rightarrow 20736$) and ≈ 0.05 (stable in absolute value, ≈ 0.87) along $d = 3$ ($D = 3375 \rightarrow 9261$), with no closing trend. The $d = 4$, $D = 20736$ trajectory was extended to $\tau = 300$ in checkpointed wall-clock chunks (a chunk boundary is an ordinary Lanczos restart; the resumed trajectory agrees with an independent step-size choice to 2×10^{-4} in the observable): its nested late-time averages saturate at $\langle \hat{n}_d \rangle_{\text{diag}} \approx 4.40 \pm 0.03$, while the microcanonical value is 3.35 and the early transient ($\tau \leq 22$) passes through 3.39, close to it. A finite-horizon coincidence with the microcanonical value is therefore not evidence of thermalization; leg (ii) is a statement about the $D \rightarrow \infty$ limit of the diagonal ensemble, and at accessible scales that limit shows no convergence (Figure 1). (The kinetic couplings destroy the diagonal dominance on which default partial pivoting relies in the shift-invert factorization; diagonal pivoting (`diag_pivot_thresh` = 0) restores tractability—the generic $d = 3$, $K = 28$ factorization drops from 257 s to 3 s—with every eigen-window certified by eigenpair residuals $\leq 6 \times 10^{-8}$ and a pivoting cross-check at $D = 14641$ agreeing to 0.002 in $\langle r \rangle$.) An $E \rightarrow \infty$ statement cannot be falsified by any finite- D simulation; a *scaling* criterion can be—and on the present evidence every scaling diagnostic is flat.

Falsifiers and test routes are listed in Section 10.

3.5 The Janus state

A natural boundary condition is the vacuum: the claim that “at the past hypothesis the phase $1^{-i\omega_0\tau} = 1$: time does not exist at $n = 1$; time turns on when the universe transitions to composite states.” Under H_P *nothing* ever transitions (Proposition 3.1), so the claim would be empty; under H_{tot} the vacuum $|1\rangle$ is *not* an eigenstate (because $H_W|1\rangle = \sum_p g_p|p\rangle$), and the statement becomes substantive and true:




Figure 1: Matched-dose ($VK^2 \approx 36$) numerics for the kinetic completion (Remark 3.9(d)–(e)). (a) Eigenstate fluctuation ratio $\sigma_{\text{ETH}}/\text{std}(a)$ versus truncation dimension: window values (filled) and exact full-spectrum values (open); no decay across a factor 17.6 in D . (b) Distance of the diagonal ensemble from the microcanonical value, normalized by K : $O(1)$ gaps with no closing trend; the $T \leq 22.5$ estimate (cross) is a transient that coincides with the microcanonical value. (c) The $d = 4, K = 11, D = 20736, V = 0.3$ trajectory extended to $\tau = 300$: observable, running time average, nested late-time averages (diagonal-ensemble estimate ≈ 4.40), microcanonical line, and the early transient. (d) Finite- T plateau versus exact diagonal ensemble at $D \leq 10^4$ (identity line).

Proposition 3.10 (Nucleation from the Janus state). *For H_{tot} with $H_W \neq 0$ and initial state $\psi(0) = |1\rangle$:*

$$\psi(\tau) = |1\rangle - \frac{i\tau}{\hbar} \sum_{p \in P_f} g_p |p\rangle + O(\tau^2), \quad (27)$$

so the total primon number grows initially at rate $d\langle \hat{N}_{\text{tot}} \rangle / d\tau = 2 \text{Im} \langle \psi | H_W | \psi \rangle / \hbar + O(\tau) = \frac{2\tau}{\hbar^2} \sum_p g_p^2 + O(\tau^2) > 0$. The unique zero-entropy state $|1\rangle$ is a regular point of the dynamics, not a fixed point; under H_P alone it is a fixed point (Proposition 3.1).

Proof. First-order time-dependent perturbation theory around the eigenstate $|1\rangle$ of H_P (the Dyson series in $H_W + H_{\text{hop}}$; H_{hop} annihilates the vacuum). $\square$

Time, in this framework, is switched on by the interaction sector, with a mechanism that exists. The stronger claims that the actual history of the universe *begins* at $|1\rangle$, and that the entropy profile is two-sided (Janus-shaped) about that event, are boundary-condition statements and are demoted to Conjectures C7 and C8; what remains as theorem is only the uniqueness and zero entropy of the floor (Theorem 2.4 and Section 6).

4 Recurrence, Periodicity, and the Fate of Eternal Return

The anti-cyclical verdict—that prime arithmetic rigorously forbids exact cyclical time and the universe cannot repeat—decomposes into a theorem, a correct application, and two qualifications. This section separates them. The resulting statement is *stronger* than the slogan, because it identifies exactly which recurrence phenomena are forbidden, which are unavoidable, and what the boundary between them depends on.

4.1 Linear independence of the prime logarithms

Lemma 4.1 (Fundamental-theorem independence). *The set $\{\log p : p \in \text{Pr}\}$ is $\mathbb{Q}$ -linearly independent. In particular, for distinct primes p, q , the ratio $\log p / \log q$ is irrational.*

Proof. Suppose $\sum_{i=1}^r q_i \log p_i = 0$ with $q_i = a_i/b_i \in \mathbb{Q}$ and distinct primes p_i . Let $B = \text{lcm}(b_i)$ and $c_i = Bq_i \in \mathbb{Z}$. Then $0 = \sum_i c_i \log p_i = \log \prod_i p_i^{c_i}$ (negative exponents handled by clearing signs into the other side), so $\prod_i p_i^{c_i} = 1$. Unique factorization forces $c_i = 0$ for all i , hence $q_i = 0$. $\square$

4.2 Exact periodicity is forbidden

Theorem 4.2 (No exact return for generic states). *Let $\psi = \sum_n c_n |n\rangle$ and let $S(\psi) = \{n : c_n \neq 0\}$ be its support. The following are equivalent:*

1. *There exists $\tau > 0$ with $e^{-i\tau H_P/\hbar} \psi = \psi$;*
2. *There exists $T > 0$ such that $\omega_0 T \log n \in 2\pi\mathbb{Z}$ for every $n \in S(\psi)$;*
3. *The set $\{\log n : n \in S(\psi)\}$ is contained in $\alpha\mathbb{Z}$ for some $\alpha > 0$ (all supports are integer multiples of a common logarithmic unit; equivalently all $n \in S(\psi)$ are powers of a single integer).*

In particular, if $S(\psi)$ contains any two integers m, n that are not powers of a common integer—for example any pair among $\{2, 3\}$, $\{6, 10\}$, $\{2, 6\}$ —then ψ is aperiodic: $e^{-i\tau H_P/\hbar} \psi \neq \psi$ for every $\tau > 0$. The same holds for H_{tot} whenever its spectrum is simple and contains two levels whose difference is irrational in units of $\hbar/\omega_0$, which holds generically.

Proof. $e^{-i\tau H_P/\hbar} \psi = \sum_n c_n e^{-i\omega_0 \tau \log n} |n\rangle$ equals ψ iff $e^{-i\omega_0 \tau \log n} = 1$ for all $n \in S(\psi)$, i.e. iff $\omega_0 \tau \log n \in 2\pi\mathbb{Z}$ for all such n . Setting $\alpha = 2\pi/(\omega_0 \tau)$, this says every $\log n$ in the support lies in $\alpha\mathbb{Z}$. If $m, n \in S$ and $\log m, \log n \in \alpha\mathbb{Z}$ with $\log m / \log n \notin \mathbb{Q}$, the condition fails for every τ ; by Lemma 4.1, $\log 2 / \log 3 \notin \mathbb{Q}$ etc. $\square$

This is the rigorous core of the anti-cyclic verdict: *eternal return in the exact sense requires the support of the universal state to lie in a single geometric progression of integers*, and no physically motivated state does.

4.3 What is *not* forbidden: three qualifications

Corollary 4.3 (Truncated systems recur). *Fix a finite truncation: primes $p \leq X$ and occupations $k_p \leq K$, so the Hilbert space is finite-dimensional and the spectrum $\{E_j\}$ of H_P (or of any self-adjoint extension) is finite. Then the evolution $U(\tau) = e^{-i\tau H/\hbar}$ is a quasi-periodic rotation, and for every state ψ and every $\varepsilon > 0$ the set $\{\tau > 0 : \|U(\tau)\psi - \psi\| < \varepsilon\}$ is relatively dense (there exists $L = L(\varepsilon, \psi)$ such that every interval of length L contains such a τ). In particular Poincaré recurrence holds in its strongest (uniform, Bohr almost-periodic) form. This is the finite-dimensional special case of Theorem 4.4 below, recorded separately because it is the form in which recurrence appears in any computation.*

Proof. On a finite-dimensional space $U(\tau) = e^{-i\tau\Lambda}$ with $\Lambda = \text{diag}(E_1, \dots, E_d)/\hbar$; the map $\tau \mapsto (e^{-i\tau E_1/\hbar}, \dots, e^{-i\tau E_d/\hbar})$ is a continuous one-parameter subgroup of the compact torus $\mathbb{T}^d$, and every orbit of a compact group rotation is uniformly recurrent: the return times to any neighborhood of the identity are syndetic. This is Kronecker's theorem in its group form [16]. $\square$

Theorem 4.4 (Uniform recurrence of the free flow, infinite system). *For every state $\psi \in \mathcal{H}_A$ and every $\varepsilon > 0$, the set of ε -return times*

$$\left\{ \tau > 0 : \|e^{-i\tau H_P/\hbar}\psi - \psi\| < \varepsilon \right\} \quad (28)$$

is relatively dense in $\mathbb{R}$. The free arithmetic flow is Bohr almost periodic for all states, with no finite invariant measure required and no truncation assumed. What fails is not ε -recurrence but its uniformity in the state: the return-time scale $L(\varepsilon, \psi)$ is unbounded as a function of ψ (it grows with the Diophantine complexity of the finite supports approximating ψ), and exact return remains forbidden by Theorem 4.2.

Proof. Fix $\varepsilon > 0$ and $\psi = \sum_n c_n |n\rangle \in \ell^2$. Choose a finite support set $F \subset \mathbb{N}_{\geq 1}$ with $\sum_{n \notin F} |c_n|^2 < \varepsilon^2/4$. The truncated flow $\tau \mapsto (e^{-i\omega_0 \tau \log n})_{n \in F}$ is a continuous one-parameter subgroup $h: \mathbb{R} \rightarrow \mathbb{T}^F$; let $H = \overline{h(\mathbb{R})}$, a compact connected subgroup of the torus. The $\mathbb{R}$ -orbit of the identity under left translation by $h(\tau)$ is dense in H by construction, and a dense orbit of an equicontinuous (group) action on a compact space is minimal; minimal equicontinuous flows are uniformly recurrent, so $T(\delta) = \{\tau : d(h(\tau), e_H) < \delta\}$ is relatively dense for every $\delta > 0$. Taking δ small enough that all phases on F deviate by less than $\varepsilon/2$ in the ℓ^2 sense (finitely many constraints), we obtain $\|U(\tau)\psi - \psi\| \leq \varepsilon/2 + \varepsilon/2$ for every $\tau \in T(\delta)$, a relatively dense set. The non-uniformity in ψ follows because F must grow as the tails of ψ shrink, and the Diophantine return times on F are unbounded in $|F|$ (for supports with rationally independent generators, Kronecker's theorem gives relative density but no uniform bound: the minimal return time for $F = \{2, 3, \dots, p_{|F|}\}$ grows at least exponentially in $|F|$, since the frequencies $\log n$ are rationally independent by Lemma 4.1 and simultaneous approximation of independent frequencies is slow). $\square$

The third qualification concerns the *classical* multiplicative walk, and it sharpens the boundary of the no-cycle verdict in an unexpected direction.

Proposition 4.5 (Pólya transience of the multiplicative walk). *Consider the classical random walk on the divisibility graph of Definition 3.2 restricted to $d = |P_f|$ active primes: from $k = (k_1, \dots, k_d) \in \mathbb{Z}_{\geq 0}^d$ step to $k + e_j$ with probability $1/(2d)$ and to $k - e_j$ (when $k_j > 0$) with probability $1/(2d)$, staying otherwise. Then the walk is recurrent for $d \leq 2$ and transient for $d \geq 3$ [17]. In particular for $d \geq 3$ the walk returns to the vacuum $k = 0$ (the Janus state) only finitely many times, almost surely.*

Proposition 4.5 has a striking interpretive reading: whether the arithmetic universe “cycles back to its origin” depends on the *number of active prime modes*. With one or two primes, the multiplicative universe is recurrent—a rigorous echo of cyclical cosmology. With three or more, it is transient, and the Janus point is a genuinely singular origin. The absolute prohibition thus becomes a structural, checkable dichotomy.

4.4 The verdict on cyclical time

Assembling the three legs: exact eternal return is impossible for states whose support spans incommensurable integers (Theorem 4.2); ε -approximate recurrence is universal for the *free* flow, at every state, with return times relatively dense but astronomically non-uniform (Theorem 4.4, and Corollary 4.3 for the trivial finite case); and the *cascading* (or classical walk) dynamics is transient for $d \geq 3$ modes (Proposition 4.5), with the interacting quantum system conjecturally

non-recurrent on every physical scale (Conjecture C4). The philosophical consequences are drawn in Section 9; the essential point is that the framework’s anti-cyclical verdict survives as a *layered* statement—exact return forbidden as a genuine theorem, practical return driven out of reach by Diophantine slowness, transience of the cascade for three or more active modes—rather than as an absolute metaphysical prohibition. The free arithmetic sector, left to itself, is the *most* recurrent object in the theory (Bohr almost periodicity for all states, Theorem 4.4); all irreversibility is therefore manufactured by the interaction sector and the boundary condition, never by the spectrum. This is the sharpest available form of the intuition that “the river never circles back,” and it is also the form most useful to the arrow-of-time analysis of Sections 5 and 9.

5 The Arithmetic Clock and the Problem of Time

5.1 The problem of time, stated precisely

The reconciliation problem for the two classical concepts of time is standard: in quantum mechanics, t is an external parameter and the Schrödinger equation evolves states with respect to it; in general relativity, elapsed time is the proper time along a worldline, $\tau_\gamma = c^{-1} \int_\gamma \sqrt{-g_{\mu\nu} dx^\mu dx^\nu}$, a geometric quantity that varies from trajectory to trajectory, while the canonical Hamiltonian of gravity is a constraint, $\hat{\mathcal{H}}\Psi = 0$, with no external evolution at all. The resolution is relational: time is a correlation between subsystems, realized quantum mechanically by the Page–Wootters mechanism [18], with the primon spectrum supplying the clock. This section turns that resolution into theorems, corrects one sign-of-life issue (the clock kernel never decorrelates, so the arrow of time cannot live in the clock), and quantifies the claim that “at the Janus point time does not exist.”

The Page–Wootters construction is standard: for a total state Ψ satisfying a constraint $(H_C + H_S)\Psi = 0$ of clock-plus-system, the conditional state of the system when the clock reads τ is $\psi_S(\tau) = \langle \tau | \Psi \rangle$, and for an ideal clock family this satisfies an effective Schrödinger equation in τ . All content therefore concentrates in the clock family. This paper’s contribution is the exact analysis of the arithmetic clock.

5.2 The zeta clock

Definition 5.1 (Zeta clock family). For $\beta > \frac{1}{2}$ define the normalized clock states

$$|\tau; \beta\rangle = \zeta(2\beta)^{-1/2} \sum_{n=1}^{\infty} n^{-\beta - i\omega_0\tau} |n\rangle, \quad (29)$$

a unit-norm family in $\mathcal{H}_A$. (The parameter β is a resolution regulator: the weights $n^{-\beta}$ confine the clock to energies of order $\hbar\omega_0/(2\beta - 1)$, as computed in Proposition 5.4.)

Proposition 5.2 (Clock covariance and two-time kernel). 1. (Covariance.) $e^{-i\tau' H_P/\hbar} |\tau; \beta\rangle = |\tau + \tau'; \beta\rangle$: clock translations are generated by H_P itself, so the clock reading τ is exactly the phase parameter conjugate to $\log \hat{N}$,

$$e^{-i\omega_0\tau \log \hat{N}} |n\rangle = n^{-i\omega_0\tau} |n\rangle. \quad (30)$$

2. (Two-time kernel.) For all τ, τ' ,

$$\langle \tau'; \beta | \tau; \beta \rangle = \frac{\zeta(2\beta - i\omega_0(\tau - \tau'))}{\zeta(2\beta)}. \quad (31)$$

3. (Positivity.) The kernel (31) is positive definite on $\mathbb{R}$: it is the Fourier transform of the probability measure $\mu_\beta = \zeta(2\beta)^{-1} \sum_n n^{-2\beta} \delta_{\log n}$ on the energy-per-mode variable $u = \log n$.

4. (Product of phases.) For a superposition $\sum_n c_n |n\rangle$ evolved for time τ , the phase acquired is $n^{-i\omega_0\tau}$, and by unique factorization

$$n^{-i\omega_0\tau} = \prod_p p^{-i\omega_0\tau k_p} : \quad (32)$$

each prime contributes an elementary phase generator, and composite evolution is the product of prime evolutions. This is the exact sense in which “primes provide the elementary generators of spectral phase order.”

Proof. (1) is immediate term by term. (2) $\langle \tau'; \beta | \tau; \beta \rangle = \zeta(2\beta)^{-1} \sum_n n^{-2\beta} e^{i\omega_0(\tau-\tau') \log n} = \zeta(2\beta)^{-1} \zeta(2\beta - i\omega_0(\tau - \tau'))$, since $e^{i\omega_0\Delta\tau \log n} = n^{i\omega_0\Delta\tau}$. (3) The kernel $K(\Delta\tau) = \int e^{i\omega_0\Delta\tau u} d\mu_\beta(u)$ is the characteristic function of μ_β composed with a dilation, hence positive definite by Bochner’s theorem. (4) follows from unique factorization applied to the logarithm of the phase. $\square$

Equation (31) is the framework’s signature formula: *the distinguishability of two readings of the arithmetic clock is governed by the Riemann zeta function*. This is remarkable; what follows adds the facts that make it usable.

Theorem 5.3 (Almost-periodicity of the clock: the arrow cannot live here). *For every $\beta > \frac{1}{2}$, the kernel $K(\Delta\tau) = \zeta(2\beta - i\omega_0\Delta\tau)/\zeta(2\beta)$ is a Bohr almost-periodic function of $\Delta\tau$ [19]. Consequently:*

1. $|K|$ does not decay along any sequence: $\limsup_{|\Delta\tau| \rightarrow \infty} |K(\Delta\tau)| = 1$.
2. For every $\varepsilon > 0$ the set $\{\Delta\tau : |K(\Delta\tau) - 1| < \varepsilon\}$ is relatively dense: clock correlations resynchronize infinitely often, in both time directions, at every resolution β .

Proof. On the line $\sigma = 2\beta > 1$, $\zeta(s) = \sum_n n^{-s}$ is an absolutely convergent Dirichlet series with $\sum_n n^{-\sigma} < \infty$, hence a (uniform limit of finite exponential sums) Bohr almost-periodic function of the imaginary part [19]. Almost-periodic functions have relatively dense ε -almost-periods, and $K(0) = 1$. $\square$

This theorem resolves a hidden tension in the framework. The zeta object appears twice: as the clock (“the distinguishability of two times is governed by the zeta function”) and as the engine of the thermodynamic arrow (“the cascade up the mountain of composite numbers”). Theorem 5.3 shows these two roles cannot be fused: an almost-periodic clock is *recurrent* and carries no arrow. In the framework the arrow is therefore relocated entirely to the interaction sector and the boundary condition (Conjecture C4 and C7), while the clock contributes the neutral, arrow-free, maximally stable phase reference. This is a structural clarification, not a loss: it says that *within the framework, time measurement and time’s arrow are functionally separated objects*, exactly as relational philosophy of time requires.

5.3 Energy–time duality of the arithmetic clock

Proposition 5.4 (Resolution duality). *For the clock family $|\tau; \beta\rangle$:*

1. The mean energy is $\langle H_P \rangle_\beta = \hbar\omega_0(-\zeta'/\zeta)(2\beta) = \frac{\hbar\omega_0}{2\beta - 1} + O(\hbar\omega_0)$ as $\beta \downarrow \frac{1}{2}$.
2. The correlation peak is asymptotically Lorentzian: as $\beta \downarrow \frac{1}{2}$, uniformly for bounded $\omega_0\Delta\tau/(2\beta - 1)$,

$$|K(\Delta\tau)|^2 = \frac{(2\beta - 1)^2}{(2\beta - 1)^2 + \omega_0^2\Delta\tau^2} (1 + o(1)), \quad (33)$$

with half-width $\Delta\tau_{1/2} \asymp (2\beta - 1)/\omega_0$.

3. Consequently $\langle H_P \rangle_\beta \Delta\tau_{1/2} \rightarrow \hbar$ as $\beta \downarrow \frac{1}{2}$: the zeta clock asymptotically saturates the Mandelstam–Tamm energy–time limit. Time resolution is bought one-for-one with clock energy, and the clock never becomes “cheap.”
4. Conversely, as $\beta \rightarrow \infty$ the state concentrates on $|1\rangle$: $|\tau; \beta\rangle \rightarrow |1\rangle$, energy $\rightarrow 0$, and $K \rightarrow 1$ for all $\Delta\tau$: the clock resolves nothing. At the Janus limit the clock degenerates—the quantified form of the source’s statement that “at $n = 1$ time does not exist.”

Proof. (1) is Theorem 2.6 at $\sigma = 2\beta \downarrow 1$. (2) Near the pole, $\zeta(\sigma - it) = (\sigma - 1 - it)^{-1} + O(1)$, so $|K|^2 = |\sigma - 1|^2 / ((\sigma - 1)^2 + t^2) + o(1)$ with $t = \omega_0 \Delta\tau$ and $\sigma - 1 = 2\beta - 1$. (3) Multiply (1) and (2). (4) $n^{-2\beta}$ concentrates on $n = 1$; $K(t) = \zeta(2\beta - it) / \zeta(2\beta) \rightarrow 1$ since $\zeta(2\beta - it) \rightarrow 1$ and $\zeta(2\beta) \rightarrow 1$. $\square$

5.4 Reconciliation of the three times

With the clock machinery in place, the tripartite reconciliation can be stated as a compact conditional statement. *Quantum-mechanical time* is the reading of a relational clock in the classical-clock limit: it is recovered when the clock subsystem is large, weakly coupled, and quasi-classical, in which case $\tau_{\text{clock}} \approx t$ for all practical observers. *Relativistic time* is the proper-time functional along worldlines: it is the geometric content of the clock’s embedding, determining which phase increments accumulate along which trajectory. *Quantum-gravitational time* is the constraint-generated phase correlation: with the primon clock, τ is conjugate to $\log \hat{N}$, and evolution is the multiplicative phase flow $n^{-i\omega_0\tau}$. The three are not rivals but levels: the block-level correlation structure ($\hat{\mathcal{H}}\Psi = 0$), its geometric projection (proper time), and its semiclassical limit (external t). All incompatibilities of principle dissolve; what remains open—and is honestly recorded—is whether the *specific* identification of the clock with the primon spectrum is realized (Conjecture C1), and how the geometric sector emerges from the arithmetic one (Conjectures C3, C9).

The synthesis deserves to be quoted, because every clause now carries a theorem or a registered conjecture:

Relativity provides the geometry; quantum mechanics provides the spectrum; the primes provide the arithmetic order of the spectrum; and time is the phase relation among prime-ordered states measured along geometric worldlines.

The measurement clause is Proposition 5.2(1) (clock covariance); the ordering clause is unique factorization, i.e. Proposition 5.2(4); the arrow clause is conspicuously absent, by Theorem 5.3, and lives in the interaction sector.

6 Thermodynamics, Gravity, and the Holographic Bridge

6.1 The exact counting law

A repeatedly used statement is “the number of states below energy E grows like $e^{E/\hbar\omega_0}$.” The exact statement is a one-line identity, and its precision is what makes the holographic bridge in Section 6.2 clean.

Theorem 6.1 (Log-count identity). *Let $N(E) = \#\{n \in \mathbb{N}_{\geq 1} : \hbar\omega_0 \log n \leq E\} = \lfloor e^{E/\hbar\omega_0} \rfloor$. Then for $E \geq \hbar\omega_0 \log 2$,*

$$\frac{E}{\hbar\omega_0} - \log 2 \leq \log N(E) \leq \frac{E}{\hbar\omega_0}, \quad (34)$$

and the defect $r(E) = \log N(E) - E/\hbar\omega_0$ satisfies $-\log 2 \leq r(E) \leq 0$ with $r(E) \rightarrow$ (sawtooth oscillations of amplitude $O(1)$); it does not converge. The microcanonical entropy $S(E) =$

$k_B \log N(E)$ satisfies

$$S(E) = \frac{E}{T_H} + O(k_B), \quad T_H = \frac{\hbar\omega_0}{k_B}, \quad (35)$$

and the smoothed microcanonical temperature equals T_H identically, at every energy: the primon gas is an exactly critical Hagedorn system.

Proof. $N(E) = \lfloor e^{E/\hbar\omega_0} \rfloor$ because $\hbar\omega_0 \log n \leq E \iff n \leq e^{E/\hbar\omega_0}$. For $y = e^{E/\hbar\omega_0} \geq 2$, $y - 1 \geq y/2$, so $\log(y/2) \leq \log \lfloor y \rfloor \leq \log y$. The sawtooth is $\log \lfloor y \rfloor - \log y = \log(\lfloor y \rfloor / y) \in (-1/y - \log y, 0]$ oscillating between $-\log 2$ and 0; it never converges because $\log \lfloor y \rfloor$ jumps at $y \in \mathbb{N}$ while $\log y$ is continuous. The temperature statement follows by differentiating the smoothing $\tilde{N}(E) = e^{E/\hbar\omega_0}$. $\square$

The constant-amplitude oscillation of $r(E)$ is not a defect but a signature: it is the microscopic granularity of the integer lattice—the “breathing” of the floor function. In Conjecture C5 this sawtooth reappears as the expected fluctuation sector of the gravitational counting function, dual to the oscillatory term of the explicit formula $\psi(x) = x - \sum_{\rho} x^{\rho}/\rho + \dots$, whose fluctuations under the Riemann hypothesis are bounded by $O(\sqrt{x} \log^2 x)$ —the content of von Koch’s equivalence [20]: the Riemann hypothesis is *equivalent* to the statement that arithmetic fluctuations about the smooth classical term are as small as square-root-small. Read in the framework’s dictionary, RH becomes: *quantum fluctuations of the arithmetic sector about the classical continuum are optimally suppressed*. That is a rigorous reformulation of “RH = spectral stability”, and it is the strongest form of that claim that mathematics currently licenses.

6.2 The holographic bridge

The derivation of the Bekenstein–Hawking entropy from prime state counting is a natural target. This paper isolates exactly one nontrivial input and shows the rest is Theorem 6.1.

Conjecture 6.1 (Arithmetic holographic bound). *There exists a universal constant $\kappa = O(1)$ such that the maximal primon excitation energy accessible in, or associated with, a spacetime region whose boundary has area A is*

$$E_{\max}(A) = \frac{\hbar\omega_0}{4\ell_P^2} A + O(\hbar\omega_0 \cdot \kappa), \quad (36)$$

independently of the detailed matter content, where $\ell_P = \sqrt{G\hbar/c^3}$ is the Planck length.

Theorem 6.2 (Conditional Bekenstein–Hawking). *Assume Conjecture 6.1. Then the arithmetic entropy of the region is*

$$S(A) = k_B \log N(E_{\max}(A)) = \frac{k_B A}{4\ell_P^2} + O(k_B), \quad (37)$$

which is the Bekenstein–Hawking entropy [21, 22] up to the $O(1)$ (logarithmic-correction class) error. Moreover, for a Schwarzschild black hole of mass M , $A = 16\pi G^2 M^2 / c^4$ and $A/4\ell_P^2 = 4\pi G M^2 / (\hbar c)$; the choice $E_{\max} = M c^2$ and $\hbar\omega_0 = 2k_B T_H^{\text{BH}}$ saturates (36) exactly, with $T_H^{\text{BH}} = \hbar c^3 / (8\pi G M k_B)$ the Hawking temperature.

Proof. Substitute (36) into Theorem 6.1. For the Schwarzschild case, $E_{\max}(A)/\hbar\omega_0 = A/4\ell_P^2$ by (36); with $E_{\max} = M c^2$ this forces $\hbar\omega_0 = 4M c^2 \ell_P^2 / A = 4M c^2 \cdot (G\hbar/c^3) / (16\pi G^2 M^2 / c^4) = \hbar c^3 / (4\pi G M) = 2k_B T_H^{\text{BH}}$. $\square$

Two honest remarks are appended to the register entry C3.

Remark 6.3 (The factor-of-two ambiguity). The Hawking identification above is exact only per-horizon: since $T_H^{\text{BH}} \propto M^{-1}$, a *universal* ω_0 cannot equal $2T_H^{\text{BH}}$ for all masses. Either ω_0 is universal and the identification holds to within $O(1)$ horizon-dependent factors, or the arithmetic scale is set per horizon. Which alternative survives is a genuine open question that the conjecture form makes explicit; a presentation that silently treats ω_0 as universal and the identification as exact would hide it. In both alternatives, the entropy itself is exact up to $O(1)$, because Theorem 6.1 controls the error.

Remark 6.4 (What the bridge does and does not derive). The bridge converts an area law into an *energy* cutoff statement plus an exact counting theorem; it does not derive the cutoff. This is precisely the logical position of the holographic principle in mainstream theory, translated into arithmetic terms. The paper’s contribution is that the conversion loss is provably zero (Theorem 6.1 has no hidden degeneracy factors, since the primon spectrum is simple, Theorem 2.4): any slack in Conjecture 6.1 is slack in the physics, not in the counting.

6.3 Singularities and the Hagedorn wall

The singularity claims swing between two poles: one places the primon gas *at* the singularity (the reported conformal system near the black-hole center), later reading declares that singularities *do not exist*. The reconciliation is a standard move of effective field theory, and it is worth stating once, cleanly, because it removes the apparent contradiction:

Remark 6.5 (Resolution of the singularity tension). “Singularity” denotes the boundary of validity of the continuum effective description, not a locus in the fundamental theory. In the reconstructed framework: the geometric (Einstein) description fails where its implicit continuum counting diverges; the arithmetic description remains finite at all energies (Theorem 2.4 has no finite accumulation point, and all quantities are finite below T_H). The pole of ζ at $s = 1$ is the exact analytic signature of the breakdown: it is a *critical* (Hagedorn) divergence, not a curvature divergence. What the Big Bang and black-hole interiors “are” in this framework is the regime where the geometric effective theory must be replaced by the arithmetic one; the primon gas is the UV-complete description of *that regime*, which is compatible with it also being located, in the geometric coordinates of the effective theory, “near the singularity.” Both readings are thereby retained, each in its proper tense. The associated physical conjecture (C6) is that the Hagedorn wall T_H replaces the singular point: approach to the singularity is approach to a critical temperature at which energy converts into primon creation rather than curvature growth.

6.4 Emergent gravity and the hierarchy problem

The resolution of the hierarchy problem—gravity as an entropic, $1/N$ -suppressed collective effect [23–25]—is the most speculative of its physics claims, and it is demoted in full. But the demoted form can be made unusually concrete, because the arithmetic sector supplies an explicit N .

Conjecture 6.2 (Arithmetic hierarchy). *The Newton constant is set by the number of arithmetic degrees of freedom below the effective cutoff E_{cut} :*

$$\frac{1}{G} \asymp \frac{\hbar}{c^3} \omega_0 N_{\text{eff}}(E_{\text{cut}}), \quad N_{\text{eff}}(E) = \lfloor e^{E/\hbar\omega_0} \rfloor, \quad (38)$$

up to a dimensionless $O(1)$ factor, so that

$$\log \frac{M_{\text{Pl}}}{M_{\text{EW}}} \asymp \frac{E_{\text{cut}}}{2\hbar\omega_0} \implies E_{\text{cut}} \asymp 2 \log\left(\frac{M_{\text{Pl}}}{M_{\text{EW}}}\right) \hbar\omega_0 \approx 74 \hbar\omega_0 \quad (39)$$

(using $M_{\text{Pl}}/M_{\text{EW}} \approx 10^{16}$).

The honest content of Conjecture 6.2 is a *relocation* of the hierarchy problem, stated with numbers: the question “why is gravity so weak?” becomes “why does the effective arithmetic universe of the electroweak regime terminate at $n \sim e^{74} \approx 10^{32}$ integers?”—which is a question about the cutoff structure of the arithmetic sector, not a fine-tuning of a Lagrangian parameter. The framework does not answer it; it converts it into a well-posed question with a single dimensionless number to explain, register entry C11. This is strictly more than a bare $1/N$ -suppression narrative with no N , and strictly less than a derivation of G .

6.5 The classical limit and the pole at unity

The classical-limit condition reads: as $s \rightarrow 1^+$, the coarse-grained arithmetic partition function should reproduce classical geometry, the pole $\zeta(s) \sim 1/(s-1)$ representing the divergent continuum limit. This paper restates this as a theorem-flanked analogy: the pole is exactly the Hagedorn divergence (Theorem 2.6); the smooth term x of $\psi(x)$ is the classical contribution and the zeros contribute the fluctuations (von Koch’s equivalence [20]); and “renormalization removing the pole” means passing to the finite part of the Laurent expansion, whose constant term is Euler’s constant γ —the arithmetic framework’s first finite counterterm. Whether the geometric sector (Einstein dynamics, $G_{\mu\nu} + \Lambda g_{\mu\nu} = 8\pi G T_{\mu\nu}$) can actually be *derived* as this coarse-graining is the deepest part of Conjecture C1 and is not claimed.

7 Entanglement: Arithmetic Superpositions and Holism

7.1 Composite integers are product states

The framework’s informal layer identifies quantum entanglement with multiplicative composition: “a composite number is mathematically entangled: it cannot be described independently of its prime factors”; “entanglement is not non-local communication, it is multiplicative inseparability”; “a composite number like $n = p_A \cdot p_B$ is a single, inseparable mathematical entity.” As quantum theory this is false. The correction relocates arithmetic entanglement to where it genuinely lives—superpositions across prime-mode bipartitions—and preserves the arithmetic content as structure rather than as a false identification.

Theorem 7.1 (Product structure of the occupation basis). *Let $P_A, P_B \subset \text{Pr}$ be disjoint with $P_A \cup P_B = \text{Pr}$, and let $\mathcal{F}_A, \mathcal{F}_B$ be the submonoids of $\mathcal{F}$ supported on P_A, P_B . The monoid isomorphism $\mathcal{F} \cong \mathcal{F}_A \times \mathcal{F}_B$ (unique factorization: $k \mapsto (k|_{P_A}, k|_{P_B})$) induces a unitary*

$$\mathcal{H}_A \cong \mathcal{H}_A \otimes \mathcal{H}_B, \quad |k\rangle \mapsto |k_A\rangle \otimes |k_B\rangle, \quad (40)$$

under which every occupation basis vector is a product state. In particular $|pq\rangle = |p\rangle_A \otimes |q\rangle_B$ for $p \in P_A, q \in P_B$: a composite integer is an unentangled configuration of its prime factors. Entanglement across the bipartition is a property of superpositions $\psi = \sum c_{ij}|i\rangle_A|j\rangle_B$ of rank > 1 in the coefficient tensor, exactly as in ordinary quantum mechanics, and of nothing else.

Proof. The restriction map is a bijection of free commutative monoids by unique factorization; the induced map on ℓ^2 bases is a bijection of orthonormal bases preserving inner products; extend linearly. Product structure of each basis vector is the definition of the tensor identification. $\square$

The composite-as-entangled picture thus becomes a precise dichotomy: the integer n *labels* a product configuration; *arithmetic entanglement* is superpositional. The dichotomy is stronger than the identification it replaces, because it carries quantitative content: entanglement entropies, Schmidt ranks, and Bell inequalities.

Definition 7.2 (Arithmetic entanglement). For a bipartition $\text{Pr} = P_A \sqcup P_B$ and a pure state $\psi \in \mathcal{H}_A \otimes \mathcal{H}_B$, the *arithmetic entanglement entropy* $S_{\text{ent}}(\psi)$ is the von Neumann entropy of

$\rho_A = \text{Tr}_B |\psi\rangle\langle\psi|$. ψ is entangled iff $S_{\text{ent}}(\psi) > 0$, iff the coefficient tensor (c_{ij}) in the occupation basis has rank > 1 when reshaped as a matrix.

Example 7.3 (The six–thirty-six Bell pair). Take $P_A = \{2\}$, $P_B = \{3\}$. Then

$$\frac{1}{\sqrt{2}}(|6\rangle + |36\rangle) = \frac{1}{\sqrt{2}}(|2\rangle_A |3\rangle_B + |4\rangle_A |9\rangle_B) \quad (41)$$

is maximally entangled across the bipartition: Schmidt rank 2, equal Schmidt values $1/\sqrt{2}$, $S_{\text{ent}} = \log 2$. The superposition of the integers 6 and 36 is an arithmetic Bell pair. More generally $\frac{1}{\sqrt{2}}(|ab\rangle + |cd\rangle)$ with a, c powers of distinct primes of P_A and b, d of P_B , $(a, b) \neq (c, d)$, is maximally entangled.

Proposition 7.4 (Bell violation in the arithmetic sector). *Restrict local measurements to the two-dimensional subspaces $V_A = \text{span}\{|2\rangle, |4\rangle\} \cong \mathbb{C}^2$ and $V_B = \text{span}\{|3\rangle, |9\rangle\} \cong \mathbb{C}^2$, identified with qubits. On the Bell state of Example 7.3, the standard spin settings along the xz -plane achieve the CHSH value $2\sqrt{2}$, the Tsirelson bound [26, 27]. Hence the arithmetic sector, equipped with superpositional states, exhibits exactly standard quantum non-locality of correlations; the violation is destroyed if the superposition is replaced by either basis vector, i.e. by any definite integer.*

Proof. The restriction of the state to $V_A \otimes V_B$ is the standard Bell state $(|00\rangle + |11\rangle)/\sqrt{2}$; the CHSH computation is the textbook one. $\square$

7.2 Splitting primes as natural entanglement resources

The Gaussian sector supplies a construction the rational sector lacks, and it realizes two images at once (the “complex primon overlay” of the five-dimensional picture and the claim that symmetry breaking is prime splitting).

Proposition 7.5 (Splitting-prime EPR pairs). *Let $p \equiv 1 \pmod{4}$ be a rational prime, so that $p = \pi\bar{\pi} = N(\pi)$ in $\mathbb{Z}[i]$ with π a Gaussian prime, $\pi \sim \bar{\pi}$ conjugate and non-associate. Define the splitting state*

$$|p\rangle_{\text{split}} = \frac{1}{\sqrt{2}}(|\pi\rangle \otimes |\bar{\pi}\rangle + |\bar{\pi}\rangle \otimes |\pi\rangle) \quad (42)$$

on the two Gaussian modes generated by π and $\bar{\pi}$. Then:

1. $|p\rangle_{\text{split}}$ is maximally entangled across the $(\pi, \bar{\pi})$ bipartition ($S_{\text{ent}} = \log 2$);
2. it is invariant under the conjugation involution C (which swaps the two tensor factors), so it is even under the arithmetic charge conjugation of Section 8;
3. it is the natural correlate, inside the arithmetic Hilbert space, of the splitting of p in $\mathbb{Z}[i]$ —the phenomenon whose statistical frequency is exactly $1/2$ by Chebotarev/Dirichlet (Section 8).

Proof. (1) The two tensor factors are qubit-spaces $\text{span}\{|\pi\rangle, |\bar{\pi}\rangle\}$; the state is the symmetric Bell state. (2) C exchanges $|\pi\rangle \leftrightarrow |\bar{\pi}\rangle$ in each factor, and the state is manifestly symmetric. (3) The correspondence $p \mapsto \{\pi, \bar{\pi}\}$ is the Dedekind–Kummer splitting theorem for $\mathbb{Z}[i]$ [12]. $\square$

Proposition 7.5 is the precise version of the “phantom layer of Gaussian primes weaving through the solid prime nodes”; the weaving is entanglement between conjugate prime modes, quantitatively $\log 2$ per split integer, and it exists only for the splitting class $p \equiv 1 \pmod{4}$, which by Dirichlet’s theorem has natural density $1/2$. The inert class $p \equiv 3 \pmod{4}$ contributes *no* entanglement resource: those primes remain prime in $\mathbb{Z}[i]$. The arithmetic sector thus possesses an intrinsic, algebraically defined binary—split versus inert—with exact statistics. Section 8 builds the force-sector dictionary on it; it is also the natural habitat for the idea that the weak (chiral) interaction lives in the complex sector.

7.3 Measurement, decoherence, and records

With entanglement defined correctly, the measurement picture—“a measurement is an interaction that creates a thermodynamically irreversible, decohered record in the prime spectrum”—translates into standard quantum mechanics with an arithmetic state space, and the standard strengths *and* standard weaknesses apply.

Remark 7.6 (Measurement as record formation). In this framework a measurement is an interaction between a system and a macroscopic apparatus/environment whose net effect is to entangle the system’s superposition across the bipartition “apparatus modes versus system modes” so strongly that the reduced state of the system is, to observational accuracy, diagonal in the pointer basis (einselection) [28]. The arrow enters through the thermodynamic cost of stabilization (Landauer’s principle [29]): records can be written reliably only in the direction of increasing multiplicity of environmental states, i.e. along the cascade of Conjecture C4. Rocks measure rocks, photons measure dust grains, and consciousness is a consumer of records, not a precondition for them—each clause survives, now attached to the correct mechanism (superposition entanglement, Theorem 7.1) rather than the incorrect one (composite inseparability).

Remark 7.7 (The improper-mixture problem). Decoherence diagonalizes reduced density matrices; it does not, by itself, select one outcome. The framework elects an Everettian resolution [30], registered as interpretive (Conjecture C12) rather than theorem-grade. What the arithmetic sector adds is structural, not probabilistic: branching in $\mathcal{H}_A$ is branching of *factorization histories*—worldlines through the divisibility graph—and the Born-rule weights are the squared amplitudes of superposed integers. The 2019 reversal experiments on small qubit registers [31] are consistent with this picture in the same sense as with standard decoherence: small closed systems can be complex-conjugated back; macroscopic entangled cascades cannot, because the required inverse unitary acts on $\sim 10^{80}$ correlated modes and its probability is beyond every Poincaré time. The experiments were performed by Lesovik et al. and implemented on an IBM quantum computer.

7.4 Bell’s trilemma and the rejection of superdeterminism

Superdeterminism can be rejected with the argument that Bell correlations arise from “multiplicative inseparability” of composites. That mechanism is false (Theorem 7.1), so the rejection must rest on other grounds: the standard Everettian one, plus the arithmetic sector’s own contribution:

1. *Realism, global not local.* The arithmetic Hilbert space and the universal state are observer-independent; individual prime modes carry no pre-existing local values (the occupation basis is a basis, not a value assignment). This is exactly the quantum theory’s own position: realism preserved globally, rejected locally.
2. *Locality, no-signaling not local action.* Arithmetic-entangled states (Examples 7.3, Proposition 7.5) exhibit Bell correlations across spacelike separation (Proposition 7.4) while the reduced states are untouched by distant operations: $\rho_B = \text{Tr}_A \rho_{AB}$ is invariant under local unitaries on A . No usable signal, energy, or conserved quantity propagates superluminally; the no-signaling theorem is a theorem of the tensor structure that Theorem 7.1 now correctly exhibits.
3. *Statistical independence, preserved.* Measurement settings and hidden variables are not pre-correlated by a cosmic conspiracy. The correlations of the Bell state are *created* by the state’s preparation (a superposition), not imposed on separate local beables. Superdeterminism saves locality only by denying independence; the arithmetic framework keeps independence and gives up local realism of modes, exactly as loophole-free experiments [32] force [26, 27].

The corrected position is the standard one, reached with the arithmetic sector’s own structure. The philosophical residue—that “space” itself is emergent from entanglement structure, so that demands for local causal explanation between entangled modes are a category error—remains available as an interpretation (register C12), in the same epistemic tier as the ER=EPR-type conjectures it echoes. What is theorem-grade is narrower and cleaner: in the arithmetic sector, *non-locality of correlations is a property of superposed integers, local realism of prime modes is false, and no-signaling is exact.*

8 The Arithmetic Gauge Sector

8.1 From Lie unification to Galois unification

The framework proposes that the Standard Model gauge group $U(1) \times SU(2) \times SU(3)$ is the “macroscopic, continuous shadow” of the Absolute Galois group $G_{\mathbb{Q}} = \text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$, with the Langlands program as the dictionary, prime splitting as symmetry breaking, and Gaussian primes as the chiral (weak) sector. As physics this is speculation; as mathematics it can be anchored on all fours, because each ingredient of the analogy corresponds to an actual theorem. This section therefore proceeds in two layers: a *theorem layer* that fixes what is true about splitting statistics and symmetry structure, and a *dictionary layer* that is demoted in its entirety (Conjecture C9).

The theorem layer rests on three classical results, which we state in the form the analogy uses them.

Theorem 8.1 (Kronecker–Weber). *Every finite abelian extension $K/\mathbb{Q}$ is contained in a cyclotomic field $\mathbb{Q}(\zeta_m)$; conversely every cyclotomic field is abelian over $\mathbb{Q}$ [12]. Combined with Dirichlet’s theorem on primes in arithmetic progressions [33] and the splitting criterion*

$$p \nmid m, \text{ } p \text{ splits completely in } \mathbb{Q}(\zeta_m) \iff p \equiv 1 \pmod{m}, \quad (43)$$

the abelian sector of $G_{\mathbb{Q}}$ is completely described by residue classes of primes: the reciprocity map of class field theory is, explicitly, $p \mapsto p \pmod{m}$.

Theorem 8.2 (Dedekind–Kummer splitting). *Let $K = \mathbb{Q}(\theta)$ with θ an algebraic integer of minimal polynomial f , and p a rational prime not dividing the index $[\mathcal{O}_K : \mathbb{Z}[\theta]]$. Then the factorization pattern of f modulo p into irreducibles of degrees $d_1, \dots, d_r$ coincides with the splitting pattern of p in $\mathcal{O}_K$: $p\mathcal{O}_K = \mathfrak{p}_1 \cdots \mathfrak{p}_r$ with residue degrees $d_1, \dots, d_r$ [12]. Symmetry breaking and prime splitting are the same operation.*

Theorem 8.3 (Chebotarev density). *Let $K/\mathbb{Q}$ be a finite Galois extension with group G , and $\sigma \in G$ any element. The set of rational primes p (unramified in K) whose Frobenius class $\text{Frob}_p \subset G$ equals the conjugacy class of σ has natural density $|\text{cl}(\sigma)|/|G|$ [34]. In particular: splitting statistics are group statistics. Dirichlet’s theorem is the abelian case (with Theorem 8.1).*

For the working example that carries the whole source dictionary, specialize to $K = \mathbb{Q}(i)$, $G = \text{Gal}(\mathbb{Q}(i)/\mathbb{Q}) = \{1, \sigma\}$ with σ the complex conjugation $a + bi \mapsto a - bi$:

Class of p	Condition	Behavior in $\mathbb{Z}[i]$	Density (Chebotarev)
Split	$p \equiv 1 \pmod{4}$	$p = \pi\bar{\pi}$, two conjugate primes	$1/2$
Inert	$p \equiv 3 \pmod{4}$	p remains prime	$1/2$
Ramified	$p = 2$	$2 = -i(1+i)^2$, one prime squared	0 (finite)

8.2 The dictionary

With the theorem layer fixed, the force-to-field dictionary can be stated so that every arithmetic entry is backed and every physical equation is demoted:

Standard Model object	Arithmetic object	Status of the pairing
U(1) gauge structure	Abelian extensions, cyclotomic fields $\mathbb{Q}(\zeta_m)$; the “photon” is the reciprocity map $p \mapsto p \pmod{m}$	Theorem layer: KW, Dirichlet (Thm 8.1); pairing demoted (C9)
Chiral $SU(2)_L$, parity violation	Quadratic/non-abelian extensions; the non-trivial automorphism σ (conjugation) acts as the arithmetic C ; split/inert dichotomy as handedness	Theorem layer: splitting table above; pairing demoted (C9, C10)
$SU(3)$ color, confinement	Inert primes ($p \equiv 3 \pmod{4}$) and their non-abelian analogues; higher-degree splitting patterns with non-commuting Frobenius classes	Theorem layer: Chebotarev (Thm 8.3); pairing demoted (C9)
Higgs mechanism, spontaneous symmetry breaking	Ramification and splitting: the unified base $\mathbb{Q}$ extends to K ; primes then <i>choose</i> splitting patterns, with exact group-valued statistics	Theorem layer: Kummer (Thm 8.2); pairing demoted (C9)
Coupling unification, charge quantization	Langlands functoriality: transfer of automorphic representations between L -groups; the dictionary itself	Mathematics: deep conjecture, proven for GL_n over function fields [35]; physics: demoted (C9)
CP violation, CKM phase	The σ -odd interaction sector: asymmetric couplings $w(\pi) \neq w(\bar{\pi})$; the non-vanishing $L(1, \chi_{-4}) = \pi/4$ as the arithmetic asymmetry functional	Theorem layer: factorization (17); model demoted (C10)

Each row now has the same structure: a true theorem about number fields, an honest conjecture about the pairing. Two entries deserve expansion because they carry the most distinctive physical claims.

8.3 Arithmetic charge conjugation, parity, and time reversal

The arithmetic model of the discrete symmetries can be defined exactly on the Gaussian sector.

Definition 8.4 (Arithmetic C , P , T). On $\mathcal{H}_{\mathbb{Z}[i]}$ (Definition 2.8):

1. C is the unitary implementing $\text{Gal}(\mathbb{Q}(i)/\mathbb{Q})$: $C|\pi\rangle = |\bar{\pi}\rangle$ on Gaussian-prime modes, extended multiplicatively. It is an involution, $C^2 = I$.
2. P is any order-two relabeling of modes that plays the role of spatial inversion (e.g. a fixed-point-free involution of a finite mode set).
3. T is the antiunitary complex conjugation K in the occupation basis: it reverses all phases, $K n^{-i\omega_0\tau} K = n^{+i\omega_0\tau}$.

Proposition 8.5 (Symmetry structure of the Gaussian sector). 1. $[C, H_{\mathbb{Z}[i]}] = 0$ and $[K, H_{\mathbb{Z}[i]}] = 0$: the free Gaussian Hamiltonian is exactly C -, T -, and CPT -symmetric, because it is built from norms $N(\pi)$, which are conjugation-invariant, with real coefficients.

2. Consequently any arithmetic model of CP violation must reside in the interaction sector: couplings $w(\pi)$ with $w(\pi) \neq w(\bar{\pi})$ in the walk/hop Hamiltonians of Section 3.3. The σ -even/ σ -odd decomposition $H = H_+ + H_-$ (eigenparts under C) is an orthogonal splitting of the interaction space, and the “amount of CP violation” is the norm ratio $\|H_-\|/\|H_+\|$ together with the relative phases of the w ’s.
3. The composition $CPT = CP \cdot K$ is antiunitary; if $w(\pi) = \overline{w(\bar{\pi})}$ (the natural hermiticity condition), the full H_{tot} is CPT -symmetric, the arithmetic echo of the CPT theorem holding by construction rather than by axiomatic field theory.

Proof. (1) $N(\bar{\pi}) = N(\pi)$; diagonal operators with real eigenvalues commute with the occupation-basis conjugation K . (2) The even/odd split under an involutive unitary is standard. (3) Direct computation: CP exchanges the couplings $w(\pi) \leftrightarrow w(\bar{\pi})$; combined with K (which conjugates the phases), hermiticity in the stated form is exactly the invariance condition. $\square$

The parallel with the Standard Model is exact in structure and honest in status: in both theories, the free (kinetic) sector is CP -symmetric; the violation lives in the charged interaction sector (the CKM phase there, the asymmetric weights $w(\pi)$ here); and CPT holds by the hermiticity structure. The claim that “the weak interaction maps to the complex sector and its rare CP violation to complex phases” becomes Conjecture C10: *the CKM phase and baryon asymmetry are the arithmetic index of the σ -odd interaction sector, accumulated along the cascade*. Its falsifiable edge is sharp: the model predicts that CP -violating and T -violating observables couple to *split-versus-inert* statistics (the only binary the Gaussian sector possesses), so any derived branching asymmetry must be expressible through L -function ratios at saddles—a constraint precise enough to check and to fail.

8.4 Vaccaro’s program, translated

Vaccaro proposes that T -violation is not a curiosity of the weak sector but the microphysical origin of dynamics itself: with exactly T -symmetric laws, the universe would be a static block without localized happening [36]. The arithmetic translation is natural in this framework: the σ -odd couplings H_- are precisely what distinguishes the interacting block from the frozen block. Under H_P alone the universe is frozen (Proposition 3.1); under $H_P + H_+$ it evolves but with exactly reciprocal processes (detailed balance holds, and the walk is unbiased); under $H_P + H_+ + H_-$ there is a microscopic bias, a ratchet, and the direction of the cascade acquires a microphysical anchor in addition to the boundary condition. Conjecture C10 subsumes this: the ratchet, compounded by the exponential state growth (Theorem 2.6), is the framework’s candidate engine for baryogenesis—a “chiral arithmetic cascade” with a mechanism that exists.

8.5 What Langlands does and does not supply

Finally, the boundary of the unification layer. The Langlands program ([35] for the proven function-field GL_n case) supplies a conjectural correspondence between Galois representations and automorphic forms: in the framework’s language, between the arithmetic sector’s symmetry data and the analytic (eigenmode) data of adelic groups. The slogan—“the continuous gauge symmetries are the low-energy shadows of discrete Galois symmetries”—requires, at minimum, that such correspondences survive specialization to the finite set of splits that the Standard Model actually uses. Nothing currently known implies this; nothing currently known forbids it. This paper’s contribution is to isolate the minimal well-posed core:

Conjecture 8.1 (Galois–gauge dictionary, minimal form). *There exists an assignment from the Standard Model’s gauge and flavor data to number fields and their Galois representations such that: (i) the observed charge lattice corresponds to a lattice of splitting restrictions of the form (43); (ii) branching ratios of force sectors equal Chebotarev densities of the associated Frobenius classes; and (iii) the hierarchy of coupling strengths corresponds to the ramification hierarchy of the fields involved.*

Condition (ii) is the conjecture’s empirical hook: Chebotarev densities are rational numbers with denominator $|G|$ —the arithmetic GUT predicts that observed branching statistics, at the unification scale, organize into ratios of small integers divided by group orders. That is a checkable structural signature, quite unlike the free parameters of Lie-group GUTs, and it is the kind of prediction that would falsify C9 outright if the statistics came out transcendental or unsplittable.

9 Arrows of Time and the Philosophy Register

The philosophical layer of the framework is unusually rich: it adjudicates between A-theory and B-theory, answers the vertiginous question, classifies the intrinsic and extrinsic camps, reconstructs the Past Hypothesis, rejects cyclical time, demotes causality, and rejects superdeterminism. The task of this section is different from the mathematical sections: the claims are not theorems and should never pretend to be; the task is to state each position *conditional on* the conjectural superstructure, mark its epistemic tier, and keep the ones that are well supported while discarding the arguments that leaned on loose mathematics. Every item below is either (a) a conditional consequence of registered conjectures, (b) an interpretive stance, or (c) a theorem already proved in Sections 3–8.

9.1 The extrinsic arrow, structurally grounded

The position in the intrinsic/extrinsic debate is extrinsic, but “not arbitrary”: the arrow is a structural asymmetry of the block, not a primitive flow. This section retains this and sharpens it into a three-source decomposition:

1. *Geometric order* (relativistic causal structure): a time-orientable spacetime supplies an earlier/later order and light cones—the riverbed, not the flow. This is standard general relativity and carries no arrow.
2. *Statistical multiplicity* (the arithmetic Hagedorn structure): Theorem 6.1 fixes $S(E) = E/T_H + O(1)$; the state space grows exponentially in one direction of complexity. This is theorem-grade but is an asymmetry of *counting*, not yet of *time*: it becomes a temporal arrow only under a boundary condition that places low multiplicity in one temporal direction.
3. *The boundary condition* (the Past Hypothesis, arithmetic form): the low-entropy past is the Janus state $|1\rangle$ or its neighborhood (unique zero-entropy state, Theorem 2.4). That the universe’s actual history occupies such a neighborhood at one end is Conjecture C7, and the two-sided (Janus-point) version—entropy increasing away from $|1\rangle$ in both temporal directions—is the arithmetic realization of the Carroll–Chen and Barbour et al. constructions [37–39], registered as C8.

The arrow formula (“causal order + time orientation + low-entropy boundary + spectral state counting”) survives verbatim, but its ingredients are sorted by tier: the first is geometry, the second is a theorem, the third is a conjecture. Nothing requires time to flow; everything requires the block to be asymmetric.

9.2 The hierarchy of arrows

The hierarchy—an arithmetic master arrow from which thermodynamic, causal, and psychological arrows derive, with the cosmological arrow as boundary condition—survives with the master arrow’s engine supplied. The version adopted here:

Arrow	In this framework	Status
Arithmetic (master)	Growth of available multiplicity $N(E)$; cascade of Conjecture C4	Conjecture (engine) + Theorem (counting)
Thermodynamic	$S = k_B \log \Omega$ along the cascade; second law as overwhelming-multiplicity drift	Conditional on C4, C7
Causal	Record formation: asymmetric decoherence structure [28, 29]; “ c causes e ” = e contains a thermodynamically stable record of c	Conditional; derivative, not fundamental (cf. [40, 41])
Psychological	Memory as record-bearing physical process; indexical “now”	Conditional; interpretive tier
Cosmological	Expansion as the geometric enabler of the arithmetic cascade; alignment of arrows contingent on epoch	Conditional; observation-compatible
Weak (CP/T)	σ -odd interaction sector (Proposition 8.5); micro-ratchet, not master arrow	Conjecture C10

Two notes attach to this table. First, the causal arrow’s derivation presupposes the thermodynamic arrow, so causal theories of time inherit the boundary-condition problem rather than solving it. Second, the weak arrow is *demoted*: nothing in the mathematics forces CP violation to be a separate fundamental arrow, but nothing derives it from the counting either; it enters as an independent interaction-sector fact (Conjecture C10), consistent with Vaccaro’s program [36] without being implied by it.

9.3 Cyclical time: the verdict

Section 4 proves the layered verdict: exact eternal return is impossible for generic states (Theorem 4.2); ε -return is nevertheless universal for the free flow, for every state, with relatively dense but astronomically non-uniform return times (Theorem 4.4); and the multiplicative cascade is transient for $d \geq 3$ modes (Proposition 4.5). The philosophical consequence—“the river can never flow in a circle”—survives in the layered form: exact cycling is a theorem-grade impossibility, practical cycling is excluded by Diophantine slowness of the returns, and the cascading dynamics genuinely never returns for three or more active modes. The framework thereby gains two things: the recognition that the *free* spectrum is maximally recurrent, so that every arrow of time must be manufactured by the interaction sector; and a *parameter-controlled* transition between recurrent and transient arithmetic universes (the number of active prime modes), which converts a metaphysical prohibition into a physically meaningful question.

9.4 Eternalism, indexicality, and the vertiginous question

The position—block universe at the fundamental level, perspectival observers inside it, rejection of both the moving spotlight and egocentric presentism—is retained, with its argument tightened at the two points where it leaned on loose physics.

Remark 9.1 (Indexical perspectival realism). The “democracy of perspectives” does not follow from the symmetry of the constraint $\hat{\mathcal{H}}\Psi = 0$ alone. A constraint’s symmetry does not by itself create perspectives; what creates them is the decomposition of Ψ into entangled clock–system–observer branches, which is exactly the Page–Wootters structure of Section 5 with the corrected entanglement of Section 7. Each perspective is an indexical fact—“this branch, this clock reading”—of the Kaplanian kind: “now” and “I” are token-reflexives, and their referents are fixed by the utterance event, not by a metaphysical spotlight [42]. Hellie’s vertiginous question (“why is this perspective the one occurring?”) [7] is thereby answered in the same move that answers “why am I here?” on a map: the question has a referent-fixing answer and no metaphysical remainder; what remains is the phenomenology, which is real but not ontology. Conitzer’s strengthened A-theory arguments [9] and Hare’s egocentric presentism [8] are then answered with the felt flow as the experience of record formation along the cascade—but now the cascade has a mechanism that exists (Section 3) and the flow’s substrate is the superpositional entanglement structure, not composite inseparability.

Remark 9.2 (What would make the block falsifiable as an interpretation). The eternalist tier of the framework is interpretation, but it is *tethered* interpretation: it inherits its standing from the constraint formalism (C1), the Page–Wootters clock (theorems), and the cascade conjecture (C4). If C4 failed—if the interacting arithmetic dynamics failed to thermalize—the psychological arrow would lose its engine and the perspectival story would need replacement. This tethering answers the rhetorical question “is the framework well-developed enough to confidently pick a side?”: yes, conditional on the register; no, unconditionally.

9.5 Past Hypothesis as arithmetic necessity: theorem and conjecture

The most striking cosmological claim here is that the low-entropy past is not a brute fact but the arithmetic floor: “the universe began at $n = 1$ because 1 is the only number with no prime factors.” Its correct and incorrect parts split cleanly. *Correct, and theorem-grade*: $|1\rangle$ is the unique state of zero energy and zero entropy (Theorem 2.4); it is the unique minimum of complexity; any trajectory through the multiplicative lattice that touches the floor must pass through it; and a state exactly at $|1\rangle$ is stationary (with the interaction sector switched off, Proposition 3.1)—the quantitative “time does not exist at the floor” (Proposition 5.4(4)). *Conjectural, and demoted*: that the actual universal state was ever at or near $|1\rangle$ (C7), and that the entropy profile is two-sided about it (C8). What the demotion buys is precision about what a derivation would need: a mechanism that places the initial condition at the floor without fine-tuning. The arithmetic framework’s candidate answer—the floor is the *only* state at which the thermodynamic and complexity orderings degenerate, so any boundary-condition principle that is symmetry-like must select it—is registered as the motivation for C7, in the same tier as Wallace’s quantum-cosmological reading of the Past Hypothesis [5] and Price’s symmetry principles [6]. The claim that the framework “dissolves” the Past Hypothesis is downgraded to: the framework *localizes* the Past Hypothesis to a single, mathematically distinguished state, which is progress, not dissolution.

9.6 The Big Crunch, Gold universes, and reverse-arrow regions

The claim under test is that the thermodynamic arrow cannot reverse in a contracting universe because “state-space volume is independent of physical volume.” The conclusion survives with a corrected argument. Entropy is a property of the state, and the available multiplicity $N(E)$

is set by the spectrum, not by the scale factor; but whether the state’s *actual* complexity continues to grow during contraction is a dynamical question, answered affirmatively only under Conjecture C4 (the cascade) plus an energy budget that does not collapse. The statement is conditional: if the cascade persists and the cutoff $E_{\max}$ does not shrink with the scale factor, the arrow persists through a Big Crunch, Gold’s reversal fails [4], and inhabitants of a putative reverse-arrow region would experience their own local multiplicity gradient as “the future”—the indexical resolution, which carries over once the cascade has an engine. If instead the cutoff is geometric (as *holographic* bounds suggest it might be, Conjecture C3), then contraction eventually forces decreasing multiplicity, and the framework itself predicts a Gold-like reversal. The two possibilities are separated cleanly by whether $E_{\max}(A)$ in (36) is a function of area alone: the conjecture form of the holographic bound thus *decides* the Big Crunch question, which a flat assertion cannot.

9.7 Superdeterminism and locality

Section 7 settles the Bell argument. The residue is short to state. The framework preserves statistical independence (no cosmic conspiracy), preserves no-signaling exactly (tensor structure, Theorem 7.1), and sacrifices only local realism of individual prime modes. Superdeterminism is rejected not because it is inconsistent (it is consistent) but because it is epistemically self-undermining and unnecessary: the arithmetic sector produces Bell correlations through superposed integers at no cost to locality of signaling. The “adjacent nodes in a non-spatial network” metaphor is retained as interpretation (C12): spacetime emergence from entanglement structure would indeed render “where” questions ill-posed at the fundamental level, but this remains a conjectural picture of emergence, kin to causal set theory’s own program [41] and to the causal theory of time it must surpass [40].

10 The Consolidated Conjecture Register

The register is the framework’s contract with honesty: every ambitious claim that lacks proof lives here, numbered, graded, and armed with a falsifier. Grades: **A** (mathematically precise; partial evidence exists; actively testable), **B** (well-formed; speculative; no current evidence beyond structural fit), **C** (interpretive; tethered to A/B entries; not empirically separable at present). The *Depends on* column lists register entries whose failure collapses the entry; the qualifier “(int.)” marks a dependency that is interpretive rather than logical — such an entry survives the parent’s failure as an autonomous mathematical problem.

ID	Statement	Grade	Depends on	Falsifier / test route
C1	<i>Spectral equivalence.</i> The renormalized near-singularity gravitational partition function of a 4-dimensional black hole equals ζ (and $\zeta_{\mathbb{Q}(i)} = \zeta L(\chi_{-4})$ in the Gaussian/5D sector), after conformal reduction to a self-adjoint H_{sing} with $H_{\text{sing}} \cong \hbar\omega_0 \log \hat{N}$.	A	— (root)	Derive H_{sing} from BKL/conformal reduction; compute its level statistics (cf. C5). Falsified if the near-singularity spectrum is provably not log-integer. Evidence base is thin—the structural Dedekind fit plus one reported program [3]; the grade reflects testability, not weight of evidence.

ID	Statement	Grade	Depends on	Falsifier / test route
C2	<i>Arithmetic cutoff.</i> The physically realized arithmetic universe is a finite or effectively finite truncation (primes $\leq X$, occupations $\leq K$), geometrically motivated, controlling the recurrence-time scale of Theorem 4.4.	B	C1	Sets the quantitative recurrence horizon (Diophantine return times); falsified by derived recurrence phenomenology inconsistent with any effective cutoff.
C3	<i>Holographic arithmetic bound.</i> $E_{\max}(A) = \frac{\hbar\omega_0}{4\ell_P^2} A + O(\hbar\omega_0\kappa)$ universal (Conj. 6.1); with Theorem 6.1 it yields $S_{\text{BH}} = A/4\ell_P^2 + O(1)$ (Thm 6.2).	A	C1	Positive evidence: the proved conditional Theorem 6.2 (S_{BH} follows exactly once the bound is granted)—the grade certifies that conditional, not independent support. Risk: black-hole entropy log-corrections. The arithmetic model fixes the $O(1)$ term's oscillation structure (sawtooth, non-convergent); mismatch with computed corrections (e.g. known $-\frac{3}{2}\log A$ terms) falsifies the pure model.

ID	Statement	Grade	Depends on	Falsifier / test route
C4	<i>Cascade/ETH, interacting completion.</i> The <i>interacting completion</i> $H_{\text{tot}}^{(U,V)} = H_P + H_W + H_{\text{hop}} + U \sum_{p<q} \hat{n}_p \hat{n}_q + V \sum_{p<q} (\hat{n}_p + \hat{n}_q)(a_p^\dagger a_q + a_q^\dagger a_p)$ (diagonal or kinetic) thermalizes on the prime lattice: $\sigma_{\text{ETH}}(D) \rightarrow 0$ and diagonal $\rightarrow$ microcanonical along truncation sequences at fixed dose (Conj. 3.1). The bare H_{tot} is excluded (Remark 3.9).	A	C1 (int.)	<i>Scaling criterion:</i> ETH diagnostics along truncation sequences with $D \rightarrow \infty$ at fixed effective dose VK^2 — sparse eigen-windows to $D = 5.9 \times 10^4$ and Krylov equilibration to $D \approx 10^5$; falsified if $\sigma_{\text{ETH}}(D)$ never decays and $\text{diag} - \text{micro}$ stays $O(1)$ for every interacting completion. Current evidence: the bare system is decided—structurally non-ergodic at all scales computed; the <i>diagonal</i> quartic localizes monotonically ($\text{PR}/D \rightarrow 10^{-3}$); the <i>kinetic</i> completion is the only one that relaxes (occupation falls from K to ≈ 3.5 within $\tau \approx 20$, GOE statistics, delocalized eigenstates at matched dose), and its finite-time plateau equals the diagonal ensemble exactly where the latter is computable ($D \leq 10^4$, $\text{plateau} - \text{diag} /K \leq 0.005$); but neither scaling leg is positive: $\sigma_{\text{ETH}}/\text{std}$ is flat at 0.95–0.98 across $D = 3375 \rightarrow 59319$ at matched dose, and the diagonal ensemble stays 0.04–0.09 of K away from microcanonical with no closing trend. Strong dose ($VK^2 \gtrsim 80$) localizes. Level statistics come free and positive (GOE at generic coupling, C5); the weak-coupling $\langle r \rangle$ is quantile- and direction-dependent (0.42–0.56; Remark 3.9(a)), so PR/D is the robust weak-coupling diagnostic.

ID	Statement	Grade	Depends on	Falsifier / test route
C5	<i>Arithmetic–chaos correspondence.</i> The spectral statistics of the quantized near-singularity (BKL-type) Hamiltonian match GUE, and its spectral determinant is arithmetic: the Hilbert–Pólya operator for the zeta zeros [43–45].	B	C1 (int.)	Compute pair correlations of quantized cosmological billiards; a proven non-GUE spectrum for the singularity sector falsifies; conversely, any rigorous Hilbert–Pólya operator would confirm the bridge (and RH’s spectral side). On the lattice model itself: GOE ratio statistics at generic coupling, verified numerically for $D = 625$ to $D = 5.9 \times 10^4$ (eigen-windows; equilibration diagnostics to $D \approx 10^5$), with a Poisson control at weak coupling (Remark 3.9).
C6	<i>Hagedorn wall as singularity resolution.</i> Approach to a spacetime singularity is approach to T_H : energy converts to primon creation, curvature growth saturates (Remark 6.5).	B	C1, C3	Near-singularity GR calculations (e.g. BKL maps) exhibiting equilibration at a critical scale; falsified by derived continua with no critical behavior.
C7	<i>Janus boundary condition.</i> The universal state’s history touches the unique zero-entropy floor $ 1\rangle$ at one end (the arithmetic Past Hypothesis).	B	C1, C4	No direct route; constrained by C1, C4; cosmological tests indirect (initial-state smoothness statistics).
C8	<i>Two-sided entropy.</i> The entropy profile is U-shaped about $ 1\rangle$ (Janus-point cosmology, arithmetic form) [37, 38].	B	C7	As for C7; additionally, CMB statistics distinguishing one-sided from two-sided low-entropy origins.

ID	Statement	Grade	Depends on	Falsifier / test route
C9	<i>Galois–gauge dictionary.</i> Minimal form of Conj. 8.1: charge lattice $\leftrightarrow$ splitting restrictions; branching ratios $\leftrightarrow$ Chebotarev densities; hierarchy $\leftrightarrow$ ramification.	B	C1 (int.)	<i>Calibrated dictionary protocol</i> (restated): fit all channel ratios of ≥ 5 independent decay channels at precision $\sigma \leq 0.3\%$ with one common group order $N \lesssim \text{few} \times 10^2$ and class sizes $c_i \mid N$; Monte-Carlo-calibrated false-positive rate $< 1\%$. No acceptable common- N fit at that precision falsifies. Present data (e.g. Z-pole branching fractions) admit no such fit—the signature is absent below unification, as the conjecture requires. Cleanest dated falsifier: observed proton decay (Section 11, item 1).
C10	<i>CP violation as the σ-odd sector.</i> CKM phase and baryon asymmetry are the arithmetic index of asymmetric couplings $w(\pi) \neq w(\bar{\pi})$ (Prop. 8.5); the ratchet compounds multiplicatively along the cascade [36].	B	C4	Structural constraint: all derived CP/T observables must factor through split/inert statistics; any derivation forcing a non-arithmetic phase breaks it. The ratchet lives in the non-conserving H_W sector (Remark 3.6).
C11	<i>Arithmetic hierarchy.</i> $\log(M_{\text{Pl}}/M_{\text{EW}}) \asymp E_{\text{cut}}/2\hbar\omega_0$, i.e. $E_{\text{cut}} \asymp 74\hbar\omega_0$ (Conj. 6.2): the hierarchy is a cutoff question.	B	C1, C3	Any derivation fixing ω_0 to a known scale (e.g. a horizon temperature, C3) closes or breaks the numerology; conversely a derived 10^{32} -state effective arithmetic universe confirms.
C12	<i>Perspectival Everett semantics.</i> Branches are factorization histories in $\mathcal{H}_{\mathcal{A}}$; “now” and “I” are indexicals; the felt flow is record formation along the cascade [30, 42].	C	C4	Interpretive: tethered to C4; not separately falsifiable at present; internal criteria of coherence and derivative phenomenology only.

Three structural observations about the register. First, its *physical depth ordering* is $\text{C1} \Rightarrow \{\text{C2}, \text{C3}, \text{C5}, \text{C6}\} \Rightarrow \{\text{C4}, \text{C7}, \text{C8}\} \Rightarrow \{\text{C9}, \text{C10}, \text{C11}\} \Rightarrow \text{C12}$, with two cross-layer edges ($\text{C11} \leftarrow \text{C3}$: the hierarchy falsifier invokes C3’s horizon temperature; $\text{C10} \leftarrow \text{C4}$: the ratchet compounds along the cascade): the zeta/black-hole identification is the root, and its confirmation or collapse

propagates deterministically through the physical plane. But the register is *two-plane*: on the mathematical plane, C4, C5, and the computational content of C9 (find a number field matching a branching table) are autonomous problems that survive any verdict on C1—which is precisely why they carry the register’s most economical tests. C4’s listed dependency on C1 is interpretive, not logical: the arithmetic sector and $H_{\text{tot}}^{(U,V)}$ exist unconditionally, so the thermalization problem is well-posed even if the spectral-identification program collapses. Second, four entries (C1, C3, C4, C5) have *purely mathematical* test routes: they can be advanced or refuted without any experiment. Third, the register is conservative: no claim appears both as a theorem and as a graded conjecture; where a strong reading is available but unproved, only the graded conjecture is carried.

11 Falsifiability, Predictions, and Test Programs

11.1 What the framework predicts that rivals do not

1. *No proton decay, by construction.* The arithmetic GUT unifies through number fields, not through a larger Lie group; baryon number is not gauged into a decay channel. Absence of proton decay, a persistent embarrassment for SU(5)-type models, is here a structural feature. Conversely, observed proton decay would falsify C9 outright—a clean, dated test.
2. *Branching statistics are group statistics.* Under C9, unification-scale branching data must organize into Chebotarev densities: rational numbers with denominators dividing a Galois group order. This is a sharp, unusual, and cheap-to-check signature. The check is now calibrated: for ≥ 5 channels at precision $\sigma \leq 0.3\%$ with $|G| \lesssim$ a few hundred, a common-group-order divisor fit has Monte-Carlo-calibrated false-positive rate below 1% (the divisor constraint is what saves the test; the weak form—each ratio near some rational—is vacuous because $\mathbb{Q}$ is dense). Current tables (e.g. the Z -pole branching fractions, five channels) admit no such fit: the signature is absent below unification, exactly as C9 requires, and no obtainable data set can decide the conjecture today. The computation validates and calibrates the instrument; the purchase waits on unification-scale data.
3. *A single new scale.* The framework has exactly one dimensionful parameter, ω_0 , with $T_H = \hbar\omega_0/k_B$. Every other number in the physics layer is either an integer, a group order, or a value of an L -function. Any phenomenological use of the framework that requires a second free scale has already left the theory.
4. *Entropy log-corrections are sawtooth-bounded, not logarithmic.* The $O(1)$ defect in Theorem 6.1 oscillates and never converges; the framework’s black-hole entropy corrections are therefore not of the standard $-\alpha \log A$ family unless additional coarse-graining is applied. This is a distinctive statistical signature of integer-lattice counting (register C3’s falsifier).
5. *Recurrence phenomenology is layered and cutoff-controlled.* The framework’s cosmology distinguishes exact recurrence (forbidden for generic states, Theorem 4.2) from ε -recurrence (universal for the free flow, Theorem 4.4, but with return times growing without bound in the state), and identifies genuine transience only in the cascading dynamics with three or more active modes (Proposition 4.5). Any completed theory of the arithmetic cutoff must fix where our universe sits among these layers—a prediction in the sense that the cutoff mechanism cannot be indifferent.

11.2 Mathematical test programs

Four problems, in increasing order of depth, would each move multiple register entries:

1. *Prove or refute thermalization for $H_{\text{tot}}^{(U,V)}$ on finite truncations (C4)*: a numerical program of ETH tests along the prime lattice, directly analogous to established kicked-rotor and Bose–Hubbard studies; every ingredient is finite-dimensional and computable. The program has been carried out at three scales and for both completions: dense diagonalization to $D = 4096$, sparse shift-invert eigen-windows to $D = 5.9 \times 10^4$, and Krylov equilibration to $D \approx 10^5$ —all laptop-scale (minutes per configuration; the interior eigensolver tier is bounded by LU fill, the evolution tier only by $D\tau$; diagonal pivoting, needed for the kinetic family, cut the generic factorization from 257 s to 3 s with eigenpair residuals $\leq 6 \times 10^{-8}$). Verdict: level statistics GOE at generic coupling throughout; every *eigenstate* ETH diagnostic flat in D ($\sigma_{\text{ETH}}/\text{std} \approx 0.95$ for the bare, quartic, and kinetic systems alike, at matched dose across $D = 3375 \rightarrow 59319$); the diagonal quartic completion localizes monotonically in U ; the kinetic completion (density-assisted hopping, Hermitian and number-conserving, validated to 10^{-15}) is the only one that relaxes—fast initial decay to ≈ 3.5 within $\tau \approx 20$ with GOE statistics and delocalized eigenstates, and a finite-time plateau that equals the diagonal ensemble exactly where the latter is computable ($D \leq 10^4$)—but the diagonal ensemble itself stays 0.04–0.09 of K from the microcanonical value at matched dose with no closing trend, and the $D = 20736$ trajectory extended to $\tau = 300$ saturates at $\langle \hat{n}_d \rangle_{\text{diag}} \approx 4.40$ against a microcanonical value of 3.35. The strong-ETH leg is flat at every accessible scale; the equilibration leg requires the $D \rightarrow \infty$ limit of the diagonal ensemble, which shows no convergence at accessible scales.
2. *Compute the level statistics of the interacting arithmetic Hamiltonian and compare with GUE (C5)*: the prime lattice with log-spaced on-site energies is a banded, incommensurate system; its statistics are accessible by dense-diagonalization experiments and by Dyson–Brownian-motion arguments. The banded-incommensurate claim is verified numerically at every scale computed: GOE ratio statistics ($\langle r \rangle = 0.45\text{--}0.58$) from $D = 625$ to 5.9×10^4 at generic coupling, with the weak-coupling limit identifiable by $\text{PR}/D \rightarrow 10^{-4}$ —though whether the correspondence is GOE or GUE in the infinite limit is exactly what finite windows cannot decide ($\langle r \rangle$ separates GOE from GUE by only 0.06).
3. *Construct a Hilbert–Pólya candidate from the arithmetic sector*: the primon Hamiltonian itself has spectrum $\log n$, not the zeros; the zeros enter as resonances of the *fluctuation* sector $-\zeta'/\zeta$. A candidate operator whose spectrum is $\{\frac{1}{2} + i\gamma\}$ built from the walk/hop sector would connect the framework to the deepest open problem in the mathematics (C5, and with it the spectral reading of RH via von Koch’s equivalence [20]).
4. *Close the Chebotarev dictionary on one completed instance (C9)*: find a number field whose splitting statistics reproduce a measured branching table. Even one instance would convert the dictionary from analogy to theory; the search space is computable and finite for any fixed table.

11.3 Assessment

The framework now satisfies three standards. Its mathematical core is genuinely theorem-grade: the spectrum, the partition function, the Hagedorn asymptotics, the counting law, the clock kernel, the no-periodicity theorem, and the entanglement structure are proved. Its analogies are tightened: every structural analogy is either a citable theorem (Kronecker–Weber, Chebotarev, Erdős–Kac, Pólya, Bohr) or an explicit register entry. And its remaining ambition—that the primes order the spectrum of spacetime itself—is stated in a form precise enough to be wrong: C1 through C12 name their evidence, their dependencies, and their falsifiers, and the two numerically testable entries (C4, C9) carry carried-out programs with stated uncertainties. A framework that can be wrong in stated ways is a research program; one that cannot is a metaphor.

A Notation

Symbol	Meaning
Pr	rational primes; $\mathcal{P}_{\mathbb{Z}[i]}$ Gaussian primes modulo units
$\mathcal{F}$	free commutative monoid on Pr (occupation vectors)
$\mathcal{H}_{\mathcal{A}} = \ell^2(\mathcal{F})$	arithmetic Hilbert space (Def. 2.1)
$ n\rangle$	occupation state of the integer $n = \prod p^{k_p}$
$\hat{n}_p, \hat{N}_{\text{tot}}, \hat{N}$	mode occupation; total primon number; multiplicative number operator
$H_P = \hbar\omega_0 \log \hat{N}$	primon Hamiltonian (Def. 2.3)
$T_H = \hbar\omega_0/k_B$	Hagedorn temperature
M_n	multiplication operator, $ k\rangle \mapsto nk\rangle$ (Def. 3.2)
H_W, H_{hop}	walk and hop Hamiltonians; the prime lattice (Sec. 3.3)
$H_{\text{kin}}(V)$	kinetic completion $V \sum_{p < q} (\hat{n}_p + \hat{n}_q)(a_p^\dagger a_q + a_q^\dagger a_p)$ (Conj. 3.1)
$\zeta, L(\chi_{-4}), \zeta_{\mathbb{Q}(i)}$	Riemann zeta; Dirichlet $L \pmod{4}$; Dedekind zeta of $\mathbb{Q}(i)$
$\omega(n), \Omega(n)$	number of distinct / total prime factors
$N(E) = \lfloor e^{E/\hbar\omega_0} \rfloor$	state counting function (Thm 6.1)
$ \tau; \beta\rangle, K(\Delta\tau)$	zeta clock family and two-time kernel (Def. 5.1, Prop. 5.2)
C, P, T	arithmetic charge conjugation, parity, time reversal (Def. 8.4)
C1–C12	the conjecture register (Sec. 10)

References

- [1] Bernard L. Julia. Statistical theory of numbers. In *Number Theory and Physics*, volume 47 of *Springer Proceedings in Physics*, pages 276–293. Springer, Berlin, 1990.
- [2] Daniel Spector. Duality, partial supersymmetry, and arithmetic number theory. *Journal of Mathematical Physics*, 39:1913–1928, 1998.
- [3] Sean A. Hartnoll, Ming Yang, and Marine De Clerck. Conformal primon gases and Gaussian primes near black-hole singularities, 2026. Work reported in 2025–2026 popular and research coverage; see the source transcript for the chain of reports. Cited here as the physical provenance of the primon–black-hole link.
- [4] Sean Carroll. *From Eternity to Here: The Quest for the Ultimate Theory of Time*. Dutton/Penguin Group, New York, 2010.
- [5] David Wallace. The nature of the past hypothesis. In Khalil Chamcham et al., editors, *The Philosophy of Cosmology*, pages 486–499. Cambridge University Press, Cambridge, 2017.
- [6] Huw Price. *Time’s Arrow and Archimedes’ Point: New Directions for the Physics of Time*. Oxford University Press, 1996.
- [7] Benj Hellie. The vertiginous question, 2013. Preprint, PhilPapers.
- [8] Caspar Hare. Egocentric presentism and perspectival realism, 2009. Position papers as discussed in the source transcript.
- [9] Vincent Conitzer. First-person perspectives and the A-theory of time, 2019. Argument referenced as source item [33] of the source transcript.
- [10] Rolf Hagedorn. Statistical thermodynamics of strong interactions at high energies. *Nuovo Cimento Supplemto*, 3:147–186, 1965.
- [11] Edward C. Titchmarsh. *The Theory of the Riemann Zeta-function*. Oxford University Press, 2nd edition, 1986. Revised by D. R. Heath-Brown.

- [12] Jürgen Neukirch. *Algebraic Number Theory*. Springer, Berlin, 1999.
- [13] G. H. Hardy and Srinivasa Ramanujan. The normal number of prime factors of a number n . *Quarterly Journal of Mathematics*, 48:76–92, 1917.
- [14] Paul Turán. On a theorem of hardy and ramanujan. *Journal of the London Mathematical Society*, 9:274–276, 1934.
- [15] Paul Erdős and Mark Kac. The gaussian law of errors in the theory of additive number theoretic functions. *American Journal of Mathematics*, 62:738–742, 1940.
- [16] G. H. Hardy and E. M. Wright. *An Introduction to the Theory of Numbers*. Oxford University Press, 6th edition, 2008.
- [17] George Pólya. Über eine aufgabe der wahrscheinlichkeitstheorie betreffend die Irrfahrt im Straßennetz. *Mathematische Annalen*, 84:149–160, 1921.
- [18] Don N. Page and William K. Wootters. Evolution without evolution: Dynamics described by stationary observables. *Physical Review D*, 27:2885–2892, 1983.
- [19] Abram S. Besicovitch. *Almost Periodic Functions*. Cambridge University Press, 1932.
- [20] Helge von Koch. Sur la distribution des nombres premiers. *Acta Mathematica*, 24:159–182, 1901.
- [21] Jacob D. Bekenstein. Black holes and entropy. *Physical Review D*, 7:2333–2346, 1973.
- [22] Stephen W. Hawking. Particle creation by black holes. *Communications in Mathematical Physics*, 43:199–220, 1975.
- [23] Ted Jacobson. Thermodynamics of spacetime: the Einstein equation of state. *Physical Review Letters*, 75:1260–1263, 1995.
- [24] Erik Verlinde. On the origin of gravity and the laws of Newton. *Journal of High Energy Physics*, 2011:29, 2011.
- [25] Gerard 't Hooft. A planar diagram theory for strong interactions. *Nuclear Physics B*, 72: 461–473, 1974.
- [26] John F. Clauser, Michael A. Horne, Abner Shimony, and Richard A. Holt. Proposed experiment to test local hidden-variable theories. *Physical Review Letters*, 23:880–884, 1969.
- [27] John S. Bell. On the Einstein Podolsky Rosen paradox. *Physics*, 1:195–200, 1964.
- [28] Wojciech H. Zurek. Decoherence, einselection, and the quantum origins of the classical. *Reviews of Modern Physics*, 75:715–775, 2003.
- [29] Rolf Landauer. Irreversibility and heat generation in the computing process. *IBM Journal of Research and Development*, 5:183–191, 1961.
- [30] David Wallace. *The Emergent Multiverse: Quantum Theory according to the Everett Interpretation*. Oxford University Press, 2012.
- [31] G. B. Lesovik et al. Arrow of time and its reversal on an IBM quantum computer. *Scientific Reports*, 9:4396, 2019.
- [32] B. Hensen et al. Loophole-free Bell inequality violation using electron spins separated by 1.3 kilometres. *Nature*, 526:682–686, 2015.

- [33] Peter G. L. Dirichlet. Beweis des Satzes, dass jede unbegrenzte arithmetische Progression, deren erstes Glied und Differenz ganze Zahlen ohne gemeinschaftlichen Theiler sind, unendlich viele Primzahlen enthält. *Abhandlungen der Königlichen Akademie der Wissenschaften zu Berlin*, pages 45–71, 1837.
- [34] Nikolai G. Chebotarev. Die Bestimmung der Dichtigkeit einer Menge von Primzahlen, welche zu einer gegebenen Substitutionsklasse gehören. *Mathematische Annalen*, 95:191–228, 1926.
- [35] Laurent Lafforgue. Chtoucas de Drinfeld et correspondance de Langlands. *Inventiones Mathematicae*, 147:57–462, 2002.
- [36] Joan A. Vaccaro. The arrow of time and the nature of spacetime, 2016. arXiv:1510.08727.
- [37] Sean M. Carroll and Jennifer Chen. Spontaneous inflation and the origin of the arrow of time. *arXiv preprint hep-th/0410270*, 2004.
- [38] Julian Barbour, Tim Koslowski, and Flavio Mercati. Janus points and arrows of time. *arXiv preprint arXiv:1601.03243*, 2016.
- [39] Julian Barbour, Tim Koslowski, and Flavio Mercati. Identification of a gravitational arrow of time. *Physical Review Letters*, 112:101302, 2014.
- [40] David Malament. Causal theories of time and the conventionality of simultaneity. *Noûs*, 11:293–300, 1977.
- [41] Rafael D. Sorkin. Causal sets: discrete gravity, 2003. arXiv:gr-qc/0309009.
- [42] David Kaplan. Demonstratives. In *Themes from Kaplan*, pages 481–563. Oxford University Press, 1989.
- [43] Michael V. Berry and Jonathan P. Keating. The Riemann zeros and eigenvalue asymptotics. *SIAM Review*, 41:236–266, 1999.
- [44] Hugh L. Montgomery. The pair correlation of zeros of the zeta function. In *Analytic Number Theory*, volume 24 of *Proceedings of Symposia in Pure Mathematics*, pages 181–193. American Mathematical Society, 1973.
- [45] Andrew M. Odlyzko. On the distribution of spacings between zeros of the zeta function. *Mathematics of Computation*, 48:273–308, 1987.