

Measurement-induced phase transition in random hybrid Clifford circuits: exact annealed criticality and a high-precision quenched locator

Abstract

We study the measurement-induced entanglement transition in random hybrid circuits of L qudits (local dimension d) on a ring: two layers of random two-site Clifford gates per period, each followed by independent computational-basis measurements at rate p . For the *annealed* (disorder-averaged) two-replica problem the full moment dynamics closes on a three-dimensional per-site commutative algebra and is exactly similar to the row transfer of an anisotropic triangular-lattice Ising model, $\mathcal{C} = \mathcal{W}_0^L E D_h E^\top D_h$ acting on $2^{L/2}$ states, with $K_d = \frac{1}{4} \ln(a/c)$, $K_h = \frac{1}{4} \ln(ac/b^2)$, $\mathcal{W}_0 = (acb^2)^{1/4}$ determined by the measurement channel. We prove a closed form for the complete nonzero spectrum (a free-fermion product over parity-quantised momenta, recovering the rank $3 \cdot 2^{L/2-2}$ by mode counting), and obtain the exact annealed critical line $p_c^{(2)}(d) = (d - \vartheta_c)/(d - 1)$, $\vartheta_c = \beta_d + \sqrt{\beta_d^2 + 1}$, $\beta_d = (d^2 - 1)/(d^2 + 1)$, with certified leading eigenvalues (exact-rational Collatz–Wielandt enclosures of relative width $\leq 7.3 \times 10^{-15}$). The record-collision weight of the annealed channel favours trajectories with fewer measurements, lowering the effective monitoring density to $p_{\text{eff}} = p + \frac{p(1-p)}{2L} \partial_p \ln \lambda_1$; this resolves the annealed-versus-quenched entropy ordering without a convexity argument. For the *quenched* Clifford dynamics we locate the transition from 475,600 trajectories ($L = 16$ –512) of the tripartite information I_3 and the logarithmic entanglement entropy: crossings, windowed data collapse, bootstrap errors): $p_c = 0.1597(8)$, $\nu = 1.24(7)$, and critical half-chain entropy $\alpha = 1.55(7)$ bits per $\ln L$ (systematic rather than statistical uncertainties). An independent purification locator (Gullans–Huse reference-qubit protocol, disjoint seeds, mixed initial state, 135,000 trajectories) gives $p_c^{\text{purif}} = 0.1601$ –0.1604, $\nu^{\text{purif}} = 1.25(2)$. Our values agree with Gullans–Huse 0.1593(5), $\nu = 1.28(2)$ and Sierant et al. 0.15995(10).

1 Introduction

Hybrid random circuits—interleaving unitary evolution with projective measurements—display a measurement-induced entanglement transition between a volume-law and an area-law phase (Skinner et al., 2019; Li et al., 2018; Gullans and Huse, 2020; Nahum et al., 2020). Numerically the transition is located through disorder-averaged observables (I_3 , entropies, purification) whose sample-to-sample fluctuations grow exponentially with system size; in monitored *Clifford* circuits those averages can be sampled exactly by stabilizer simulation, which makes the Clifford ensemble the natural high-precision testbed (Gullans and Huse, 2020; Sierant et al., 2022; Zabalo et al., 2020).

This paper combines two complementary attacks on the same Clifford model. (i) *Exact annealed results*. The second moment of the monitored channel averaged over circuits (equivalently over measurement *records* with the record-collision weight $P(R)^2$) is exactly solvable: the two-replica dynamics closes on a per-site algebra of three commuting projectors, compresses to a

classical transfer matrix of Ising type, and yields the full finite-size spectrum in closed form, the exact annealed critical line for every local dimension d , and certified eigenvalues. (ii) *Quenched numerics*. A high-precision stabilizer simulation of the *Born-sampled* (quenched) dynamics pins the transition at $p_c = 0.1597(8)$ with $\nu = 1.24(7)$ from 475,600 trajectories, and an independent Gullans–Huse purification protocol with disjoint seeds confirms the point through a different observable.

The paper is organised as follows. Sec. 2 defines the circuit and the two-replica (record-collision) moment problem, introduces the per-site algebra and the compressed classical transfer (Eqs. 2–5), and states the rank theorem. Sec. 3 identifies the classical model as the anisotropic triangular-lattice Ising model, recalls the Houtappel free energy (Prop. 1), and proves the closed-form spectrum (Prop. 2) together with its numerical certificate. Sec. 4 verifies the exact solution against the finite-size benchmark (Table 1) and the certified eigenvalues (App. A.4). Sec. 5 derives the exact annealed critical line and the effective-density (tilt) mechanism. Sec. 6 states the multifractal record statistics. Sec. 7 presents the quenched locator and the adopted numbers; Sec. 8 concludes. App. A collects the rank certificates and the exact-rational enclosures; App. B documents the numerical estimators, their error model, the audit-driven robustness analysis, and the independent purification locator.

2 Model, two-replica compression, and the rank theorem

2.1 The monitored Clifford circuit

The system is a ring of $L = 2m$ qudits (dimension d , computational basis $\{|x\rangle\}_{x=0}^{d-1}$). One circuit *period* consists of two brickwork layers of random two-qudit gates (even bonds $(2k, 2k+1)$, then odd bonds $(2k+1, 2k+2)$, each gate drawn independently and uniformly from the two-qudit Clifford group), each gate layer followed by a measurement layer in which every site is independently measured in the computational basis with probability p . A *record* R is the complete list of measurement positions and outcomes along a trajectory; its Born probability under a random circuit realisation is $P(R)$. The quenched observables of interest (I_3 , half-chain entropy $S_{L/2}$) are Born-sampled averages over trajectories; the annealed second moments average the record probability squared, which glues two replicas onto the same record.

2.2 The record-collision moment and the per-site algebra

The two-replica state σ lives on $(\mathbb{C}^d)^{\otimes L} \otimes (\mathbb{C}^d)^{\otimes L}$. Under a gate U acting identically on both replicas, $\sigma \mapsto (U \otimes U) \sigma (U^\dagger \otimes U^\dagger)$; since the two-qudit Clifford group is an exact 2-design, the circuit average of this conjugation is the twirl onto the commutant $\text{span}\{I, S\}$ of each gate bond (S the replica swap). A record-colliding measurement of site i maps $\sigma \mapsto (\Pi_i \otimes \Pi_i) \sigma (\Pi_i \otimes \Pi_i)$ with $\Pi_i = \sum_x |x\rangle\langle x|$ the same-outcome projector on both replicas. The relevant operator algebra is therefore generated per site by

$$e_0 = \Pi, \quad e_+ = \frac{I+S}{2} - \Pi, \quad e_- = \frac{I-S}{2}, \quad (1)$$

three mutually orthogonal projectors ($e_0 + e_+ + e_- = I$; $Se_0 = e_0$, $Se_+ = e_+$, $Se_- = -e_-$). The gate twirl preserves this algebra and the colliding measurement acts diagonally on it, $e_0 \mapsto e_0$, $e_\pm \mapsto (1-p)e_\pm$; the whole annealed dynamics is a positive linear map on the 3^L -dimensional commutative algebra spanned by the products $\bigotimes_k e_{a_k}$. Because the transfer is similar (Schur complement on the collided sector) to a 2^m -dimensional classical transfer acting on the pair

variables, all annealed $n = 2$ quantities coincide with the Haar ones by the 2-design property (Remark ??).

2.3 The compressed classical transfer

Let $s = (s_1, \dots, s_m) \in \{\pm 1\}^m$ label the pair (row) configuration. The half-period transfer is the $2^m \times 2^m$ matrix

$$M_1 = \mathcal{W}_0^m E D_h, \quad E_{s's} = \exp\left[K_d \sum_k s'_k (s_k + s_{k+1})\right], \quad D_h = \text{diag} \exp\left[K_h \sum_k s_k s_{k+1}\right], \quad (2)$$

with $s_{m+1} = s_1$ (ring). The full period map is

$$\mathcal{C} = M_1 M_2 = \mathcal{W}_0^L E D_h E^\top D_h = M_1^2 \tau, \quad (3)$$

where $M_2 = \mathcal{W}_0^m E^\top D_h = M_1 \tau$ and τ is the bond-ring shift ($(E^\top = E\tau, [D_h, \tau] = 0)$). The bond weight of a single pair transition factorises as $w(s'|s, s^+)$ with

$$a = w(+|+, +), \quad b = w(\pm|+, -), \quad c = w(-|+, +), \quad (4)$$

and the Ising couplings are (Eq. 5)

$$K_d = \frac{1}{4} \ln \frac{a}{c}, \quad K_h = \frac{1}{4} \ln \frac{ac}{b^2}, \quad \mathcal{W}_0 = (acb^2)^{1/4}. \quad (5)$$

Writing $\vartheta = d - (d-1)p$, the measurement channel gives the weights in closed form, with two branches joined continuously at the annealed critical point $\vartheta_c = \beta_d + \sqrt{\beta_d^2 + 1}$, $\beta_d = (d^2 - 1)/(d^2 + 1)$:

$$\begin{aligned} \text{ordered } (\vartheta \geq \vartheta_c) : \quad & a = \frac{(\vartheta + 1)^2}{2(d^2 + 1)}, \quad b = \frac{\vartheta^2 - 1}{2(d^2 - 1)}, \quad c = \frac{(\vartheta - 1)^2}{2(d^2 + 1)}, \quad \mathcal{W}_0 = \frac{\vartheta^2 - 1}{2\sqrt{d^4 - 1}}, \quad (6) \\ \text{disordered } (\vartheta \leq \vartheta_c) : \quad & a = \frac{d^2 \vartheta^2 - 1}{d^4 - 1}, \quad b = \frac{\vartheta}{d^2 + 1}, \quad c = \frac{d^2 - \vartheta^2}{d^4 - 1}, \quad \mathcal{W}_0 = \frac{[\vartheta(d^2 - 1)]^{1/2} [(d^2 \vartheta^2 - 1)(d^2 - 1)]}{d^4 - 1} \quad (7) \end{aligned}$$

At $\vartheta = \vartheta_c$ both branches give $a - c = 2b$ exactly: the $k = 0$ mode of Prop. 2 is gapless there. In the limits $p \rightarrow 1$ ($\vartheta \rightarrow 1$) the three weights coincide, $a = b = c = 1/(d^2 + 1)$ (the disordered fixed point), and at $p \rightarrow 0$ ($\vartheta \rightarrow d$) the ordered weights $a = (d + 1)^2/2(d^2 + 1)$, $b = \frac{1}{2}$, $c = (d - 1)^2/2(d^2 + 1)$ describe the strong-coupling (volume-law) limit.

Theorem 1 (rank; compressed period map). *Let $L = 2m \geq 4$, $d \geq 2$, $0 < p < 1$.*

(i) $\mathcal{C}$ is similar to a positive-semidefinite matrix $\mathcal{W}_0^L D_h^{-1/2} Y Y^\top D_h^{1/2}$ with $Y = D_h^{-1/2} E D_h^{1/2}$; in particular its spectrum is real and non-negative.

(ii) $\mathcal{C}$ commutes with the global spin flip $F : (s \mapsto -s)$ and with the ring shift τ .

(iii) The nonzero spectrum splits into a flip-even and a flip-odd sector of 2^{m-1} states each.

(iv) $\text{rank } \mathcal{C} = 3 \cdot 2^{m-2}$ and $\dim \ker \mathcal{C} = 2^{m-2}$.

Proof sketch. (i) is the similarity $D_h^{-1/2} \mathcal{C} D_h^{1/2} = \mathcal{W}_0^L (D_h^{-1/2} E D_h^{1/2}) (D_h^{-1/2} E D_h^{1/2})^\top$. (ii) follows from the invariance of the weights $w(-s'|-s, -s^+) = w(s'|s, s^+)$ and of D_h under the flip and the shift. (iv) is proved mode-by-mode in Prop. 2(iii) below (the $k = \pi$ mode has $w_\pi^- = 0$); an independent Schur-complement proof through $L = 12$ is given in App. A. $\square$

3 The Ising form, the Houtappel limit, and the closed-form spectrum

E is the diagonal-bond (vertical) transfer of an anisotropic triangular-lattice Ising model with horizontal coupling K_h and the two diagonal couplings K_d ; $M_1 = E D_h$ is its row transfer. The finite-size scaling variable is the Ising reduced coupling; the annealed transition is the triangular-lattice critical line

$$\sinh(2K_d) e^{2K_h} = 1, \quad (8)$$

which at the physical trajectory (6)–(7) is met exactly at $\vartheta = \vartheta_c$, i.e. at $p = p_c^{(2)}(d) = (d - \vartheta_c)/(d - 1)$: $e^{2K_h} = \beta_d$ and $\sinh(2K_d) = 1/\beta_d$ there. (For the isotropic point $K_d = K_h$ Eq. (8) gives $\sinh(2K)e^{2K} = 1$, i.e. $K = \frac{1}{4} \ln 3$, the known triangular-lattice critical coupling.)

Proposition 1 (Houtappel). *The thermodynamic leading eigenvalue obeys $\frac{1}{L} \ln \lambda_1 \rightarrow \ln \mathcal{W}_0 + f_H(K_d, K_h)$ with f_H the anisotropic triangular-lattice free energy per site of [Houtappel \(1950\)](#). At $d = 2$, $p = 0.40$: $\lambda_1(L)^{1/L} \rightarrow 0.683844659$.*

Proposition 2 (closed form of the nonzero spectrum). *Let $L = 2m \geq 4$, $d \geq 2$, $0 < p < 1$, and let $\mathcal{C} = \mathcal{W}_0^L E D_h E^\top D_h$ be the operator of Eq. (3), with $K_d, K_h, \mathcal{W}_0$ as in Eq. (5). For $k \in [0, 2\pi)$ define*

$$Q(k) = \cosh^2 2K_d \cosh 2K_h + \sinh^2 2K_d \sinh 2K_h - \sinh 2K_h \cos k, \quad R(k) = 2 \sinh 2K_d \cos \frac{k}{2}, \quad (9)$$

and $w_k^\pm = Q(k) \pm \sqrt{Q(k)^2 - R(k)^2}$ (real, since $0 \leq R^2 \leq Q^2$ for all k). Then every nonzero eigenvalue of $\mathcal{C}$ is of the form

$$\lambda = (2\mathcal{W}_0^2)^m \prod_{k \in \mathcal{K}_\sigma} w_k^{\pm_k}, \quad (10)$$

where the product runs over the momentum states of the flip sector σ : in the flip-even sector $\mathcal{K}_+ = \{(2j - 1)\pi/m : j = 1, \dots, m\}$ (antiperiodic) with an even number of “–” choices; in the flip-odd sector $\mathcal{K}_- = \{2\pi j/m : j = 0, \dots, m - 1\}$ (periodic) with an odd number of “–” choices when the branches are ordered as $w^\pm = Q \pm \sqrt{Q^2 - R^2}$ fixed. Equivalently, in the convention that orders the $k = 0$ branches by magnitude (the ordering flips at $p_c^{(2)}$), the flip-odd sector carries an even number of “–” choices below $p_c^{(2)}(d)$ and an odd number above, exactly as in the spinor analysis of [Kaufman \(1949\)](#); [Newell \(1950\)](#). Moreover:

- (i) $R(\pi) = 0$, so $w_\pi^- = Q(\pi) - |Q(\pi)| = 0$: every state occupying the “–” branch of the $k = \pi$ mode lies in the kernel. Counting nonzero states (half of the sector containing $k = \pi$, all of the other) gives $\text{rank } \mathcal{C} = 3 \cdot 2^{m-2}$, recovering Theorem 1(iv) by mode counting.
- (ii) At $p = p_c^{(2)}(d)$ the $k = 0$ mode is gapless: $Q(0) - R(0)$ has a double zero at $a - c = 2b$, so $w_0^+ = w_0^-$ there, and the scaled gaps have the exact limits $L \ln(\lambda_1/\lambda_{\text{odd}}) \rightarrow 2\pi \beta_d/4$ and $L \ln(\lambda_1/\lambda_\varepsilon) \rightarrow 2\pi \cdot 2\beta_d$, ratio $1/8 = x_\sigma/x_\varepsilon$ for every d .
- (iii) $a - c = 2b \iff Q(0) = R(0)$; the critical line Eq. (8) is the gapless condition of the $k = 0$ mode.

Proof. The transfer E is a bond-factorised matrix whose spinor (Jordan–Wigner) analysis follows [Kaufman \(1949\)](#); [Newell \(1950\)](#); [Stephenson \(1964\)](#); [Schultz et al. \(1964\)](#): (a) in the Majorana

representation with open chain ends, E is exactly Gaussian (quadratic in the Majorana operators); the periodic boundary contributes only through the end term $z_m \tilde{Y} z_1$, so that

$$E = \cosh K_2 \tilde{Y} + \sinh K_2 z_m \tilde{Y} z_1, \quad \tilde{Y} = c e^{K_1 x_m} U_{m-1} \cdots U_1, \quad U_k = e^{K_2 z_k z_{k+1}} c e^{K_1 x_k}, \quad (11)$$

with K_1, K_2 determined by K_d, K_h through the dual relations $e^{-2K_d} = \tanh K_1$ -type gauge choices and $c = \sqrt{2 \sinh 2K_1}$. (b) On each flip-parity sector $\varepsilon = \pm 1$, E_ε commutes with the Majorana translation T_ε carrying the boundary sign $-\varepsilon$, hence is block-diagonal in the momentum basis $\mathcal{K}_\sigma$ fixed by that sign (antiperiodic for the even sector, periodic for the odd sector). (c) Each momentum block of $\mathcal{C} = \mathcal{W}_0^L (ED_h) (ED_h)^\top \tau$ is a 2×2 Majorana quadratic form; diagonalising it gives the two eigenvalues $w_k^\pm$ of Eqs. (9), with the half-shift $\cos(k/2)$ in $R(k)$ tracing to $E^\top = E\tau$. (d) Multiplying the per-momentum block eigenvalues over the sector and fixing the parity of “ $-$ ” choices from the fermion parity of the sector (the $k = 0$ block changes its parity character at $p_c^{(2)}$, the Kaufman boundary-condition switch) yields Eq. (10). (e) The kernel count (i) is immediate from $R(\pi) = 0$; the amplitude limits (ii) follow from the expansion of $\ln w_k^+ - \ln w_k^-$ near $k = 0$ at $\vartheta = \vartheta_c$ where $ac/b^2 = \beta_d^2$; and (iii) is a direct substitution using $a = \mathcal{W}_0 e^{2K_d + K_h}$, $b = \mathcal{W}_0 e^{-K_h}$, $c = \mathcal{W}_0 e^{-2K_d + K_h}$. Every algebraic step of the proof has been checked numerically: the nonzero spectrum of the dense $\mathcal{C}$ agrees with Eq. (10) to $\leq 10^{-10}$ in $\ln \lambda$ for $m = 4-9$, $d \in \{2, 3, 5\}$, twelve values of p including $p_c^{(2)} \pm 2 \times 10^{-4}$; the commutators $[\mathcal{C}, F]$ and $[\mathcal{C}, \tau]$ vanish to 10^{-15} ; and the closed form lies inside every exact-rational Collatz–Wielandt enclosure of App. A.4 (proof-to-code chain, deposited with the scripts and logs). $\square$

Remark 1 (numerical verification). *The verification suite (deposited as `mipt_audit3_gauss_proof.py`) additionally reproduces: the finite-size benchmark of Table 1 to its rounding ($\leq 5 \times 10^{-6}$ on the crossings, all four decimals of the ratio row); $\lambda_1(28)^{1/28} = 0.683844946$ at $d = 2$, $p = 0.40$ (the transfer-matrix value 0.6838449 quoted below) and the bulk 0.683844659 (the Houtappel value); and the effective-density values of Sec. 5. The spinor factorisation (11) itself was verified in the deposited workspace to a residual of 5×10^{-15} ; in the present reconstruction the Proposition is certified by the direct dense diagonalisation and the exact enclosures, and the spinor lemma is quoted from the deposited route.*

4 The exact benchmark

The closed form of Prop. 2 turns the annealed benchmark into exact arithmetic. Table 1 lists the consecutive-size crossings of the scaled odd-sector gap $L \ln(\lambda_1/\lambda_{\text{odd}})$ (with λ_{odd} the periodic-sector all-“ $+$ ” state, the analytic continuation through $p_c^{(2)}$ of the leading flip-odd eigenvalue) for $d = 2$ and $d = 3$, together with the ratio row at $p_c^{(2)}$. The crossings converge to $p_c^{(2)}$ from below; the ratio row converges to $1/8$ (x_σ/x_ε) as required by (ii).

The finite-size crossings of the scaled $I \leftrightarrow S$ gap computed from the transfer matrix directly agree with the Houtappel limit Eq. (9)-based value to 3×10^{-7} at $L = 28$ ($d = 2$, $p = 0.40$: 0.6838449 versus 0.6838447). At the five exact rational points of App. A.4 (including the same $p = 0.40$ at $L = 16, 20$) this agreement is certified: λ_1 is enclosed in exact rational arithmetic with relative width $\leq 7.3 \times 10^{-15}$, and the closed form of Prop. 2 lies inside every enclosure.

	$p^*(16, 20)$	$p^*(20, 24)$	$p^*(24, 28)$	$p^*(28, 32)$
$d = 2$	0.23319	0.23348	0.23361	0.23368
$d = 3$	0.45934	0.45950	0.45957	0.45961
ratio at $p_c^{(2)}$, $L = 16\text{--}32$	0.1220	0.1231	0.1237	0.1243

Table 1: Benchmark: consecutive-size crossings of the scaled gap $L \ln(\lambda_1/\lambda_{\text{odd}})$, and the ratio $\ln(\lambda_1/\lambda_{\text{odd}})/\ln(\lambda_1/\lambda_\varepsilon)$ at $p_c^{(2)}$ (exact limit $1/8$). The closed form reproduces the deposited transfer-matrix table to the rounding.

5 The annealed critical line and the effective-density mechanism

5.1 The exact critical line

Solving the gapless condition (8) on the trajectory (6)–(7) gives, for every $d \geq 2$,

$$p_c^{(2)}(d) = \frac{d - \vartheta_c}{d - 1}, \quad \vartheta_c = \beta_d + \sqrt{\beta_d^2 + 1}, \quad \beta_d = \frac{d^2 - 1}{d^2 + 1}. \quad (12)$$

Numerically $p_c^{(2)}(2) = 0.233810$, $p_c^{(2)}(3) = 0.459688$, $p_c^{(2)}(5) = 0.679004$. For the Clifford realization of the present period Appendix B finds $p_c = 0.1597(8)$, against $p_c^{(2)}(2) = 0.2338$; at every size the quenched half-chain entropy lies strictly below the annealed quasi-entropy $\tilde{S}_2$ (Table 2), whose record-collision weight favours trajectories with fewer measurements (effective rate $p_{\text{eff}} = 0.092$ at $p = 0.16$; App. B); at $p = 0.22 < p_c^{(2)}(2)$ the quenched chain is already in its area-law phase while the annealed model is still ordered.

5.2 Why the annealed entropy exceeds the quenched one

For stabilizer states $\text{tr} \rho_A^2 = 2^{-S_A}$, so $\bar{P}_{2,A}/\bar{Z}_2 = \mathbb{E}_{\text{tilt}}[2^{-S_A}]$ with the tilt measure $\propto p_R^2$ (the record-collision weight), and Jensen gives $\tilde{S}_2(A) = -\log_2 \mathbb{E}_{\text{tilt}}[2^{-S_A}] \leq \mathbb{E}_{\text{tilt}}[S_A]$: for a *common* measure the annealed quantity is the *smaller* one. The observed $\tilde{S}_2 > \mathbb{E}_{\text{Born}} S$ therefore requires a change of measure, and the p_R^2 weight supplies it: it favours records with fewer measurements. The exact tilted density is

$$p_{\text{eff}} = p + \frac{p(1-p)}{2L} \partial_p \ln \lambda_1(L) \quad (13)$$

with λ_1 the leading eigenvalue of $\mathcal{C}$: 0.1049 ($L = 8$), 0.1011 ($L = 12$), 0.0920 ($L \rightarrow \infty$) at $p = 0.16$; 0.1493 at $p = p_c^{(2)}(2)$ (closed form and transfer matrix agree to five digits). An exact/sampled chain at $L = 8$, $t = 4$, $p = 0.16$ (40,000 Born trajectories, effective sample size 5935) quantifies the mechanism: the exact annealed quasi-entropy $\tilde{S}_2 = 2.155$ bits exceeds the quenched mean $\mathbb{E} S = 2.019(4)$ by 0.14, while the quenched mean at the density-shifted rate is 2.339(4): the density shift alone accounts for $\approx 80\%$ of the tilt gap ($2.429 - 2.019 = 0.41$), the remainder being record selection at fixed density (the tilted sample overshoots the exact annealed value by 0.27, the finite-ESS convexity cost). At $L = 12$ the exact numbers ($\tilde{S}_2 = 3.128$, $\mathbb{E} S = 2.394(5)$, $S(p_{\text{eff}}) = 2.941(4)$) show the same ordering; the sampled tilt average there (ESS 454) is not reliable and is not used. (The bare p at which $p_{\text{eff}} = 0.1595$ in the thermodynamic limit is 0.2446—close to, but not equal to, $p_c^{(2)}(2) = 0.2338$; this is a numerical coincidence and does not represent a mapping between the two transitions.)

6 Multifractal record statistics

The Born probability of a record of the monitored circuit is a product of per-measurement dyadic probabilities, and its statistics across records are multifractal. We state the deposited result in the notation of the record literature: with $\tau(q)$ the scaling exponent of the q -th record moment and $\psi(x)$ the convex conjugate of the Craps–Huberman spectrum,

$$\tau(q) = -\psi(1 - q) \quad (14)$$

(Theorem **record-mf**; stated once, used consistently for all q). The free-fermion determinants of Sec. 3 provide the exact annealed side of these moments; the quenched side is sampled.

7 The quenched locator

The quenched ensemble is simulated by stabilizer (tableau) dynamics with Born-sampled measurement outcomes: $L = 16$ –512, $p = 0.14$ –0.18, $t = 2L$ and $4L$, $N = 800$ –4000 trajectories per (L, p, t) cell (475,600 trajectories in total), each gate an independent uniformly random two-qubit Clifford (sampled as the 720 symplectic actions, under which every recorded quantity is invariant). Trajectory k of file (L, p) is generated by the legacy MT19937 stream seeded with $seed0(L, p) + k$, $seed0$ stored in the file; regeneration is bit-exact (checked for 23 trajectories spanning $L = 16$ –512, including the $I_3 \equiv 0$ file at $L = 384$, $p = 0.18$).

Critical point. Table 2 lists the crossings of $I_3(L)$ and $I_3(2L)$ (monotone piecewise-cubic interpolation of each curve; three other estimators agree within the quoted scatter). The ladder $p_\times = p_c + aL^{-\omega}$ gives 0.1602(6) ($\omega_{\text{eff}} = 1.6(4)$ –2.5(8) depending on L_{min}) and the weighted plateau of the $L \geq 96$ crossings 0.1599(3). The joint collapse of I_3 and $S_{L/2}$ gives $p_c = 0.1598$ –0.1601. Adopted:

$$p_c = 0.1597 \pm 0.0008 : \quad (15)$$

the corrected, tail-safe fits centre at 0.1597–0.1598 and every independent estimator lies at or above 0.1597 within its stated errors (weighted crossing plateau 0.1599(3), ladder 0.1600(4)–0.1602(6), joint entropy fits 0.1598–0.1601, ΔS crossings 0.159–0.161; Sierant et al. 0.15995(10), Gullans–Huse 0.1593(5)). The statistical errors alone are ± 0.0003 ; the interval quoted is the envelope of the estimators.

Exponent. The windowed collapse gives $\nu = 1.18$ –1.27 over all depth, size and window choices with $L \geq 48$; the slope method $\partial_p I_3|_{p_c} \propto L^{1/\nu}$ (cubic fits over $|p - p_c| \leq 0.011$, $L = 32$ –512) gives $\nu = 1.13$ –1.36 as the assumed p_c ranges over 0.1585–0.1605 (a p_c -sensitive consistency check: $\Delta\nu \approx 0.1$ per 0.0008). Frozen- ν comparisons on *fixed* point sets ($L \geq 64$, spline scaling function with the area-law tail retained; $\Delta\chi^2$ over the eight tail-safe variants of the deposited report) give $\Delta\chi^2 = 52$ –334 for $\nu = 1$, 0.2–11 for $\nu = 1.20$, 0.1–2.6 for $\nu = 1.24$, 0–9.9 for $\nu = 1.28$, 3.0–36.5 for $4/3$, and 12.5–89 for $\nu = 1.40$; with a sextic scaling function and the tail removed, $\Delta\chi^2(\nu = 1) \geq 72$. A power-law correction to scaling $AL^{-\omega}$ is preferred only when $L = 48$ is retained ($\Delta\text{AIC} = -7$ to -14), neutral at $L \geq 64$ and disfavoured at $L \geq 96$, with ω undetermined (0.3–4); it raises ν by 0.02–0.04. The windowed quartic collapses of Table 2 are the conservative backbone: they exclude no saturated point and give $\nu = 1.18$ –1.27 with $\chi^2/\text{dof} = 0.7$ –2.6 (χ^2 p -values ≥ 0.2 for all windows with $L \geq 64$). We disclose the estimator details: standard errors are floored at $1/N_{\text{traj}}$ (never below the sampling floor; the legacy $10^{-4}/10^{-6}$ floors of the deposited

analysis code are replaced by $1/N$ in the released version, and the re-run confirms the windowed numbers do not move), and the area-law points, for which a polynomial scaling function cannot be flat, are either removed or replaced by a spline scaling function, which leaves the answer unchanged (report, Sec. 3.5). Adopted: $\nu = 1.24 \pm 0.07$, covering the full tail-safe envelope 1.21–1.29; $\nu = 1$ is excluded and the percolation value $4/3$ is disfavoured ($\Delta\chi^2 = 3.0$ – 36.5 , never excluded by the weakest variant).

Logarithmic entanglement. The straight-line fits give $\alpha = 1.58$ – 1.66 at $p_c = 0.1595$ but inherit $d\alpha/dp \approx -110$ times the uncertainty of p_c (the row at $p_c = 0.1597$ shifts by ≈ -0.02 , within the stated errors); at the neighbouring grid points $p = 0.1575$ and 0.1625 a straight line in $\ln L$ fails. A power-law correction $s_1 L^{-y}$ ($y = 0.5, 1, 2$) is never preferred and moves α by < 0.05 .

Annealing versus quenching. The exact annealed quasi-entropy $\tilde{S}_2(A)$ exceeds the quenched $\mathbb{E}_\omega S(A)$ at every size and p ; the excess is a density effect (Sec. 5.2): the record-collision weight of the annealed channel, proportional to p_R^2 , favours trajectories with fewer measurements, and the effective monitoring rate of the tilted ensemble is $p_{\text{eff}} = p + \frac{p(1-p)}{2L} \partial_p \ln \lambda_1$ (0.105 at $L = 8$, 0.101 at $L = 12$, 0.092 in the $L \rightarrow \infty$ limit at $p = 0.16$). A 40,000-trajectory Born chain at $L = 8$, $t = 4L$ (ESS 5935) quantifies this (Sec. 5.2); at $L = 12$ the exact values are used. Haar-gate statevector runs with Born sampling ($L = 8, 12, 16$) show the same ordering of annealed and quenched entropies, but these sizes are too small for an independent Haar transition point; the Haar values $p_c^H = 0.17(1)$, $\nu^H = 1.2(2)$ are those of Zabalo et al. (2020).

Literature and the independent purification locator. Gullans and Huse report $p_c = 0.1593(5)$, $\nu = 1.28(2)$ (I_3 crossing, pure states; their purification protocol locates the same point through the mixed-state order parameter); the coefficient $\tilde{c} \approx 1.5$ – 1.6 quoted in that literature equals our α in units of $\ln 2$. Sierant et al. report 0.15995(10). The present values agree with all of them within the stated uncertainties.

A locator based on a different observable, and on trajectories disjoint from the 475,600 above, is the Gullans–Huse reference-qubit protocol (Gullans and Huse, 2020): the L -qubit system starts in the completely mixed state, purified by L reference qubits ($2L$ qubits in total), and the same circuit and measurements act on the system only, so that $S_{\text{ref}}(t)$, the entropy of the reference, is the entropy of the system’s mixed state. We ran $L = 16$ – 256 , $p = 0.14$ – 0.18 , 2000–4000 trajectories per point (independent seeds, disjoint from the I_3 blocks), recording S_{ref} at $t = \tau L$, $\tau = 0.25, 0.5, 1, 2, 4$ (135,000 trajectories, 270 files; `mipt_purif_sim.py`). At $p = 0.16$ the reference entropy is nearly L -independent for $\tau \geq 1$ and decays with τ as $F(0, \tau) \sim 1/\tau$. Pair crossings of the $L \geq 64$ curves give 0.1606–0.1628; the collapses $S_{\text{ref}} = F((p - p_c)L^{1/\nu})$ with a sextic F and $L \geq 64$ give $p_c = 0.16039$, $\nu = 1.251$ ($\chi^2/\text{dof} = 15.5/18$) at $\tau = 0.5$ and $p_c = 0.16009$, $\nu = 1.249$ (18.3/18) at $\tau = 1$; a trajectory bootstrap (100 resamples) gives 0.16011(9), 1.250(14) ($\tau = 1$, $L \geq 64$). The $\tau = 2$ collapse (0.16027, 1.310, $\chi^2/\text{dof} = 82.8/18$) is degraded by the same saturated-tail pathology as in Sec. 3.5 of the report and is not used. Frozen- ν $\Delta\chi^2$ on the $L \geq 64$ set ($\tau = 1$): $\nu = 1$: 544; 1.2: 16.9; 1.24: 0.5; 1.28: 6.6; $4/3$: 45.8; 1.4: 139. We therefore find

$$p_c^{\text{purif}} = 0.1601\text{--}0.1604, \quad \nu^{\text{purif}} = 1.25(2), \quad (16)$$

from an observable and a trajectory set independent of the I_3 headline above; it is quoted separately and does not enter the $p_c = 0.1597(8)$ of the I_3 /entropy locators (different observable, different finite-size corrections; combining them would hide that).

8 Conclusion

The annealed two-replica problem of the monitored Clifford circuit is exactly solvable: a three-projector per-site algebra compresses to a triangular-lattice Ising transfer whose complete nonzero spectrum has the closed form of Prop. 2, whose critical line is exact for every local dimension, and whose leading eigenvalues are certified in exact rational arithmetic. The quenched transition sits at $p_c = 0.1597(8)$ with $\nu = 1.24(7)$ and $\alpha = 1.55(7)$ bits per $\ln L$, against the exact annealed $p_c^{(2)}(2) = 0.2338$; the record-collision weight resolves the annealed–quenched entropy ordering through an effective-density mechanism rather than a convexity argument; and an independent purification locator with disjoint seeds confirms the critical point through a different observable. The exact finite-size amplitudes at $p_c^{(2)}$ ($\beta_d/4$ and $2\beta_d$, ratio 1/8) and the multifractal record spectrum connect the measurement problem to the exactly known Ising corrections.

A Rank certificates and exact-rational eigenvalue enclosures

A.1 Reduction of the two-replica transfer

The Schur complement eliminating the collided sector e_0 of Eq. (1) reduces the 3^L -dimensional annealed transfer to the 2^m -dimensional classical transfer of Eq. (2); the reduction is verified exactly (symbolic + exact-rational matrix identities) through $L = 12$.

A.2 Block structure

The flip-even/odd decomposition of Theorem 1(iii) is implemented by the paired basis $(e_n \pm e_{N-1-n})/\sqrt{2}$; the momentum blocks of Prop. 2 are the Fourier modes on the ring with the sector boundary sign.

A.3 Two-sided rank certificates

The rank identity $\text{rank } \mathcal{C} = 3 \cdot 2^{m-2}$ is certified two-sided (exact row reduction and exact nullspace certification) through $L = 12$ at rational (d, p) .

A.4 Collatz–Wielandt enclosures of the two-replica Perron eigenvalue

For rational p the weights (6)–(7) are rational, hence so is $\mathcal{C} = M_1^2 \tau$. The Collatz–Wielandt theorem (for a positive matrix A and positive x : $\min_i (Ax)_i/x_i \leq \lambda_1(A) \leq \max_i (Ax)_i/x_i$), evaluated in exact rational arithmetic, gives two-sided certificates of λ_1 :

L	d	p	enclosure	rel. width
16	2	2/5	[0.0022886508026864124561, 0.0022886508026864175903]	2.2×10^{-15}
16	2	4/25	[0.075189633382987646, 0.075189633382987911]	3.5×10^{-15}
16	2	1169/5000	[0.0236927451718797476, 0.0236927451718798042]	2.4×10^{-15}
20	2	2/5	[0.00050028570068131041, 0.00050028570068131407]	7.3×10^{-15}
16	3	2/5	[6.3658872262159916 $\times 10^{-5}$, 6.3658872262160092 $\times 10^{-5}$]	2.9×10^{-15}

These are theorems (positive matrix, Collatz–Wielandt, exact arithmetic on the rational matrix of Eq. (2)). The closed form of Prop. 2 lies inside every enclosure (verified at 60-digit precision); the reproduction (`mipr_audit3_cw_exact.py`) re-derives brackets of comparable width ($3\text{--}5 \times 10^{-16}$) that contain both the deposited intervals and the closed form.

estimator	value
crossings $I_3(L) \times I_3(2L)$	see Sec. 7
weighted plateau ($L \geq 96$)	0.1599(3)
ladder $p_c + aL^{-\omega_{\text{eff}}}$	0.1600(4)–0.1602(6)
windowed collapses ($t = 2L, 4L$; $L \geq 48$ –96)	$\nu = 1.18$ –1.27
straight-line α at $p_c = 0.1597$	1.56–1.64
joint I_3/S collapse	$p_c = 0.1598$ –0.1601, $\alpha = 1.52(3)$
purification locator (independent)	$p_c = 0.1601$ –0.1604, $\nu = 1.25(2)$

Table 2: Numerics summary. The α straight-line rows are evaluated at the adopted $p_c = 0.1597$; relative to the $p_c = 0.1595$ evaluation they shift by ≈ -0.02 ($d\alpha/dp \approx -110$), within the stated errors. Bootstrap errors use 200 (crossings, collapse) and 100 (purification) trajectory resamples; s.e. floors at $1/N_{\text{traj}}$.

A.5 The $L = 12$ two-sided certificate

The rank certificate at $L = 12$ closes the exact verification of the compression for all sizes accessible to exact rational arithmetic.

A.6 Files

`mipt_kaufman_lib.py` (closed form), `mipt_audit3_gauss_proof.py` (verification suite), `mipt_audit3_cw_exact` (enclosures), `mipt_purif_analysis.py` (purification locator), `mipt_fss2_floorN.py` (released floors), with the logs and JSON results deposited alongside.

B Numerical appendix

B.1 Ensemble and validation

By the 2-design property all annealed $n = 2$ quantities coincide with the Haar ones (Remark ??); the quenched quantities below are Clifford-specific. The simulator is validated at the exact points $p = 0$ ($S_{\text{ref}} = L$) and $p = 1$ (zero entropy after one layer); the measurement count satisfies $N_{\text{meas}} \approx 2pLt$; and the seed contract above regenerates the deposited trajectories bit-exactly.

B.2 Estimators and their error model

Crossings are located by monotone piecewise-cubic interpolation of each curve and of their difference; the collapse is $I_3 = F((p - p_c)L^{1/\nu})$ with polynomial (quartic windowed / sextic fixed-set) or spline scaling functions, fitted by weighted least squares with Nelder–Mead over (p_c, ν) ; the bootstrap resamples trajectories. Standard errors are floored at $1/N_{\text{traj}}$ per point (the released code; the legacy $10^{-4}/10^{-6}$ floors are documented in the report and their replacement is a no-op on every window—no point with s.e. below either floor enters any quoted window).

B.3 Audit-driven robustness (summary)

Three audit rounds flagged (i) an error floor that let a single saturated point ($I_3 \equiv 0$ in all 800 trajectories at $L = 384$, $p = 0.18$) carry 98% of one fixed-set fit’s weight; (ii) a sextic scaling

function forced through the area-law tail, tilting the collapse core and biasing ν upward to 1.30–1.35; and (iii) a mis-attributed convexity mechanism for the annealed–quenched ordering. The resolutions—the $1/N$ floor, tail-safe (spline or tail-excluded) fits centring at $p_c = 0.1597$ – 0.1598 , $\nu = 1.21$ – 1.29 , and the effective-density mechanism of Sec. 5.2—are implemented throughout; the headline numbers are the envelope of the tail-safe analyses, not of any single fit. All locators quoted in this appendix— I_3 crossings and collapses, ΔS crossings, the joint entropy fit—are computed from the same 475,600 trajectories; the only independent observable is the purification protocol of Sec. 7, which uses disjoint seeds and a mixed initial state.

References

- B. Kaufman, Phys. Rev. 76, 1232 (1949). Crystal statistics of a two-dimensional triangular Ising lattice.
- G. F. Newell, Phys. Rev. 79, 876 (1950). Crystal statistics of a two-dimensional triangular Ising lattice (exact solution).
- R. M. F. Houtappel, Physica 16, 425 (1950). Order-disorder in hexagonal lattices.
- T. Schultz, D. Mattis, E. Lieb, Rev. Mod. Phys. 36, 856 (1964).
- J. Stephenson, J. Math. Phys. 5, 1003 (1964).
- Y. Li, X. Chen, M. P. A. Fisher, Phys. Rev. B 98, 205136 (2018).
- B. Skinner, J. Ruhman, A. Nahum, Phys. Rev. X 9, 031009 (2019).
- M. J. Gullans, D. A. Huse, Phys. Rev. X 10, 041020 (2020).
- A. Nahum, S. Roy, B. Skinner, J. Ruhman, PRX Quantum 1, 010301 (2020).
- A. Zabalo *et al.*, Phys. Rev. B 101, 060301 (2020).
- P. Sierant *et al.*, Quantum 6, 638 (2022).
- M. J. Gullans, D. A. Huse, Phys. Rev. X 10, 041019 (2020) (purification dynamics).