

Week 10: Multinomial regression

Monica Alexander

18/03/2022

Packages:

```
library(tidyverse)
library(here)
library(nnet) # for multinomial
```

Multinomial

Question of interest: how does infant mortality cause of death vary by race, mother's age and prematurity?

Data prep

Read in infant data and do some cleaning:

```
d <- read_rds(here("data/infant.RDS"))
d <- d %>%
  mutate(neo_death = ifelse(aged<=28, 1, 0),
         cod_group = case_when(
           str_starts(cod, "peri") ~ "perinatal",
           cod %in% c("other", "unknown") ~ "other/unknown",
           cod %in% c("sids", "maltreatment", "infection") ~ "exogenous",
           cod %in% c("resp", "heart") ~ "respiratory/heart",
           TRUE ~ "congenital malformations"
         ),
         preterm = ifelse(gest<37, 1, 0)) %>%
  filter(gest<99, !is.na(mom_age_group))


infant <- d %>% select(race, mom_age, gest, preterm, cod_group)
infant
```

```
## # A tibble: 17,024 x 5
##   race mom_age gest preterm cod_group
##   <chr>   <dbl> <dbl>   <dbl> <chr>
## 1 NHW      30    27       1 perinatal
## 2 NHW      32    36       1 congenital malformations
## 3 NHW      25    44       0 perinatal
## 4 NHB      29    21       1 perinatal
## 5 NHB      23    26       1 perinatal
```

```
## 6 NHW      34    39      0 congenital malformations
## 7 NHB      27    26      1 perinatal
## 8 NHB      24    40      0 exogenous
## 9 NHB      39    36      1 congenital malformations
## 10 NHB     30    39      0 congenital malformations
## # ... with 17,014 more rows
```

Graph by race from lecture:

```
infant %>%
  group_by(race, cod_group) %>%
  tally() %>%
  group_by(race) %>%
  mutate(prop = n/sum(n)) %>%
  mutate(cod_group = fct_reorder(cod_group, -prop)) %>%
  ggplot(aes(cod_group, prop, fill = race)) + geom_bar(stat = "identity", position = 'dodge') +
  labs(title = "Proportion of infant deaths by cause", x = "cause", y = "proportion") +
  theme_bw() +
  scale_fill_brewer(palette = "Set1")
```



Making into wide format

```
infant_wide <- infant %>%
  group_by(race, mom_age, gest, preterm, cod_group) %>%
  tally(name = "deaths") %>%
  pivot_wider(names_from = cod_group, values_from = deaths) %>%
```

```
mutate_all(.funs = funs(ifelse(is.na(.), 0, .)))
head(infant_wide)
```

```
## # A tibble: 6 x 9
## # Groups:   race, mom_age, gest, preterm [6]
##   race mom_age gest preterm perinatal exogenous 'other/unknown'
##   <chr>   <dbl> <dbl>   <dbl>   <dbl>   <dbl>       <dbl>
## 1 NHB      14    19       1       1       0         0
## 2 NHB      14    21       1       1       0         0
## 3 NHB      14    22       1       1       0         0
## 4 NHB      14    23       1       1       0         0
## 5 NHB      14    24       1       3       1         1
## 6 NHB      14    25       1       1       0         0
## # ... with 2 more variables: congenital malformations <dbl>,
## #   respiratory/heart <dbl>
```

Making the Y variable:

```
infant_wide$Y <- as.matrix(infant_wide[,c("perinatal",
                                           "exogenous",
                                           "congenital malformations",
                                           "respiratory/heart", "other/unknown")])
head(infant_wide$Y)
```

```
##      perinatal exogenous congenital malformations respiratory/heart
## [1,]         1         0                     0                     0
## [2,]         1         0                     0                     0
## [3,]         1         0                     0                     0
## [4,]         1         0                     0                     0
## [5,]         3         1                     0                     0
## [6,]         1         0                     0                     0
##      other/unknown
## [1,]              0
## [2,]              0
## [3,]              0
## [4,]              0
## [5,]              1
## [6,]              0
```

Regression

```
library(nnet)
mod_mn <- multinom(Y ~ race+ mom_age+ preterm, data = infant_wide)
```

```
## # weights:  25 (16 variable)
## initial value 27399.071021
## iter  10 value 20149.661320
## iter  20 value 19437.349750
## final value 19436.462463
## converged
```

```
summary(mod_mn)
```

```
## Call:
## multinom(formula = Y ~ race + mom_age + preterm, data = infant_wide)
##
## Coefficients:
##              (Intercept)      raceNHW      mom_age      preterm
## exogenous              2.56320808  0.088345261 -0.05692035 -3.429460
## congenital malformations -0.01647076  0.621524245  0.01916732 -2.423940
## respiratory/heart        -0.15823646 -0.004845986 -0.01780013 -2.251658
## other/unknown            1.10771251  0.145290756 -0.02245255 -3.137589
##
## Std. Errors:
##              (Intercept)      raceNHW      mom_age      preterm
## exogenous              0.1235975  0.05354744  0.004365804  0.06000498
## congenital malformations  0.1093744  0.04840430  0.003501902  0.05449309
## respiratory/heart        0.1811151  0.07928810  0.006287607  0.08451183
## other/unknown            0.1361523  0.06022037  0.004717569  0.06546394
##
## Residual Deviance: 38872.92
## AIC: 38904.92
```

Pull out coefficients:

```
coef(mod_mn)
```

```
##              (Intercept)      raceNHW      mom_age      preterm
## exogenous              2.56320808  0.088345261 -0.05692035 -3.429460
## congenital malformations -0.01647076  0.621524245  0.01916732 -2.423940
## respiratory/heart        -0.15823646 -0.004845986 -0.01780013 -2.251658
## other/unknown            1.10771251  0.145290756 -0.02245255 -3.137589
```

```
exp(coef(mod_mn))
```

```
##              (Intercept)      raceNHW      mom_age      preterm
## exogenous            12.9773831  1.0923652  0.9446693  0.03240443
## congenital malformations  0.9836641  1.8617637  1.0193522  0.08857195
## respiratory/heart        0.8536479  0.9951657  0.9823574  0.10522463
## other/unknown            3.0274253  1.1563757  0.9777976  0.04338727
```

Exercise: plot coefficient estimates and standard errors.

Predicted probabilities

```
predict_df <- tibble(race = rep(c("NHW", "NHB"), each = 2),
                     mom_age = 30,
                     preterm = rep(c(0,1),2))
preds <- bind_cols(predict_df, as_tibble(predict(mod_mn, newdata = predict_df, type = 'probs')))
```

```
## # A tibble: 4 x 8
##   race mom_age preterm perinatal exogenous 'congenital malfor~ 'respiratory/he~
##   <chr>   <dbl>   <dbl>   <dbl>   <dbl>         <dbl>         <dbl>
## 1 NHW     30     0    0.110    0.282         0.357         0.0547
## 2 NHW     30     1    0.666    0.0555        0.192         0.0349
## 3 NHB     30     0    0.140    0.329         0.245         0.0700
## 4 NHB     30     1    0.740    0.0564        0.115         0.0390
## # ... with 1 more variable: other/unknown <dbl>
```

Plot:

```
preds %>%
  pivot_longer(`perinatal`:`other/unknown`, names_to = "cod_group", values_to = "probability") %>%
  mutate(preterm = ifelse(preterm==1, "pre-term", "full-term")) %>%
  ggplot(aes(race, probability, fill = cod_group)) +
  geom_bar(stat = "identity") +
  facet_grid(~preterm) +
  ggtitle("Predicted probabilities of infant death by race, prematurity and cause\nMothers aged 30")
```

