

SOC6707: Intermediate Data Analysis

Monica Alexander

Week 4: Data Visualization

Announcements

Data visualization principles

- ▶ Choose the right graph
- ▶ Know your audience
- ▶ Emphasize important patterns without being misleading
- ▶ Clear, effective designs

Choose the right graph

Choosing the right graph primarily depends on the type of variables that you are trying to visualize:


- ▶ Quantitative variables e.g. histograms, scatter plots
- ▶ Qualitative variables e.g. barcharts

Choose the graph based on the kind of data and the message to be conveyed.


- ▶ Do not use different graphs just for variety, as specific graphs convey certain types of information more effectively than others.
- ▶ If not required, do not use any chart — show only numbers.

Pie charts

Girl's names by starting letter, 1990



Girl's names by starting letter, 2010




Pie charts

?pie

Pie charts are a very bad way of displaying information. The eye is good at judging linear measures and bad at judging relative areas. A bar chart or dot chart is a preferable way of displaying this type of data.

Alternative



Know your audience

Graphs can be used for

- ▶ our own exploratory data analysis
- ▶ to convey a message to experts,
- ▶ to help tell a story to a general audience.

Make sure that the intended audience understands each element of the plot.

Examples: spiral plot, log scales


- ▶ Think of the color blind. In R, `viridis` and `brewer` palettes give colorblind-friendly options

Emphasize important patterns without being misleading

There is no such thing as information overload. There is only bad design. — Edward Tufte

- ▶ Eliminate distractions
- ▶ Highlight the essential
- ▶ Use color and text strategically
- ▶ Avoid pseudo-3D plots

Highlight the essential




Source:

<https://link.springer.com/article/10.1007/s11524-021-00573-8>

When to start the axis at zero?

When to start the axis at zero?



Source

When to start the axis at zero?



National Review


@NRO

Follow



The only [#climatechange](#) chart you need to see. natl.re/wPKpro


(h/t [@powerlineUS](#))



When to start the axis at zero?

Average global temperature by year


Data from NASA/GISS.




When to include zeroes

- ▶ With bar plots, we are implying the length is proportional to the quantities being displayed. By avoiding 0, relatively small differences can be made to look much bigger than they actually are.
- ▶ With line plots or plots that use position, it is not necessary to start the axis at zero (and could be misleading)

Life expectancy (years), 2010




Life expectancy (years), 2010



Emphasize important patterns without being misleading

COVID-19 CASES VS. DEATHS LAST 7 DAYS

DAILY CASES (7-DAY MOVING AVERAGE)



DEATHS (7-DAY DEATH RATE)




Source: CDC

Clear, effective designs

Life expectancy bounce-backs amid continued losses

Life expectancy changes since the start of the COVID-19 pandemic

Estimates for 2021 are adjusted for the weeks with missing data in 2021



Important types of graphs: next week

- ▶ Histograms
- ▶ Bar charts
- ▶ Boxplots
- ▶ Line plots
- ▶ Scatter plots

...but data visualization need not be a graph



Hobart, Tasmania, Australia



Toronto, Ontario, Canada

Important types of graphs

Important types of graphs

- ▶ Histograms
- ▶ Bar charts
- ▶ Boxplots
- ▶ Line plots
- ▶ Scatter plots

Example datasets used here


1. TTC subway delays (from above)
2. Country-level indicators, 2009-2017
 - ▶ Uploaded onto Quercus
 - ▶ TFR = total fertility rate
 - ▶ GDP = gross domestics product
 - ▶ dataset also has life expectancy (females), child mortality, maternal mortality

Histograms


Shows the distribution of a **quantitative** variable

- ▶ Histograms show the frequency (count) of observations by value
- ▶ The range of values of a variables is divided into intervals ('bins') and then the number of observations in each bin is tabulated
- ▶ A histogram shows the count of observations in each bin with a rectangle of height equal to the count
- ▶ The x axis is the value bins, the y axis is the count/frequency (or proportion)


Female life expectancy, 2017



Delay times, TTC subway 2019



Making the histogram more informative




Bar charts


Shows summary measures across values of a **categorical** (qualitative) variable

- ▶ Illustrate the value of a particular outcome in a particular category
- ▶ The 'value' can be counts, but could also be a summary measure (e.g. mean)
- ▶ The value is again shown by a rectangle of height equal to the value
- ▶ Bar charts can be plotted vertically or horizontally
- ▶ In the vertical setting, the x axis is the categories and the y axis is the value of the quantitative variable


Number of delays by line, 2019



Same but horizontal




Showing mean delay time



More complicated example

Top 5 station by median delay




Box plots

Good for showing summaries of **quantitative** variables across different **categorical** groups.


- ▶ Visualizing quartiles (25/50/75 percentiles) of quantitative data
- ▶ Boxes show the IQR and median
- ▶ Whiskers show values outside IQR (in R/ggplot, default is $1.5 \times \text{IQR}$)
- ▶ Outliers may be shown with individual dots
- ▶ In the vertical case, the x axis is the categories and the y axis is the quantitative variable

Life expectancy (years) by region of the world



Could also do horizontal

Life expectancy (years) by region of the world




Line plots

Best used to describe values of a **quantitative** variable (on y axis) across sequential values of another **quantitative** variable on the x axis

- ▶ Plots a series of values of a quantitative variable connected together by a line
- ▶ Useful to visualize trends over time

Total fertility rate, Canada and the US




Scatter plots


Shows relationship between two different **quantitative** variables

- ▶ Uses dots to represent values for two different **quantitative** values
- ▶ The position of each dot on the x and y axis indicates values for an individual data point
- ▶ Extremely useful in visualizing the relationship between two quantitative variables

TFR versus life expectancy, 2017



TFR versus life expectancy, 2017



Further introduction to ggplot


ggplot

- ▶ ggplot is the graphing package that goes with the tidyverse in R
- ▶ Very powerful to make a wide range of graphics
- ▶ Every graph so far this lecture was done in ggplot
- ▶ ggplot code works in layers, with each layer adding complexity
 - ▶ start with defining dataset and different variables
 - ▶ add on type of plot
 - ▶ scales
 - ▶ layout (facets)
 - ▶ themes, fonts, sizes...

More practice in lab, but here's a starting example

Reproducing the TFR versus life expectancy chart, colored by region

TFR versus life expectancy, 2017



Data

```
# read in the data
```

```
country_ind <- read_csv("../data/country_indicators.csv")
```

```
country_ind
```

```
## # A tibble: 1,584 x 9
```

```
##   country_code country      region      year   tfr life_expectancy child_mort
##   <chr>         <chr>      <chr>      <dbl> <dbl>         <dbl>      <dbl>
## 1 AFG          Afghanistan Southern Asia  2009  6.18          61.9       93.9
## 2 AFG          Afghanistan Southern Asia  2010  5.98          62.5       90.0
## 3 AFG          Afghanistan Southern Asia  2011  5.77          63        86.3
## 4 AFG          Afghanistan Southern Asia  2012  5.56          63.5       82.9
## 5 AFG          Afghanistan Southern Asia  2013  5.36          64.0       79.6
## 6 AFG          Afghanistan Southern Asia  2014  5.16          64.5       76.6
## 7 AFG          Afghanistan Southern Asia  2015  4.98          64.9       73.8
## 8 AFG          Afghanistan Southern Asia  2016  4.80          65.3       71.2
## 9 AFG          Afghanistan Southern Asia  2017  4.63          65.7       68.8
## 10 ALB         Albania      Developed re~ 2009  1.65          79.0       16.7
## # ... with 1,574 more rows, and 2 more variables: maternal_mort <dbl>,
## #   gdp <dbl>
```

```
# filter to just be 2017
```


```
country_ind_2017 <- country_ind %>% filter(year==2017)
```

A blank canvas

`aes` stands for aesthetic and tells `ggplot` the main characteristics of your plot (`x`, `y`, and if the color or fill vary by group)

```
plot1 <- ggplot(data = country_ind_2017, aes(x = tfr, y = life_expectancy))
```

```
#print  
plot1
```




Add the points

Add layers with ggplot using the +

```
plot1 <- ggplot(data = country_ind_2017, aes(x = tfr, y = life_expectancy)) +  
  geom_point()
```


plot1



Tidy up labels

```
plot1 <- ggplot(data = country_ind_2017, aes(x = tfr, y = life_expectancy)) +  
  geom_point()+  
  xlab("TFR (births per woman)") +  
  ylab("life expectancy (years)")
```


plot1



Title

```
plot1 <- ggplot(data = country_ind_2017, aes(x = tfr, y = life_expectancy)) +  
  geom_point()+  
  xlab("TFR (births per woman)") +  
  ylab("life expectancy (years)") +  
  ggtitle("TFR versus life expectancy, 2017")
```

plot1




Change color of points

to see all colors, type `colors()`

```
plot1 <- ggplot(data = country_ind_2017, aes(x = tfr, y = life_expectancy)) +  
  geom_point(color = "red")+  
  xlab("TFR (births per woman)") +  
  ylab("life expectancy (years)") +  
  ggtitle("TFR versus life expectancy, 2017")
```

plot1




Coloring by group

This goes in the `aes()` because it **depends on the data**

```
plot1 <- ggplot(data = country_ind_2017, aes(x = tfr, y = life_expectancy, color = region)) +  
  geom_point()+  
  xlab("TFR (births per woman)")+  
  ylab("life expectancy (years)")+  
  ggtitle("TFR versus life expectancy, 2017")  
  
plot1
```

TFR versus life expectancy, 2017




Change theme (optional) and size of points

```
plot1 <- ggplot(data = country_ind_2017, aes(x = tfr, y = life_expectancy, color = region)) +  
  geom_point(size = 2)+  
  xlab("TFR (births per woman)") +  
  ylab("life expectancy (years)") +  
  ggtitle("TFR versus life expectancy, 2017") +  
  theme_bw(base_size = 14)
```

plot1

TFR versus life expectancy, 2017




Change color scheme

viridis and brewer both good options

```
plot1 <- ggplot(data = country_ind_2017, aes(x = tfr, y = life_expectancy, color = region)) +  
  geom_point(size = 2) +  
  xlab("TFR (births per woman)") +  
  ylab("life expectancy (years)") +  
  ggtitle("TFR versus life expectancy, 2017") +  
  theme_bw(base_size = 14) +  
  scale_color_viridis_d()
```

plot1

TFR versus life expectancy, 2017



Summary

- ▶ EDA and data visualization is often just as informative and important as statistical analysis
- ▶ It is essential to understand the structure of your data, missing-ness, any outliers/issues, and the raw patterns in your data before deciding on your statistical analysis
- ▶ Plot, plot, plot
- ▶ Practice, practice, practice

Summary

Plots:

- ▶ Bar charts for categorical/qualitative variables
- ▶ Histograms, boxplots for one quantitative variable (potentially across multiple categories)
- ▶ Line plots and scatter plots for two quantitative variables (line plot when one is sequential)

Data ideas

- ▶ IPUMS: <https://ipums.org/>
- ▶ ICPSR:
<https://www.icpsr.umich.edu/web/pages/ICPSR/thematic-collections.html>
- ▶ CHASS SDA: <https://datacentre.chass.utoronto.ca/>
- ▶ Toronto Open Data Portal: <https://open.toronto.ca/> or use opendatatoronto R package (ask for code)
- ▶ UN WPP: <https://population.un.org/wpp/>
- ▶ NBER:
<https://www.nber.org/research/data?page=1&perPage=50>