



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



## Original short

## Influence of influenza vaccination and comorbidity on the evolution of patients hospitalised for COVID-19



José Manuel Fernández Ibáñez<sup>a,\*</sup>, María del Carmen Morales<sup>Ballesterosa</sup>, Manuel José Fernández<sup>Anguitab</sup>, María Ángeles Galindo<sup>Andúgard</sup>, Ángel Arias<sup>Arias</sup> and José Ramón<sup>Barberá-Farré</sup>

<sup>a</sup> Internal Medicine Department, Geriatrics Section, Hospital General Mancha Centro, Alcázar de San Juan, Spain.

<sup>b</sup> Cardiology Department, Hospital General Mancha Centro, Alcázar de San Juan, Spain.

<sup>c</sup> Research Unit, Hospital General Mancha Centro, Alcázar de San Juan, Spain.

<sup>d</sup> Internal Medicine Department, Hospital General Mancha Centro, Alcázar de San Juan, Spain.

## ARTICLE INFORMATION

## History of the article:

Received 25 February 2021

Accepted 2 June 2021 On-line 16 June 2021

## Keywords:

Influenza vaccine  
COVID-19  
Inpatients Age  
Comorbidity  
Mortality

## RESUMEN

**Background and objective:** The outbreak of COVID-19 disease is evolving worldwide. The aim of the study is to assess the association between influenza vaccination and the risk of mortality in hospitalised COVID-19 patients, as well as other risk factors.

**Materials and methods:** Retrospective observational study. It was conducted among hospitalised patients with COVID-19 at Hospital La Mancha Centro between 5 and 25 March 2020. We used multivariate logistic regression to explore the association between influenza vaccination and mortality due to COVID and other risk factors.

**Results:** 410 patients were included. Influenza vaccination had no effect among patients hospitalised for COVID-19 (OR 1.55 [95% CI 0.96–2.48,  $p=0.071$ ]). Increased in-hospital mortality was associated with older age (OR 1.05 [95% CI 1.02–1.07]), for each increase in age;  $p < 0.001$ , Charlson  $\geq 3$  (OR: 1.84 [95% CI: 1.07–3.15,  $p = 0.027$ ]) and heart failure on admission (OR: 6.00 [95% CI: 1.07–3.15,  $p = 0.027$ ]). [95% CI: 1.6–21.7;  $p = 0.007$ ].

**Conclusions:** Influenza vaccination had no effect on the outcome of patients hospitalised with COVID-19. Risk factors identified were older age, higher comorbidity and heart failure on admission.

© 2021 Elsevier España, S.L.U. All rights reserved.

## Influence of influenza vaccine and comorbidity on the evolution of hospitalised COVID-19 patients

## ABSTRACT

**Background and objective:** The COVID-19 coronavirus disease outbreak is evolving around the world. The aim of this study is to evaluate the association between influenza vaccination and the risk of mortality in hospitalized COVID-19 patients, as well as other risk factors.

**Materials and methods:** Retrospective observational study. This study was conducted among hospitalized patients with COVID-19 at Hospital La Mancha Centro between March 5 and 25, 2020. Information on influenza vaccination was extracted from electronic medical records. We used a multivariate logistic regression to explore the association between influenza vaccination and mortality from COVID and other risk factors.

**Results:** 410 patients were included. Influenza vaccine had no effect among COVID-19 hospitalized patients [OR: 1.55 (95%CI: 0.96 - 2.48;  $p=0.071$ )]. Increasing hospital mortality was associated with older age [OR: 1.05 (95% CI 1.02–1.07), per year increase;  $p<0.001$ ], Charlson  $\geq 3$  [OR: 1.84 (95%CI: 1.07–3.15,  $p=0.027$ )] and heart failure on admission [OR: 6 (95%CI: 1.6 - 21.7;  $p=0.007$ )].

## Keywords:

Influenza vaccine  
COVID-19  
Hospitalized patients  
Age  
Comorbidity  
Mortality

\* Corresponding author.

E-mail: [jmfernandez@seccam.jccm.es](mailto:jmfernandez@seccam.jccm.es) (J.M. Fernández Ibáñez).

**Conclusions:** Influenza vaccine had no effect among COVID-19 hospitalized patients. The risk factors identified were older age, higher comorbidity and heart failure on admission.

© 2021 Elsevier España, S.L.U. All rights reserved.

## Introduction

COVID-19 is a disease caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and has resulted in a pandemic that started in Wuhan, China, in December 2019. As of February 2021, the number of cases worldwide has exceeded 103 million with more than 2.2 million deaths<sup>1</sup>.

There is currently no treatment with a sufficient level of evidence. Most countries have resorted to public health measures such as isolation, quarantine and social distancing to prevent the spread of the disease.

Until vaccination for SARS-CoV-2 becomes universally available, several authors suggest that influenza vaccination may offer cross-protection against respiratory viruses other than influenza and minimise the severity of COVID-19<sup>2</sup> disease.

Other authors propose increasing anti-influenza vaccination rates to prevent co-infection of influenza with SARS-CoV-2<sup>3</sup>. However, others suggest that vaccination may increase susceptibility to SARS-CoV-2<sup>4</sup> infection.

In this article we analyse the influence of anti-influenza vaccination and other prognostic factors on mortality in COVID-19 patients admitted to Mancha Centro hospital during the first wave of the pandemic.

## Material and methods

### Design and participants

Retrospective observational cohort study. We consecutively included 410 patients admitted with a diagnosis of COVID-19 at Hospital Mancha Centro from 5 to 25 March 2020. Information on influenza vaccination in autumn 2019 was obtained from Primary Care records.

The diagnosis of COVID-19 was confirmed by polymerase chain reaction (PCR) test. Patients with negative PCR, indeterminate PCR or no PCR but with high clinical suspicion of disease were also included.

### Variables studied

The main outcome variable was mortality. Influenza vaccination was the main independent variable. Control covariates were age, sex, residence (community or institution), functionality (Barthel scale), comorbidity (Charlson index) and underlying comorbidities at admission: obesity, chronic respiratory diseases, ischaemic heart disease and/or heart failure, renal failure, chronic liver disease, haematological malignancies, other malignancies, following immunosuppressive treatment, diabetes, arterial hypertension (AHT),

**Table 1**  
Main characteristics of the patients included in the study and differences between vaccinated and unvaccinated patients

	Global (n = 410)	Unvaccinated (n = 201)	Vaccinated (n = 209)	p
Mean age in years (SD; range)	70,7 (13,9; 28 - 100)	65,6 (14,3)	75,6 (11,6)	< 0,001
<i>Age groups</i>				
Under 65 years of age	132 (32,2%)	103 (51,2%)	29 (13,9%)	< 0,001
Between 65 and 75 years of age	103 (25,1%)	45 (22,4%)	58 (27,8%)	
Between 75 and 85 years of age	113 (27,6%)	30 (14,9%)	83 (39,7%)	
Over 85 years old	62 (15,1%)	23 (11,4%)	39 (18,7%)	
<i>Sex</i>				
Man	202 (49,3%)	101 (50,2%)	101 (48,3%)	0,697
Woman	208 (50,7%)	100 (49,8%)	108 (51,7%)	
<i>Barthel</i>				
> 60	319 (77,8%)	167 (89,8%)	152 (79,2%)	0,004
≤ 60	59 (14,4%)	19 (10,2%)	40 (20,8%)	
Not available	32 (7,8%)	-	-	
<i>Charlson</i>				
< 3	232 (78,8%)	176 (87,6%)	147 (70,3%)	< 0,001
≥ 3	87 (21,2%)	25 (12,4%)	62 (29,7%)	
<i>Comorbidities</i>				
Obesity	88 (21,5%)	40 (19,9%)	48 (23%)	0,450
Chronic Respiratory Disease	145 (35,4%)	49 (24,4%)	96 (45,9%)	< 0,001
Ischaemic heart disease / heart failure	50 (12,2%)	14 (7%)	36 (17,2%)	0,002
Kidney failure	49 (12%)	20 (10%)	29 (13,9%)	0,221
Chronic liver disease	7 (1,7%)	4 (2%)	3 (1,4%)	0,719
Haematological neoplasia	9 (2,2%)	4 (2%)	5 (2,4%)	0,999
Other neoplasms	60 (7,3%)	12 (6%)	18 (8,6%)	0,304
Diabetes	109 (26,6%)	37 (18,4%)	72 (34,4%)	< 0,001
HTA	256 (62,4%)	106 (52,7%)	150 (71,8%)	< 0,001
Cognitive impairment	49 (12%)	16 (8%)	33 (15,8%)	0,015
Other neurological diseases	96 (23,4%)	41 (20,4%)	55 (26,3%)	0,157
Depressive syndrome	61 (14,9%)	27 (13,4%)	34 (16,3%)	0,420
Heart failure on admission	20 (4,9%)	6 (3%)	14 (6,7%)	0,081
Cardiovascular event during admission	6 (1,5%)	2 (1%)	4 (1,9%)	0,686
Immunosuppressive treatment	14 (3,4%)	6 (3%)	8 (3,8%)	0,639
Institutionalised	32 (7,8%)	10 (5%)	22 (10,5%)	0,036
ICU admission	24 (5,9%)	18 (9%)	6 (2,9%)	0,009
Hospital Exitus	142 (34,6%)	49 (24,4%)	93 (44,5%)	< 0,001

**Table 2**

Factors associated with in-hospital mortality according to bivariate analysis

	No exitus (n = 268)	Exitus (n =142)	P
Mean age in years (SD)	67,4 (13,8)	77 (12)	< 0,001
<i>Age groups</i>			
Under 65 years of age	113 (85,6%)	19 (14,4%)	< 0,001
Between 65 and 75 years of age	67 (65%)	36 (35%)	
Between 75 and 85 years of age	64 (56,6%)	49 (43,4%)	
Over 85 years old	24 (38,7%)	38 (61,3%)	
<i>Sex</i>			
Man	131 (64,9%)	71 (35,1%)	0,829
Woman	137 (65,9%)	71 (34,1%)	
<i>Barthel</i>			
> 60	225 (70,5%)	94 (29,5%)	< 0,001
≤ 60	27 (45,8%)	32 (54,2%)	
Not available	-	-	
<i>Charlson</i>			
< 3	230 (71,2%)	93 (28,8%)	< 0,001
= >3	38 (43,7%)	49 (56,3%)	
<i>Obesity</i>			
Yes	51 (58%)	37 (42%)	0,099
No	217 (67,4%)	105 (32,6%)	
<i>Respiratory Disease</i>			
Yes	86 (59,3%)	59 (40,7%)	0,057
No	182 (68,7%)	83 (31,3%)	
<i>Ischaemic Heart Disease / Heart Failure</i>			
Yes	22 (44%)	28 (56%)	< 0,001
No	246 (68,3%)	114 (31,7%)	
<i>Kidney failure</i>			
Yes	23 (46,9%)	26 (53,1%)	0,004
No	245 (67,9%)	116 (32,1%)	
<i>Chronic liver disease</i>			
Yes	5 (71,4%)	2 (28,6%)	0,999
No	263 (65,3%)	140 (34,7%)	
<i>Haematological neoplasia</i>			
Yes	4 (44,4%)	5 (55,6%)	0,286
No	264 (65,8%)	137 (34,2%)	
<i>Other neoplasms</i>			
Yes	11 (36,7%)	189 (63,3%)	0,001
No	257 (67,6%)	123 (32,4%)	
<i>Diabetes</i>			
Yes	68 (62,4%)	41 (37,6%)	0,445
No	200 (66,4%)	101 (33,6%)	
<i>HTA</i>			
Yes	148 (57,8%)	108 (42,2%)	< 0,001
No	120 (77,9%)	34 (22,1%)	
<i>Cognitive impairment</i>			
Yes	29 (59,2%)	20 (40,8%)	0,332
No	239 (66,2%)	122 (33,8%)	
<i>Other neurological diseases</i>			
Yes	58 (60,4%)	38 (39,6%)	0,244
No	210 (66,9%)	104 (33,1%)	
<i>Depressive syndrome</i>			
Yes	35 (57,4%)	26 (42,6%)	0,155
No	233 (66,8%)	116 (33,2%)	
<i>Heart failure during admission</i>			
Yes	3 (15%)	17 (85%)	< 0,001
No	265 (67,9%)	125 (32,1%)	
<i>Cardiovascular event during admission</i>			
Yes	2 (33,3%)	4 (66,7%)	0,190
No	262 (65,5%)	138 (34,5%)	
<i>Immunosuppressive treatment</i>			
Yes	9 (64,3%)	5 (35,7%)	0,999
No	259 (65,4%)	137 (34,6%)	
<i>Institutionalised</i>			
Yes	15 (46,9%)	17 (53,1%)	0,022
No	253 (66,9%)	125 (33,1%)	
<i>Flu vaccine</i>			
Yes	116 (55,5%)	93 (44,5%)	< 0,001
No	152 (75,6%)	49 (24,4%)	
<i>ICU admission</i>			
Yes	14 (58,3%)	10 (41,7%)	0,456
No	254 (65,8%)	132 (34,2%)	

cognitive impairment and other neurological diseases and depression. Heart failure and cardiovascular events during admission were also included.

### Statistical analysis

Quantitative variables were described by means of mean and standard deviation (SD) and qualitative variables by absolute and relative frequencies.

Bivariate analysis was used to identify the factors implicated in mortality, using Student's *t*-test for quantitative variables AND<sup>2</sup> (or Fisher's exact test) for qualitative variables.

A multivariate analysis (two-way logistic regression) was performed to independently identify possible risk factors and the role of influenza vaccination in the mortality of admitted COVID-19 patients.

All analyses were performed using the SPSS v18 statistical software and a *p*-value < 0.05 was taken as statistically significant.

### Results

A total of 410 patients admitted for COVID-19 were included. The mean age was 70.7 years (SD: 13.9; range 28 - 100). Males were 49.3% and females 50.7%; 209 (51%) were vaccinated against influenza (101 males and 108 females) and 201 (49%) were not. The scale of

Barthel was > 60 in 84.4% of cases and ≤ 60 in 15.6%. The Charlson index was < 3 in 78.8% and ≥ 3 in 21.2% of cases. The most prevalent chronic diseases were hypertension (62.4% of cases) and hypertension (62.4% of cases).

patients), chronic respiratory diseases (35.4%), neurological diseases (35.4% including cognitive impairment), diabetes (26.6%) and obesity (21.5%). The PCR test was positive in 354 patients (86.3%), negative in 35 (8.5%), indeterminate in 6 (1.5%) and not performed in 15 (3.7%). Intensive care unit (ICU) admissions were 5.9% of patients (24) and in-hospital mortality was 34.6% (142) (table 1).

Vaccinated patients are more often older than 65 years (86.2%), more dependent (Barthel < 60 20.8 vs. 10.2%; *p* = 0.004 compared to non-vaccinated), with greater comorbidity (Charlson > 3 29.7 vs. 12.4%; *p* < 0.001), with a higher percentage of chronic respiratory diseases (45.9 vs. 24.4%; *p* < 0.001), ischaemic heart disease (17.2 vs. 7%; *p* = 0.002), diabetes (34.4 vs. 18.4%; *p* < 0.001), HTN (71.8 vs. 52.7%; *p* < 0.001), cognitive impairment (15.8 vs. 8%; *p* = 0.001), cognitive impairment (15.8 vs. 8%; *p* = 0.015) and are institutionalised patients (10.5 vs. 5%; *p* = 0.036). Vaccinated patients were admitted to the ICU less often (2.9 vs. 9%; *p* = 0.009) (table 1).

The factors associated with mortality were: age being more significant with increasing age (those over 85 years died 61.3%), Barthel scale (being more at risk of death than those over 85 years), the Barthel scale (being more at risk of death than those over 85 years) and the Barthel scale (being more at risk of death than those over 85 years of age).

those patients with Barthel ≤ 60 [54.2 vs. 29.5%; *p* < 0.001]),

Charlson index ≥ 3 (56.3 vs. 28.8%; *p* < 0.001), history of ischaemic heart disease/heart failure (56 vs. 31.7%);

*p* < 0.001), having non-hematological malignancies (63.3 vs. 32.4%; *p* = 0.001), HTN (42.2 vs. 22.1%; *p* < 0.001), renal failure (53.1

vs. 32.1%; *p* = 0.004), being institutionalised (53.1 vs. 33.1%; *p* = 0.004), being in an institution (53.1 vs.

0.022) and influenza vaccination (44.5 vs. 24.4%; *p* < 0.001).

Another risk factor for mortality was the presence of heart failure during admission (85 vs. 32.1%; *p* < 0.001) (table 2).

Multivariate analysis finally identified age (OR 1.05 [95% CI 1.02-1.07 per year increase; *p* < 0.001]), Charlson index > 3 (OR

1.84 [95% CI 1.07-3.15; *p* = 0.027]) and having heart failure during admission (OR 6 [95% CI 1.6-21.7; *p* = 0.007]) as independent risk factors for in-hospital mortality. Influenza vaccination was not associated with mortality (OR: 1.55 [0.96-2.48, *p* = 0.071]) (table 3).

**Table 3**  
Multivariate analysis of factors associated with mortality

Variables	OR (95% CI)	P
Age	1,045 (1,024-1,066)	< 0,001
Charlson ( $\geq 3$ vs <3)	1,835 (1,07-3,148)	0,027
Heart failure on admission	5,993 (1,625-21,659)	0,007
Flu vaccine	1,545 (0,963-2,477)	0,071

## Discussion

In this study we show that influenza vaccination has no effect on mortality in patients admitted for COVID-19, in agreement with other authors<sup>5-6</sup>.

Vaccinated patients are older and have higher comorbidity, which are the factors associated with higher mortality in COVID-19 patients without influencing influenza vaccination.

Age, a high number of pre-admission comorbidities and the presence of heart failure on admission were the risk factors independently associated with higher in-hospital mortality in COVID-19 patients.

The vast majority of studies find age as an independent risk factor for mortality in COVID-19 patients which could be explained in relation to [immunosenescence](#)<sup>7</sup>.

Consistent with other studies, our patients with high comorbidity have a significantly higher mortality than those with low [comorbidity](#)<sup>8</sup>.

The presence of heart failure on admission was shown to be an important independent predictor of mortality in our study. According to some studies, left ventricular diastolic impairment appears to be common in acute SARS infection, even among those without underlying cardiac [disease](#)<sup>9</sup>.

Our study has some limitations. We did not include analytical data that may be associated with increased mortality in various [studies](#)<sup>8</sup>, but our aim was to assess influenza vaccination and other comorbidities in the risk of in-hospital mortality. Nor have we taken into account the treatments administered during admission, due to their heterogeneity and low level of evidence in the published [studies](#)<sup>10</sup>. In addition, the epidemiological situation may have conditioned admission criteria and bed availability and may have influenced mortality outcomes.

The strengths of our study were that the vast majority of COVID-19 cases were laboratory confirmed and all patient data were systematically collected, so we believe that the sample is representative of COVID-19 cases treated in our area.

## Conclusions

Influenza vaccination does not seem to have an effect on in-hospital mortality in COVID-19 patients admitted to our hospital. Age, high comorbidity and the presence of cardiac insufficiency on admission are independent prognostic factors for mortality, which could help physicians to identify patients with poor prognosis for management and treatment.

## Funding

This work has not received any funding.

## Conflict of interest

The authors declare that they have no conflicts of interest.

## Bibliography

1. COVID-19 Map - Johns Hopkins Coronavirus Resource Center. [Accessed 04 January 2021] Available at: <https://coronavirus.jhu.edu/map.html>.
2. Zheng J, Perlman S. Immune responses in influenza A virus and human corona- virus infections: an ongoing battle between the virus and host. *Curr Opin Virol.* 2018;28:43–52, <http://dx.doi.org/10.1016/j.coviro.2017.11.002>.
3. Zanettini C, Omar M, Dinalankara W, Luidy E, Colantuoni E, Luidy Imada E, et al. Influenza Vaccination and COVID-19 Mortality in the USA: An Ecological Study. *Vaccines (Basel).* 2021;9:427, <http://dx.doi.org/10.3390/vaccines9050427>.
4. Tetro JA. Is COVID-19 receiving ADE from other coronaviruses? *Microbes Infect.* 2020;22:72–3, <http://dx.doi.org/10.1016/j.micinf.2020.02.006>.
5. Skowronski DM, Zou M, Clarke Q, Chambers C, Dickinson JA, Sabaiduc S, et al. Influenza Vaccine Does Not Increase the Risk of Coronavirus or Other Noninfluenza Respiratory Viruses: Retrospective Analysis From Canada, 2010–2011 to 2016–2017. *Clin Infect Dis.* 2020;71:2285–8, <http://dx.doi.org/10.1093/cid/ciaa626>.
6. Del Riccio M, Lorini C, Bonaccorsi G, Paget J, Caini S. The association between influenza vaccination and the risk of SARS-COV-2 infection, severe illness, and death: A systematic review of the literature. *Int J Environ Res Public Health.* 2020;17:1–9, <http://dx.doi.org/10.3390/ijerph17217870>.
7. Sun H, Ning R, Tao Y, Yu C, Deng X, Zhao C, et al. Risk Factors for Mortality in 244 Older Adults With COVID-19 in Wuhan, China: A Retrospective Study. *J. Am. Geriatr. Soc.* 2020;68:E19–23, <http://dx.doi.org/10.1111/jgs.16533>.
8. Guan WJ, Liang WH, Zhao Y, Liang HR, Chen ZS, Li YM, et al. Comorbidity and its impact on 1590 patients with COVID-19 in China: a nationwide analysis. *Eur Respir J.* 2020;55, <http://dx.doi.org/10.1183/13993003.00547-2020>.
9. Shi S, Qin M, Shen B, Cai Y, Liu T, Yang F, et al. Association of Cardiac Injury with Mortality in Hospitalized Patients with COVID-19 in Wuhan, China. *JAMA Cardiol.* 2020;5:802–10, <http://dx.doi.org/10.1001/jamacardio.2020.0950>.
10. Song Y, Zhang M, Yin L, Wang K, Zhou Y, Zhou M, et al. COVID-19 treatment: close to a cure? A rapid review of pharmacotherapies for the novel coronavirus (SARS-CoV-2). *Int J Antimicrob Agents.* 2020;56:106080, <http://dx.doi.org/10.1016/j.ijantimicag.2020.106080>.