

Vaccines and cardiovascular outcomes: lessons learned from influenza epidemics

Siva H. Yedlapati^{1*}, Anuradha Mendu¹, Venkat R. Tummala², Sowmith S. Maganti², Khurram Nasir³, and Safi U. Khan³

¹Department of Medicine, Erie County Medical Center, 462 Grider Street, Buffalo, NY 14215, USA; ²Department of Biology, Virginia Commonwealth University, 1000 W Cary St, Richmond, VA 23284, USA; and ³Department of Cardiology, DeBakey Heart and Vascular Center, 6565 Fannin St, Houston, TX 77030, USA

KEYWORDS

Cardiovascular disease;
Influenza vaccine;
Prevention

Cardiovascular disease (CVD) is the leading cause of death in the world and is largely preventable. An increasing amount of evidence suggests that annual influenza vaccination reduces CVD-related morbidity and mortality. Despite various clinical guidelines recommending annual influenza vaccination for the general population for influenza-like illness risk reduction, with a particular emphasis on people with CVD, vaccination rates fall consistently below the goal established by the World Health Organization. This review outlines the importance of influenza vaccination, mechanisms of cardiovascular events in influenza, summarizing the available literature on the effects of influenza vaccine in CVD and the benefits of influenza vaccine during the COVID-19 pandemic.

For the audio file associated with this article, please visit the HTML version of the article online (https://academic.oup.com/eurheartjsupp/issue/25/Supplement_A)

Introduction

While the global estimates of influenza disease frequently emphasize the burden of respiratory symptoms, epidemiological data have shown the effects of influenza on cardiovascular disease (CVD) outcomes.¹ According to the World Health Organization (WHO), influenza kills up to 650 000 people annually and remains among the top 10 leading causes of mortality for all ages, particularly for those with one or more comorbid conditions like CVD.² Before the coronavirus disease 2019 (COVID-19), an estimated 29 million people in the USA contracted influenza between 2018 and 2019 flu season, hospitalizations numbered close to 380 000 and fatalities exceeded 27 000.³ All-cause influenza-attributable mortality was estimated to be 25.4 per 100 000 population and 118.2 per 100 000 for adults aged 65 years in the 2017-18 influenza season in Europe.⁴

Therefore, the WHO and the Centers for Disease Control and Prevention (CDC) Advisory Committee on Immunization

Practices make annual recommendations for the composition of the influenza vaccine to manufacturers. Various clinical guidelines recommend annual influenza vaccination for the general population for influenza-like illness risk reduction, with particular emphasis on people with CVD.⁵⁻⁷ The European Society of Cardiology guidelines (ESC-2019) and the American Heart Association and the American College of Cardiology (AHA/ACC-2006) joint guidelines suggest a Class I (Level of Evidence B) recommendation for influenza vaccination for all patients with CVD.^{5,8}

The economic burden of seasonal influenza is substantial concerning direct healthcare and indirect societal costs, stemming from the associated morbidity and mortality with estimated productivity losses.⁴ Therefore, given the critical relevance of this issue, we reviewed the sources of evidence to understand the mechanism of CVD complications associated with influenza, the effects of influenza vaccine on CVD based on randomized and observational studies, global trends and predictors associated with the vaccination uptake, and potential implications and relevance to the current COVID-19 pandemic.

*Corresponding author. Tel: +1 716 961 6995, Fax: +1 716 898 5193, Email: syedlapati@ecmc.edu

Mechanism of cardiovascular events in influenza

Studies conducted in the early 1900s following the influenza pandemic revealed a relationship between the increase in major adverse cardiovascular events (MACE), notably myocardial infarction (MI) and stroke, coinciding with or soon following an influenza epidemic.^{9,10} A century later, in 2017, influenza remains a significant cause of CVD complications. One in eight patients hospitalized with influenza had an acute cardiovascular (CV) event, with 31% of those requiring intensive care and 7% ultimately dying in the USA.¹¹ Atherosclerosis is an inflammatory response culminating in a plaque comprised of a core rich in lipids, pro-inflammatory cells and cytokines, and a fibrous cap. It is thought that influenza acts by a cascade of biological events, including the inflammatory release of cytokines that causes a pro-thrombotic state, atherosclerotic plaque destabilization and subsequent thrombosis, immune complex deposition in atherosclerotic plaques, and increased macrophage circulation leading to coronary vascular events. In addition, physiological effects such as hypoxia and tachycardia can further mediate ischaemia in subcritical stenosed vessels (Figure 1).¹²⁻¹⁴ Other mechanisms include sympathetic nervous system activation with subsequent effects on vascular tone with vasoconstriction; epithelial dysfunction; insufficient coronary artery blood flow with fever and tachycardia, volume overload, and arrhythmia.^{2,15} Finally, influenza produces direct effects on the heart to cause myocarditis. Histopathological and molecular studies on influenza-infected mice have shown that the virus can be isolated from heart tissue and that its presence leads to local inflammatory changes.¹⁶

Effects of influenza vaccine on CVD

Randomized controlled trials

Multiple randomized controlled trials (RCTs) have shown inconsistent results regarding the effects of influenza vaccination in patients with CVD (Table 1). The Flu Vaccination Acute Coronary Syndrome (FLUVACS) study was a single-blind RCT of 301 patients with MI or stable CAD scheduled for PCI.¹⁷ The FLUVACS showed a significantly lower CV death rate with the vaccine than the placebo at 1 year (6.2 vs. 17.7%, $P=0.002$).^{17,37,38} On the other hand, Influenza Vaccination in Secondary Prevention from Coronary Ischemic Events in Coronary Artery Disease (FLUCAD) study found no improvement in CV death after influenza vaccination at 12 months (3.0 vs. 5.8%, $P=0.13$). However, the rates of coronary ischaemic events over 12 months were significantly lower with the vaccine than placebo (6.02 vs. 9.97%, $P=0.04$).¹⁸ The Efficacy of Influenza Vaccine in Reducing Cardiovascular Events in Patients with Coronary Artery Disease (IVCAD) trial found no reduction in CV death or MI at 1 year with the vaccine vs. placebo (29 vs. 26%, $P=0.60$).³⁹ In contrast, the study by Phrommintikul and colleagues²⁰ was a prospective randomized open-label trial that showed a lower risk of major CV events [unadjusted hazard ratio (HR) 0.70 (95% CI: 0.57-0.86)] but no reduction in CV death.

Evidence exists that elderly patients with underlying CVD may mount a less vigorous response to standard-dose influenza vaccination.^{40,41} The Influenza Vaccine to Effectively Stop CardioThoracic Events and Decompensated Heart

Failure (INVESTED) study compared standard-dose with high-dose influenza vaccination in patients with recent MI or heart failure (HF) hospitalization in patients (mean age: 66 years) and found no reduction in all-cause mortality or cardiopulmonary hospitalizations compared with the standard dose [HR, 1.06 (95% CI: 0.97-1.17)].²¹ The Influenza Vaccine to Prevent Adverse Vascular Events (IVVE) trial evaluated influenza vaccination among patients with symptomatic HF and reported that influenza vaccination did not reduce the primary composite outcome at 36 months [HR, 0.93 (95% CI: 0.81-1.07)]. The IAMI (Influenza Vaccination After Myocardial Infarction) study, a double-blind, randomized clinical trial to test whether influenza vaccination given early after admission for MI or high-risk CAD can reduce MACE within 12 months of the index event, has shown that the primary outcome was lower in the vaccine vs. control [HR, 0.72 (95% CI: 0.52-0.99)]. The vaccinated group also experienced fewer secondary outcomes of all-cause mortality [HR, 0.59 (95% CI: 0.39-0.89)] and CV mortality [HR, 0.59 (95% CI: 0.39-0.90)].²⁴ The authors subsequently pooled their results with the FLUCAD, FLUVACS, and study by Phrommintikul *et al.* to find a 49% relative risk (RR) reduction in CV mortality [HR, 0.51 (0.36-0.71)].^{17,18,20,24}

Observational studies

Several observational studies have studied the association between the vaccine and CV outcomes (Table 1).^{31,35,42-46} A self-controlled case-series study has shown reduction in acute MI [RR, 0.68 (95% CI: 0.60-0.78)] at 1-14 days after vaccination compared with the baseline period [RR, 0.82 (95% CI: 0.75-0.90)] at 29-59 days. Early seasonal vaccinations showed more pronounced reductions in MI.⁴² Another population-based case-control study examined the association of influenza vaccination with a reduced risk of out-of-hospital primary cardiac arrest [odds ratio (OR), 0.51 (95% CI: 0.33-0.79)].⁴⁵ In the older cohort (>65 years), during the 1998-99 and 1999-2000 influenza seasons. Vaccination was associated with a reduction in the risk of hospitalization for cardiac disease [OR, 0.81 (95% CI: 0.73-0.89)], cerebrovascular disease [OR, 0.84 (95% CI: 0.72-0.97)] during the 1998-99 season as well as 1999-2000 season [OR, 0.77 (95% CI: 0.66-0.89)]. There was also a reduction in the risk of all-cause death [OR, 0.52 (95% CI: 0.47-0.57)] during the 1998-99 and [OR, 0.50 (95% CI: 0.46-0.55)] 1999-2000 seasons.⁴⁶

The United Kingdom General Practice Research Database (2001 and 2007) showed that influenza vaccination was associated with a reduced rate of acute MI [adjusted OR, 0.81 (95% CI: 0.77-0.85)]. Early seasonal influenza vaccination was associated with a lower rate of acute MI [adjusted OR, 0.79 (95% CI: 0.75-0.83)] than vaccination during the late season [adjusted OR, 0.88 (95% CI: 0.79-0.97)].⁴⁴ The Denmark data showed that for patients with HF, vaccination was associated with a reduced risk of both all-cause [adjusted HR, 0.82 (95% CI: 0.81-0.84)] and CV death [adjusted HR, 0.82 (95% CI: 0.81-0.84)].

Meta-analyses of randomized clinical trials and observational studies

Given considerable variability in the results of randomized and observational studies regarding the efficacy of the influenza vaccine for CVD prevention, meta-analyses of




Figure 1 Influenza and CV complications.

randomized and observational studies have been conducted. A Cochrane review showed a reduction in CV mortality with the influenza vaccine in patients with CVD [RR, 0.45 (95% CI: 0.26-0.76)].⁴⁷ A more recent meta-analysis of 4 randomized clinical trials and 12 observational studies in patients with CVD showed a significant reduction in all-cause [RR, 0.75 (95% CI: 0.60-0.93)] and CV mortality [RR, 0.82 (95% CI: 0.80-0.84)], and MACE [RR, 0.87 (95% CI: 0.80-0.94)] at a median follow-up of 20 months.⁴⁸ Even more up-to-date meta-analysis of 8 trials (14 420 patients) confirmed that influenza vaccine vs. control was associated with a significant 25% lower risk of MACE [RR, 0.75 (95% CI: 0.57-0.97)].⁴⁹ In relative terms, these risk reductions are comparable in size effect with guideline-directed therapy in CVD (Table 2).⁴⁹

Prevalence and predictors of influenza vaccine uptake

The WHO sets a global target of 75% vaccination uptake,⁵⁶ however, only 50% of US adults were vaccinated during influenza season between 2020 and 2021.⁵⁷ Similarly, the prevalence of vaccination was only 44% in Europe among elderly adults (≥65 years) between 2016 and 2017,⁵⁸ and 15% in the general population and 37.3% among high-risk groups in Asian counties between 2008 and 2018.⁵⁶ Figure 2 demonstrates a global review of vaccine uptake between 2009 and 2020. A US Behavioral Risk Factor Surveillance System (BRFSS, 2016-19) study estimated that the prevalence of influenza vaccine in individuals with atherosclerotic CVD was 54% for Whites, 45% for Blacks, and 42% for Hispanic adults ($P < 0.001$).⁶⁰ Various studies have discussed barriers associated with the low uptake of vaccines (Table 3). For instance, the BRFSS studies

suggested that fear of adverse reactions was the primary deterrent for the under-utilization of the influenza vaccine. Besides persistent vaccine hesitancy and misinformation, under-utilization was also considered attributable to the limited effectiveness of the existing influenza vaccine, which has ranged from 10 to 60% in estimated effectiveness in recent years.⁷⁰ Due to natural antigenic drift, predicting the viral strain makeup to include in the standard seasonal influenza vaccine is challenging and is an imprecise science.⁷¹ If the prediction is inaccurate, the effectiveness of the seasonal influenza vaccine may be low.

The effectiveness of the influenza vaccine also depends on other clinical factors, such as the age and comorbidities of the recipient. In 2018-19, the adjusted vaccine effectiveness for all influenza vaccines across all age groups was only 29%, creating more vaccine hesitancy among the general public.⁷² Due to current vaccine development strategies, it has become challenging to facilitate the rapid mass production of vaccines in response to new circulating mutations of influenza viruses.² Finally, racial and geographic disparities may also explain gaps in vaccine uptake.

Relevance during the COVID-19 pandemic

COVID-19 was declared a pandemic by the WHO on 11 March 2020, with an overall case-fatality ratio of ~2.3%.² Influenza infection and COVID-19 share a similar initial clinical presentation,⁷³ though COVID-19 appears more transmissible.⁷⁴ Unlike influenza, COVID-19 is associated with a ~10-fold higher hospitalization rate and a ~5 to 10 times higher mortality rate.^{73,75,76} One study showed the highest mortality (69.4%) among patients with elevated troponin-T and underlying CVD, followed by patients with

Table 1 Baseline characteristics of the studies

Study	Groups	n	Age	Females (%)	HTN (%)	DM (%)	HLD (%)	Obesity (%)	Smokers (%)	Follow-up (months)	Primary outcome
Randomized clinical trials											
FLUVACS ¹⁷ (2004)	Vaccine	145	64.0	29.7	59.3	19.3	—	—	45.5	12	Favourable
	Control	147	65.0	26.5	45.6	17.0	—	—	42.9		
FLUCAD ¹⁸ (2008)	Vaccine	325	58.8	28.9	69.7	19.8	—	—	20.7	12	Unfavourable
	Control	333	58.1	26.1	63.4	20.7	—	—	16.3		
IVCAD ¹⁹ (2009)	Vaccine	141	54.9	34.0	82.0	—	83.0	—	—	12	Unfavourable
	Control	137	54.5	33.0	84.0	—	90.0	—	—		
Phrommintikul <i>et al.</i> ²⁰ (2011)	Vaccine	221	65.0	39.0	63.1	29.0	44.3	—	13.7	12	Favourable
	Control	218	67.0	48.0	61.6	32.1	49.5	—	10.3		
INVESTED ²¹ (2021)	Vaccine	2630	65.5	27.3	75.7	37.2	68.4	48.4	18.4	28	Unfavourable
	Control	2630	65.5	28.8	78.4	37.1	69.4	48.8	16.0		
IVVE ^{22,23} (2022)	Vaccine	2560	57.4	52.1	64.9	22.3	16.4	—	—	36	Unfavourable
	Control	2569	57.0	50.8	64.9	23.0	16.6	—	—		
IAMI ²⁴ (2021)	Vaccine	1272	60.1	18.6	52.0	22.4	34.0	—	35.5	12	Favourable
	Control	1260	59.6	17.9	47.6	19.7	32.7	—	35.4		
Observational studies											
Diego <i>et al.</i> ²⁵ (2009)	Vaccine	860	76.7	52.0	66.7	32.1	—	24.0	6.5	40	Favourable
	Control	480	75.5	53.8	60.6	32.7	—	17.7	8.1		
Liu <i>et al.</i> ²⁶ (2012)	Vaccine	2760	74.8	41.7	80.5	54.4	42.9	—	—	48	Favourable
	Control	2288	75.7	48.2	76.2	53.2	38.8	—	—		
Wu <i>et al.</i> ²⁷ (2014)	Vaccine	2087	71.9	1.5	—	—	—	—	—	12	Favourable
	Control	429	68.3	1.6	—	—	—	—	—		
Kopel <i>et al.</i> ²⁸ (2014)	Vaccine	501	75.8	44.0	73.0	45.0	—	23.0	32.0	48	Unfavourable
	Control	1463	74.1	45.0	72.0	45.0	—	24.0	29.0		
Blaya-Nova'kova' <i>et al.</i> ²⁹ (2016)	Vaccine	1016	76.0	60.3	70.0	25.9	42.6	22.7	—	48	Favourable
	Control	1016	76.5	61.9	69.7	24.7	42.3	22.1	—		
Vardeny <i>et al.</i> ³⁰ (2016)	Vaccine	1769	67.9	19.8	68.9	40.9	—	—	—	27	Favourable
	Control	6630	62.7	22.3	71.2	32.9	—	—	—		
Modin <i>et al.</i> ³¹ (2019)	Vaccine	78	73.7	43.6	36.7	16.7	—	—	—	44.4	Favourable
	Control	379	72.8	44.8	40.3	14.9	—	—	—		
Wu <i>et al.</i> ³² (2019)	Vaccine	4350	76.3	35.1	90.1	53.7	57.6	—	—	12	Favourable
	Control	4350	76.2	34.6	89.9	53.5	56.6	—	—		
Kaya <i>et al.</i> ³³ (2017)	Vaccine	265	60.0	28.3	35.1	24.0	—	—	—	15	Favourable
	Control	391	63.0	27.6	34.3	21.0	—	—	—		
Jackson <i>et al.</i> ³⁴ (2002)	Vaccine	1016	—	34.9	49.3	24.1	—	—	23.3	27.6	Unfavourable
	Control	362	—	27.3	42.8	17.1	—	—	41.4		
Lavalley <i>et al.</i> ³⁵ (2014)	Vaccine	5054	70.0	39.7	82.7	29.5	51.4	—	19.3	24	Unfavourable
	Control	5054	69.9	40.0	82.9	29.9	51.6	—	19.5		
Mohseni <i>et al.</i> ³⁶ (2017)	Vaccine	59	74.7	49.9	—	—	—	—	—	12	Favourable
	Control	202	74.7	49.9	—	—	—	—	—		

HTN, hypertension; DM, diabetes mellitus; HLD, hyperlipidemia; n, number of patients; FLUVACS, Flu Vaccination Acute Coronary Syndrome; FLUCAD, Influenza Vaccination in Secondary Prevention from Coronary Ischemic Events in Coronary Artery Disease; IVCAD, Influenza Vaccine in Reducing Cardiovascular Events in Patients with Coronary Artery Disease; INVESTED, Influenza Vaccine to Effectively Stop CardioThoracic Events and Decompensated Heart Failure; IVVE, Influenza Vaccine to Prevent Adverse Vascular Events; IAMI, Influenza Vaccination After Myocardial Infarction.

elevated troponin-T and no underlying CVD (37.5%).⁷⁷ Another Danish study showed that the incidence of ischaemic stroke was 10 times higher, and the incidence of acute MI was 5 times higher during the 14 days after COVID-19 diagnosis compared with the control interval.⁷⁸


Benefits of influenza vaccine during COVID-19

Some studies have postulated the underlying mechanisms of COVID-19-related CVD injury overlapping with influenza infection, such as cytokine-mediated plaque destabilization, direct myocardial injury, and increased metabolic

demand.^{2,78} Studies have also shown an association between influenza vaccination status and COVID-19-related morbidity, hospitalization, and mortality.⁷⁹⁻⁸¹ A Brazilian study reported that patients with COVID-19 who received inactivated influenza vaccine experienced on average 7% lower odds of needing intensive care treatment, 17% lower need for invasive respiratory support, and 16% lower mortality than non-vaccinated patients.⁸¹ Protective effects were more prominent when the vaccine was administered after the onset of symptoms as well as among younger patients.⁸¹ An Italian study found that the higher influenza vaccination rates were associated with fewer deaths

Table 2 Barriers of influenza vaccine uptake

Study	Desire to avoid medication	Vaccine not obligatory/not needed/lack of healthcare worker advice	Fear of side effects	Belief that vaccine is ineffective and/or unsafe
Sagor (2018) ⁵⁰	✓	✓	✓	✓
Korani (2015) ⁵¹			✓	✓
Abu-Rishey (2016) ⁵²				✓
El Khoury (2015) ⁵³		✓		
Kravos (2015) ⁵⁴		✓	✓	
Bodeker (2015) ⁵⁵		✓	✓	
Sheldenkar (2019) ⁵⁶				✓

Figure 2 Percentage of population aged 65 and over vaccinated for influenza, 2009, 2019 (or nearest years) and 2020**Figure 2** Percentage of the population, aged 65 years and above, vaccinated for influenza, 2009, 2019 (or nearest years), and 2020. Note: A 3-year average for Iceland and Luxembourg for all years but 2020. Data are estimated for Norway. Source: OECD Health Statistics (2021).⁵⁹**Table 3** Efficacy of guideline-directed interventions and influenza vaccine in the prevention of major adverse cardiovascular events and/or mortality

Coronary intervention	Prevention	Intervention efficacy (%)
Smoking cessation ^{61,62}	Secondary	32-43
Statins ⁶³	Secondary	19-30
Beta-blockers ^{64,65}	Secondary	15-31
Angiotensin receptor inhibitors ⁶⁵⁻⁶⁹	Secondary	17-25
Influenza ^{48,49}	Secondary	18-41

from COVID-19; with each 1% increase in influenza vaccination among adults aged >65 years, the regional death rate of COVID-19 decreased by 0.34.⁸⁰ A meta-analysis showed that influenza vaccination was associated with a reduced risk of COVID-19 infection [RR, 0.83 (95% CI: 0.76-0.90)] and hospitalization [RR, 0.71 (95% CI: 0.59-0.84)]. Further analysis suggested that the tetravalent influenza vaccine may be associated with a reduced risk of COVID-19 infection [RR, 0.74 (95% CI: 0.65-0.84)].⁸² A Brazilian study among pregnant women showed that the influenza vaccines could confer protection against severe COVID-19 infection.⁸³

The benefit of influenza vaccination appears more critical during the COVID-19 pandemic, particularly if patients with COVID-19 are at risk of superimposed secondary infection with influenza. There are reports of respiratory viral and bacterial coinfections,⁸⁴⁻⁸⁷ and vaccination reduces the risk of influenza and may offer some incremental cardiorespiratory protection.² Some studies have suggested that influenza vaccination could be used as a temporary measure to reduce the severity of COVID-19, especially in conjunction with vaccines against SARS-CoV-2.^{79,80} Certain vaccines lead to protection against other infections through trained immunity for up to 1 year and in the case of live vaccines for up to 5 years.⁸⁸ Furthermore, influenza vaccination itself would generate sustained immunity that enhances immunity against SARS-CoV-2.⁸²

Conclusion

Influenza remains a global healthcare challenge, and a syndemic of COVID-19 and influenza would create significant issues for clinical management, diagnosis, and strain upon global health resources. While various time-series assessments have shown downstream CV effects with every future worldwide respiratory virus epidemic,^{9,10} our knowledge of predicting the next pandemic and its agent of origin remains insufficient.⁸⁹ This focused review evaluated the impact of influenza on CV morbidity and mortality and reviewed the evidence establishing the

efficacy of the influenza vaccine in patients with CVD. Furthermore, studies have highlighted co-protection mediated by the influenza vaccine among patients with COVID-19, providing a unique opportunity to control the burden of CVD complications during the current pandemic.

Funding

This paper was published as part of a supplement financially supported by Sanofi. Manuscripts were accepted after rigorous peer review process that was managed by an expert Guest Editor independently appointed by the Editor-in-Chief. The findings and conclusions contained within are those of the authors and do not necessarily reflect positions or policies of Sanofi.

Conflict of interest: None declared.

Data Availability

Data sharing not applicable to this article as no datasets were generated or analyzed during the current study.

References

- Macias AE, McElhaney JE, Chaves SS, Nealon J, Nunes MC, Samson SI, *et al.* The disease burden of influenza beyond respiratory illness. *Vaccine* 2021;**39**:A6-A14.
- Behrouzi B, Araujo Campoverde MV, Liang K, Talbot HK, Bogoch II, McGeer A, *et al.* Influenza vaccination to reduce cardiovascular morbidity and mortality in patients with COVID-19: JACC state-of-the-art review. *J Am Coll Cardiol* 2020;**76**:1777-1794.
- Centers for Disease Control and Prevention. Estimated flu illnesses, medical visits, hospitalizations, and deaths in the United States—2018-2019 flu season. 2022. <https://www.cdc.gov/flu/about/burden/2018-2019.html>. (4 August 2022).
- Nielsen J, Vestergaard LS, Richter L, Schmid D, Bustos N, Asikainen T, *et al.* European all-cause excess and influenza-attributable mortality in the 2017/18 season: should the burden of influenza B be reconsidered? *Clin Microbiol Infect* 2019;**25**:1266-1276.
- Knuuti J, Wijns W, Saraste A, Capodanno D, Barbato E, Funck-Brentano C, *et al.* 2019 ESC guidelines for the diagnosis and management of chronic coronary syndromes. *Eur Heart J* 2020;**41**:407-477.
- Maddox TM, Stecker EC, Bozkurt B, DeMichelis N, Doherty JU, Freeman A, *et al.* COVID-19 clinical guidance for the cardiovascular care team. *Am Coll Cardiol* 2020;4088. <https://digitalcommons.psjhealth.org/publications/4088>
- Smith SC, Allen J, Blair SN, Bonow RO, Brass LM, Fonarow GC, *et al.* AHA/ACC guidelines for secondary prevention for patients with coronary and other atherosclerotic vascular disease: 2006 update. *Circulation* 2006;**113**:2363-2372.
- Davis MM, Taubert K, Benin AL, Brown DW, Mensah GA, Baddour LM, *et al.* Influenza vaccination as secondary prevention for cardiovascular disease: a science advisory from the American heart association/American college of cardiology. *Circulation* 2006;**114**:1549-1553.
- Blackburn R, Zhao H, Pebody R, Hayward A, Warren-Gash C. Laboratory-Confirmed respiratory infections as predictors of hospital admission for myocardial infarction and stroke: time-series analysis of English data for 2004-2015. *Clin Infect Dis* 2018;**67**:8-17.
- Collins SD. Excess mortality from causes other than influenza and pneumonia during influenza epidemics. *Public Health Rep* 1932;**47**:2159-2179.
- Chow EJ, Rolfes MA, O'Halloran A, Anderson EJ, Bennett NM, Billing L, *et al.* Acute cardiovascular events associated with influenza in hospitalized adults: a cross-sectional study. *Ann Intern Med* 2020;**173**:605-613.
- Conti CR. Vascular events responsible for thrombotic occlusion of a blood vessel. *Clin Cardiol* 1993;**16**:761-762.
- Hebsur S, Vakili E, Oetgen WJ, Kumar PN, Lazarous DF. Influenza and coronary artery disease: exploring a clinical association with myocardial infarction and analyzing the utility of vaccination in prevention of myocardial infarction. *Rev Cardiovasc Med* 2014;**15**:168-175.
- Kinlay S, Ganz P. Role of endothelial dysfunction in coronary artery disease and implications for therapy. *Am J Cardiol* 1997;**80**:111-161.
- Corrales-Medina VF, Madjid M, Musher DM. Role of acute infection in triggering acute coronary syndromes. *Lancet Infect Dis* 2010;**10**:83-92.
- Fislova T, Gocnik M, Sladkova T, Ďurmanová V, Rajčáni J, Varečková E, *et al.* Multiorgan distribution of human influenza A virus strains observed in a mouse model. *Arch Virol* 2009;**154**:409-419.
- Gurfinkel EP, de la Fuente RL, Mendiz O, Mautner B. Flu vaccination in acute coronary syndromes and planned percutaneous coronary interventions (FLUVACS) study. *Eur Heart J* 2004;**25**:25-31.
- Ciszewski A, Bilinska ZT, Brydak LB, Kepka C, Kruk M, Romanowska M, *et al.* Influenza vaccination in secondary prevention from coronary ischaemic events in coronary artery disease: FLUCAD study. *Eur Heart J* 2008;**29**:1350-1358.
- Keshkar-Jahromi M, Vakili H, Rahnavardi M, Gholamin S, Razavi SM, Eskandari A, *et al.* The efficacy of influenza vaccination in reducing cardiovascular events in patients with coronary artery diseases: IVCAD study. *Clin Microbiol Infect* 2009;**15**:S396.
- Phrommintikul A, Kuanprasert S, Wongcharoen W, Kanjanavanit R, Chaiwarith R, Sukonthasarn A. Influenza vaccination reduces cardiovascular events in patients with acute coronary syndrome. *Eur Heart J* 2011;**32**:1730-1735.
- Vardeny O, Kim K, Udell JA, Joseph J, Desai AS, Farkouh ME, *et al.* Effect of high-dose trivalent vs standard-dose quadrivalent influenza vaccine on mortality or cardiopulmonary hospitalization in patients with high-risk cardiovascular disease: a randomized clinical trial. *JAMA* 2021;**325**:39-49.
- Loeb M, Dokainish H, Dans A, Palileo-Villanueva LM, Roy A, Karaye K, *et al.* Randomized controlled trial of influenza vaccine in patients with heart failure to reduce adverse vascular events (IVVE): rationale and design. *Am Heart J* 2019;**212**:36-44.
- Loeb M. Influenza vaccine to prevent adverse vascular events—IVVE. Presented at the American College of Cardiology Annual Scientific Session (ACC 2022), Washington, DC, April 3, 2022. 2021; 2022.
- Fröbert O, Götberg M, Erlinge D, Akhtar Z, Christiansen EH, MacIntyre CR, *et al.* Influenza vaccination after myocardial infarction: a randomized, double-blind, placebo-controlled, multicenter trial. *Circulation* 2021;**144**:1476-1484.
- de Diego C, Vila-Corcoles A, Ochoa O, Rodriguez-Blanco T, Salsench E, Hospital I, *et al.* Effects of annual influenza vaccination on winter mortality in elderly people with chronic heart disease. *Eur Heart J* 2009;**30**:209-216.
- Liu IF, Huang CC, Chan WL, Huang PH, Chung CM, Lin SJ, *et al.* Effects of annual influenza vaccination on mortality and hospitalization in elderly patients with ischemic heart disease: a nationwide population-based study. *Prev Med* 2012;**54**:431-433.
- Wu WC, Jiang L, Friedmann PD, Trivedi A. Association between process quality measures for heart failure and mortality among US veterans. *Am Heart J* 2014;**168**:713-720.
- Kopel E, Klempfner R, Goldenberg I. Influenza vaccine and survival in acute heart failure. *Eur J Heart Fail* 2014;**16**:264-270.
- Blaya-Novakova V, Prado-Galbarro FJ, Sarria-Santamera A. Effects of annual influenza vaccination on mortality in patients with heart failure. *Eur J Public Health* 2016;**26**:890-892.
- Vardeny O, Claggett B, Udell JA, Packer M, Zile M, Rouleau J, *et al.* Influenza vaccination in patients with chronic heart failure: the PARADIGM-HF trial. *JACC Heart Fail* 2016;**4**:152-158.
- Modin D, Jorgensen ME, Gislason G, Jensen JS, Kober L, Claggett B, *et al.* Influenza vaccine in heart failure. *Circulation* 2019;**139**:575-586.
- Wu HH, Chang YY, Kuo SC, Chen YT. Influenza vaccination and secondary prevention of cardiovascular disease among Taiwanese elders-A propensity score-matched follow-up study. *PloS One* 2019;**14**:e0219172.
- Kaya H, Beton O, Acar G, Temizhan A, Cavusoğlu Y, Guray U, *et al.* Influence of influenza vaccination on recurrent hospitalization in patients with heart failure. *Herz* 2017;**42**:307-315.

34. Jackson LA. Influenza vaccination is not associated with a reduction in the risk of recurrent coronary events. *Am J Epidemiol* 2002;**156**: 634-640.
35. Lavalley PC, Labreuche J, Fox KM, Lavados P, Mattle H, Steg PG, *et al*. Influenza vaccination and cardiovascular risk in patients with recent TIA and stroke. *Neurology* 2014;**82**:1905-1913.
36. Mohseni H, Kiran A, Khorshidi R, Rahimi K. Influenza vaccination and risk of hospitalization in patients with heart failure: a self-controlled case series study. *Eur Heart J* 2017;**38**:326-333.
37. Gurfinkel EP, de la Fuente RL. Two-year follow-up of the FLU vaccination acute coronary syndromes (FLUVACS) registry. *Tex Heart Inst J* 2004;**31**:28-32.
38. Gurfinkel EP, de la Fuente RL, Mendiz O, Mautner B. Influenza vaccine pilot study in acute coronary syndromes and planned percutaneous coronary interventions: the FLU vaccination acute coronary syndromes (FLUVACS) study. *Circulation* 2002;**105**:2143-2147.
39. Keshtkar-Jahromi M, Vakili H, Rahnnavardi M, Gholamin S, Razavi S, Eskandari A, *et al*. The efficacy of influenza vaccination in reducing cardiovascular events in patients with coronary artery diseases: IVCAD study: P1418. *Clin Microbiol Infect* 2009;**15**:S396.
40. Gross PA, Hermogenes AW, Sacks HS, Lau J, Levandowski RA. The efficacy of influenza vaccine in elderly persons. A meta-analysis and review of the literature. *Ann Intern Med* 1995;**123**:518-527.
41. Vardeny O, Sweitzer NK, Detry MA, Moran JM, Johnson MR, Hayney MS. Decreased immune responses to influenza vaccination in patients with heart failure. *J Card Fail* 2009;**15**:368-373.
42. Gwini SM, Coupland CA, Siriwardena AN. The effect of influenza vaccination on risk of acute myocardial infarction: self-controlled case-series study. *Vaccine* 2011;**29**:1145-1149.
43. Naghavi M, Barlas Z, Siadat S, Naguib S, Madjid M, Casscells W. Association of influenza vaccination and reduced risk of recurrent myocardial infarction. *Circulation* 2000;**102**:3039-3045.
44. Siriwardena AN, Gwini SM, Coupland CA. Influenza vaccination, pneumococcal vaccination and risk of acute myocardial infarction: matched case-control study. *Cmaj* 2010;**182**:1617-1623.
45. Siscovick DS, Raghunathan TE, Lin D, Weinmann S, Arbogast P, Lemaitre RN, *et al*. Influenza vaccination and the risk of primary cardiac arrest. *Am J Epidemiol* 2000;**152**:674-677.
46. Nichol KL, Nordin J, Mullooly J, Lask R, Fillbrandt K, Iwane M. Influenza vaccination and reduction in hospitalizations for cardiac disease and stroke among the elderly. *N Engl J Med* 2003;**348**:1322-1332.
47. Clar C, Oseni Z, Flowers N, Keshtkar-Jahromi M, Rees K. Influenza vaccines for preventing cardiovascular disease. *Cochrane Database Syst Rev* 2015;**5**.
48. Yedlapati SH, Khan SU, Talluri S, Lone AN, Khan MZ, Khan MS, *et al*. Effects of influenza vaccine on mortality and cardiovascular outcomes in patients with cardiovascular disease: a systematic review and meta-analysis. *J Am Heart Assoc* 2021;**10**:e019636.
49. Maniar YM, Al-Abdoh A, Michos ED. Influenza vaccination for cardiovascular prevention: further insights from the IAMI trial and an updated meta-analysis. *Curr Cardiol Rep* 2022;**24**:1327-1335.
50. Sagor KH, AlAtteeq MA. Beliefs, attitudes, and barriers associated with the uptake of the seasonal influenza vaccine among patients visiting primary healthcare clinics. *Saudi Med J* 2018;**39**:690-696.
51. Korani MF. Assessment of seasonal flu immunization status among adult patients visiting al-sharaee primary health care center in makkahal-mokarramah. *Int J Med Sci Public Health* 2015;**4**:117-123.
52. Abu-Rish EY, Elayeh ER, Mousa LA, Butanji YK, Albsoul-Younes AM. Knowledge, awareness and practices towards seasonal influenza and its vaccine: implications for future vaccination campaigns in Jordan. *Fam Pract* 2016;**33**:690-697.
53. Khoury GE, Salameh P. Influenza vaccination: a cross-sectional survey of knowledge, attitude and practices among the Lebanese adult population. *Int J Environ Res Public Health* 2015;**12**:15486-15497.
54. Kravos A, Kračun L, Kravos K, Iljaž R. The impact of patient's socio-demographic characteristics, comorbidities and attitudes on flu vaccination uptake in family practice settings. *Zdr Varst* 2015;**54**:204.
55. Bödeker B, Remschmidt C, Schmich P, Wichmann O. Why are older adults and individuals with underlying chronic diseases in Germany not vaccinated against flu? A population-based study. *BMC Public Health* 2015;**15**:1-10.
56. Sheldenkar A, Lim F, Yung CF, Lwin MO. Acceptance and uptake of influenza vaccines in Asia: a systematic review. *Vaccine* 2019;**37**: 4896-4905.
57. Centers for Disease Control and Prevention. Flu Vaccination Coverage, United States, 2020-21 Influenza Season 2021; 2022. [https://www.cdc.gov/flu/fluview/cv/cv-2022estimates.htm#:~:text=Overall%2C%20among%20adults%20E2%89%A518,%2D20%20season%20\(48.4%25\)\(12%20November%202022\).](https://www.cdc.gov/flu/fluview/cv/cv-2022estimates.htm#:~:text=Overall%2C%20among%20adults%20E2%89%A518,%2D20%20season%20(48.4%25)(12%20November%202022).)
58. Eurostat. 2019. <https://ec.europa.eu/eurostat/web/products-eurostat-news/-/DDN-20191209-2> (7 August 2022).
59. OECD. Percentage of population aged 65 and over vaccinated for influenza, 2009, 2019 (or nearest years) and 2020, *Health at a Glance 2021: OECD Indicators*. Paris: OECD Publishing, 2021.
60. Al Rifai M, Khalid U, Misra A, Liu J, Nasir K, Cainzos-Achirica M, *et al*. Racial and geographic disparities in influenza vaccination in the U.S. Among individuals with atherosclerotic cardiovascular disease: renewed importance in the setting of COVID-19. *Am J Prev Cardiol* 2021;**5**:100150.
61. Critchley JA, Capewell S. Smoking cessation for the secondary prevention of coronary heart disease. *Cochrane Database Syst Rev* 2012;**2**: CD003041.
62. Shah AM, Pfeffer MA, Hartley LH, Moyé LA, Gersh BJ, Rutherford JD, *et al*. Risk of all-cause mortality, recurrent myocardial infarction, and heart failure hospitalization associated with smoking status following myocardial infarction with left ventricular dysfunction. *Am J Cardiol* 2010;**106**:911-916.
63. Afllalo J, Duque G, Steele R, Jukema JW, de Craen AJ, Eisenberg MJ. Statins for secondary prevention in elderly patients: a hierarchical Bayesian meta-analysis. *J Am Coll Cardiol* 2008;**51**:37-45.
64. A randomized trial of propranolol in patients with acute myocardial infarction. I. Mortality results. *JAMA* 1982;**247**:1707-1714.
65. Freemantle N, Cleland J, Young P, Mason J, Harrison J. B blockade after myocardial infarction: systematic review and meta regression analysis. *BMJ* 1999;**318**:1730-1737.
66. Ferrari R, Bertrand ME, Remme WJ, Simoons ML, Deckers JW, Fox KM. Insight into ACE inhibition in the prevention of cardiac events in stable coronary artery disease: the EUROPA trial. *Expert Rev Cardiovasc Ther* 2007;**5**:1037-1046.
67. Flather MD, Yusuf S, Køber L, Pfeffer M, Hall A, Murray G, *et al*. Long-term ACE-inhibitor therapy in patients with heart failure or left-ventricular dysfunction: a systematic overview of data from individual patients. *Lancet* 2000;**355**:1575-1581.
68. Yusuf S, Pepine CJ, Garces C, Pouleur H, Rousseau M, Salem D, *et al*. Effect of enalapril on myocardial infarction and unstable angina in patients with low ejection fractions. *Lancet* 1992;**340**:1173-1178.
69. Yusuf S, Sleight P, Pogue J, Bosch J, Davies R, Dagenais G. Effects of an angiotensin-converting-enzyme inhibitor, ramipril, on cardiovascular events in high-risk patients. *N Engl J Med* 2000;**342**:145-153.
70. Vardeny O, Udell JA, Joseph J, Farkouh ME, Hernandez AF, McGeer AJ, *et al*. High-dose influenza vaccine to reduce clinical outcomes in high-risk cardiovascular patients: rationale and design of the INVESTED trial. *Am Heart J* 2018;**202**:97-103.
71. Harding AT, Heaton BE, Dumm RE, Heaton NS. Rationally designed influenza virus vaccines that are antigenically stable during growth in eggs. *mBio* 2017;**8**:e00669-17.
72. Flannery B, Kondor RJG, Chung JR, Gaglani M, Reis M, Zimmerman RK, *et al*. Spread of antigenically drifted influenza A(H3N2) viruses and vaccine effectiveness in the United States during the 2018-2019 season. *J Infect Dis* 2020;**221**:8-15.
73. Guan WJ, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX, *et al*. Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med* 2020;**382**: 1708-1720.
74. Liu Y, Gayle AA, Wilder-Smith A, Rocklöv J. The reproductive number of COVID-19 is higher compared to SARS coronavirus. *J Travel Med* 2020;**27**:taaa021.
75. Lai CC, Shih TP, Ko WC, Tang HJ, Hsueh PR. Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and coronavirus disease-2019 (COVID-19): the epidemic and the challenges. *Int J Antimicrob Agents* 2020;**55**:105924.
76. Ruan S. Likelihood of survival of coronavirus disease 2019. *Lancet Infect Dis* 2020;**20**:630-631.
77. Guo T, Fan Y, Chen M, Wu X, Zhang L, He T, *et al*. Cardiovascular implications of fatal outcomes of patients with coronavirus disease 2019 (COVID-19). *JAMA Cardiol* 2020;**5**:811-818.
78. Modin D, Claggett B, Sindet-Pedersen C, Lassen MCH, Skaarup KG, Jensen JUS, *et al*. Acute COVID-19 and the incidence of ischemic stroke and acute myocardial infarction. *Circulation* 2020;**142**:2080-2082.

79. Arokiaraj MC. Considering interim interventions to control COVID-19 associated morbidity and mortality—perspectives. *Front Public Health* 2020;**8**:444.
80. Marín-Hernández D, Schwartz RE, Nixon DF. Epidemiological evidence for association between higher influenza vaccine uptake in the elderly and lower COVID-19 deaths in Italy. *J Med Virol* 2021;**93**:64.
81. Fink G, Orlova-Fink N, Schindler T, Grisi S, Ferrer APS, Daubenberger C, *et al.* Inactivated trivalent influenza vaccination is associated with lower mortality among patients with COVID-19 in Brazil. *BMJ Evid Based Med* 2020;**26**:192-193.
82. Su W, Wang H, Sun C, Li N, Guo X, Song Q, *et al.* The association between previous influenza vaccination and COVID-19 infection risk and severity: a systematic review and meta-analysis. *Am J Prev Med* 2022;**63**:121-130.
83. Paganoti CF, Rodrigues AS, Francisco RPV, Costa RAD. The influenza vaccine may protect pregnant and postpartum women against severe COVID-19. *Vaccines (Basel)* 2022;**10**:206.
84. Ding Q, Lu P, Fan Y, Xia Y, Liu M. The clinical characteristics of pneumonia patients coinfecting with 2019 novel coronavirus and influenza virus in Wuhan, China. *J Med Virol* 2020;**92**:1549-1555.
85. Khodamoradi Z, Moghadami M, Lotfi M. Co-infection of coronavirus disease 2019 and influenza: a report from Iran. *Arch Iran Med* 2020;**23**: 239-243.
86. Kim D, Quinn J, Pinsky B, Shah NH, Brown I. Rates of co-infection between SARS-CoV-2 and other respiratory pathogens. *Jama* 2020;**323**: 2085-2086.
87. Wu X, Cai Y, Huang X, Yu X, Zhao L, Wang F, *et al.* Co-infection with SARS-CoV-2 and influenza A virus in patient with pneumonia, China. *Emerg Infect Dis* 2020;**26**:1324.
88. Netea MG, Domínguez-Andrés J, Barreiro LB, Chavakis T, Divangahi M, Fuchs E, *et al.* Defining trained immunity and its role in health and disease. *Nat Rev Immunol* 2020;**20**:375-388.
89. Neumann G, Kawaoka Y. Predicting the next influenza pandemics. *J Infect Dis* 2019;**219**:S14-S20.