

Image Formation

Chetan Arora

Disclaimer: The contents of these slides are taken from various publicly available resources such as research papers, talks and lectures. To be used for the purpose of classroom teaching, and academic dissemination only.



Image Formation: Luminance

- Light is emitted by light source
- Light is reflected from objects
- Reflected light is sensed by eye or by camera







Image Capture


- Place a piece of film in front of an object: Do we get a reasonable image?





Pinhole camera

- Add a barrier to block off most of the rays
 - This reduces blurring
 - The opening is known as the aperture





Effects of the Aperture Size

- In an ideal pinhole, only one ray of light reaches each point on the film
 - The image can be very dim
- Making aperture bigger makes the image blurry









Image Capturing using a Converging Lens

- A lens focuses light onto the film.
- Rays passing through the ‘Optical Center’ are not deviated.
- All rays parallel to the ‘Optical Axis’ converge at the ‘Focal Point’.





Thin Lens Equation




- Similar Triangles

$$\left. \begin{aligned} \frac{B}{A} &= \frac{e}{z} \\ \frac{B}{A} &= \frac{e-f}{f} = \frac{e}{f} - 1 \end{aligned} \right\} \frac{e}{f} - 1 = \frac{e}{z} \Rightarrow \frac{1}{f} = \frac{1}{z} + \frac{1}{e}$$



Blur Circle

- Object is out of focus \Rightarrow Blur Circle has radius: $R = \frac{L\delta}{2e}$
- A minimal L (pinhole) gives minimal R
- To capture a ‘good’ image, adjust camera settings, such that R remains smaller than the image resolution





Focus and Depth of Field (DOF)



- Although a lens can precisely focus at only one distance at a time, the decrease in sharpness is gradual on each side of the focused distance, so that within the DOF, the un-sharpness is ‘imperceptible’ under normal viewing conditions (depends upon pixel width: usually less than a pixel)
- Depth of field (DOF) is the distance between the nearest and farthest objects in a scene that appear “acceptably” sharp in an image.





DOF and Aperture


- How does the aperture affect the depth of field?
- A smaller aperture increases the range in which the object appears approximately in focus but reduces the amount of light into the camera






Focal Length and Field of View (FOV)

- As f gets smaller, image becomes more wide angle
 - more world points project onto the finite image plane
- As f gets larger, image becomes more narrow angle
 - smaller part of the world projects onto the finite image plane





Focal Length and Field of View (FOV)







Focal Length and Field of View (FOV)

FOCAL LENGTH

& ANGLE OF VIEW




LENS TYPES





Recall: Thin Lens Equation



- Similar Triangles

$$\left. \begin{aligned} \frac{B}{A} &= \frac{e}{z} \\ \frac{B}{A} &= \frac{e-f}{f} = \frac{e}{f} - 1 \end{aligned} \right\} \frac{e}{f} - 1 = \frac{e}{z} \Rightarrow \frac{1}{f} = \frac{1}{z} + \frac{1}{e}$$




Pinhole Approximation

- What happens if $z \gg f$?
- We need to adjust the image plane such that objects at infinity are in focus.

$$\frac{1}{f} = \frac{1}{z} + \frac{1}{e} \Rightarrow \frac{1}{f} \approx \frac{1}{e} \Rightarrow f \approx e$$

$$\frac{h'}{h} = \frac{f}{z} \Rightarrow h' = \frac{f}{z} h$$




- The dependence of the apparent size of an object on its depth (i.e. distance from the camera) is known as **perspective**



Perspective Effects

- Far away objects appear smaller





Perspective Effects: Foreshortening



28mm Lens



50mm Lens



70mm Lens

$$\frac{h'}{h} = \frac{f}{z}$$



Perspective Effects

- Parallel lines in the world intersect in the image at a “vanishing point”



Color




Colorimetry

- Light is **Electromagnetic Energy** in the 400nm to 700nm wavelength.
- Pure or **Monochromatic Light** is perceived as one color in the range violet-indigo-blue-green-yellow-orange-red of the rainbow.
- The amount of energy present at each wavelength is represented by the **Spectral Energy Distribution** of the light.



Colorimetry


- The spectral-energy-distributions of sunlight (what we call white light) is as shown on the right.
- Note: The dominant wavelength is not necessarily that with largest spectral component.





Color and Human Eye


- The human retina has 3 kind of color sensors (called **cones**) with approx. peak sensitivity at red, green and blue.
- The spectral response function R, G and B of these cones to monochromatic light is on the right.
- Response to blue light is less strong. The sensitivity peaks of R and G are in the yellow range.





Color Sensitivity

- Luminous efficiency function (the eye's response to light of constant luminance at a given wavelength) corresponds to the sum of the R, G, and B spectral response curves.
- Human eye seems to be most sensitive to the green hues.





Reproducing Color

- Every human eye distinguishable color can be produced by a additive mixture of red, green and blue. This is the principle of color reproduction.

Experimental RGB Mix Identification:

- The observer would alter the brightness of each of the three primary beams until a match to the test color was observed.




Reproducing Color

- Not all test colors could be matched using this technique.
- When this was the case, a variable amount of one of the primaries could be added to the test color, and a match with the remaining two primaries was carried out with the variable color spot.
- For these cases, the amount of the primary added to the test color was considered to be a negative value. In this way, the entire range of human color perception could be covered.



Reproducing Color

- Negative values of red in the range from 450nm to 525nm.
- This amount of red has to be added to the given color in order that this new color can be matched by the described values of G and B.
- Certain colors **can not** be produced by R-G-B-mixes







Color Models

- $\text{color} = xR + yG + zB = [x \quad y \quad z] \begin{bmatrix} R \\ G \\ B \end{bmatrix}$

- $\begin{bmatrix} C \\ M \\ Y \end{bmatrix} = 1 - \begin{bmatrix} R \\ G \\ B \end{bmatrix}$

- LUV, HSV, ...


- Gray





Color Sensing in Digital Cameras

- The Bayer pattern (Bayer 1976) places green filters over half of the sensors (in a checkerboard pattern), and red and blue filters over the remaining ones.
- RGB values at a pixel estimated using neighboring individual sensor values (demosaicing)





Digital Pictures

- A Matrix of numbers (Greylevel image)
- A Matrix of triplets (RGB Color, etc.)

1	2	3	4	5	6	7	8	9	10	9	8	7	6	5
2	4	5	6	7	8	9	10	11	12	11	10	9	8	7
3	5	6	7	8	9	10	11	12	13	12	11	10	9	8
4	6	7	8	9	10	11	12	13	14	13	12	11	10	9
5	7	8	9	10	11	12	13	14	15	14	13	12	11	10
6	8	9	10	11	12	13	14	15	16	15	14	13	12	11
7	9	10	11	12	13	14	15	16	17	16	15	14	13	12
8	10	11	12	13	14	15	16	17	18	17	16	15	14	13
9	11	12	13	14	15	16	17	18	19	18	17	16	15	14
10	12	13	14	15	16	17	18	19	20	19	18	17	16	15
9	11	12	13	14	15	16	17	18	19	18	17	16	15	14
8	10	11	12	13	14	15	16	17	18	17	16	15	14	13
7	9	10	11	12	13	14	15	16	17	16	15	14	13	12
6	8	9	10	11	12	13	14	15	16	15	14	13	12	11
5	7	8	9	10	11	12	13	14	15	14	13	12	11	10
4	6	7	8	9	10	11	12	13	14	13	12	11	10	9
3	5	6	7	8	9	10	11	12	13	12	11	10	9	8
2	4	5	6	7	8	9	10	11	12	11	10	9	8	7
1	3	4	5	6	7	8	9	10	11	10	9	8	7	6



Image Sampling


- Sampling the Image Plane
 - Finite number of Pixels
 - Do we always want maximum number of pixels?




Original



$\times 1/4$



$\times 1/16$





$\times 1/32$



Camera Shutter


Rolling Shutter

- Pixels are exposed roll by roll
- Good for still or slow objects
- Distorted image for very fast moving objects



Global Shutter

- All pixels are exposed simultaneously
- Good for moving objects
- No image distortion





Chetan Arora

Department of Computer Science and Engineering, IIT Delhi

Camera Shutter




Operations on Color Histogram






The Histogram

- Frequency counting of gray levels or colors








Histogram - Example







Histogram Equalization

- Equal usage of all gray levels







Histogram Equalization





Histogram Equalization



1. Given image $I(x, y)$, create a histogram H :



$$\forall x, y: H(I(x, y)) = H(I(x, y)) + 1$$

2. Create cumulative histogram $S(k)$:

$$S(0) = H(0); \quad S(k + 1) = S(k) + H(k + 1)$$



Histogram Equalization



Let m be first index for which $S(m) \neq 0$

3. Create **Look Up Table (LUT)** $T(k)$:

$$T(k) = \{ [S(k) - S(m)] / [S(255) - S(m)] \} * 255$$


4. Apply LUT T to I , get equalized image J

$$J(x, y) = T(I(x, y))$$




Adaptive Histogram Equalization

- For each pixel
 - Compute Equalization LUT in local region
 - Transform by LUT only the center pixel
- Go to next pixel





Adaptive Histogram Equalization






Other Operations with LUT


- Threshold

In	0	1	2	3	4	5	6	7
Out	0	0	0	0	7	7	7	7




- Negative

In	0	1	2	3	4	5	6	7
Out	7	6	5	4	3	2	1	0






Color Quantization





24-bit RGB color




Optimal 16 colors (4 bits)



Color Quantization



24-bit RGB



Optimal 16 colors (4 bits)



Color/Gray Level Quantization

- Quantizing the color/gray-level
 - Finite number of colors



256 Levels



8 Levels



4 Levels



2 Levels