



# Probability Theroy based on Measure Theory

STAT 517

Dr. John Smith



Copyright © 2013 John Smith

PUBLISHED BY PUBLISHER

BOOK-WEBSITE.COM

Licensed under the Creative Commons Attribution-NonCommercial 3.0 Unported License (the “License”). You may not use this file except in compliance with the License. You may obtain a copy of the License at <http://creativecommons.org/licenses/by-nc/3.0>. Unless required by applicable law or agreed to in writing, software distributed under the License is distributed on an “AS IS” BASIS, WITHOUT WARRANTIES OR CONDITIONS OF ANY KIND, either express or implied. See the License for the specific language governing permissions and limitations under the License.

*First printing, March 2013*

# Contents

I	Part One	
1	Probability Measure .....	7
1.1	Overview	7
1.2	Probability on a Field	7
1.3	Extention of Probability Measure to a $\sigma$ -field	14
1.4	Probabilities Concerning Sequences of Events	28
1.5	General Measure on a Field	37
1.6	Extension of General Measure to $\sigma$ -Field	48
1.7	Measure in Euclidean	58
II	Part Two	
2	Integration with Respect to a Measure .....	69
3	Random Variable .....	71
4	Convergence in Probability/Limit Theorem .....	73
5	Radon-Nikodym Derivative Theorem .....	75
6	Special Topics .....	77
	Index .....	79





# Part One

<b>1</b>	<b>Probability Measure .....</b>	<b>7</b>
1.1	Overview	
1.2	Probability on a Field	
1.3	Extension of Probability Measure to a $\sigma$ -field	
1.4	Probabilities Concerning Sequences of Events	
1.5	General Measure on a Field	
1.6	Extension of General Measure to $\sigma$ -Field	
1.7	Measure in Euclidean	



# 1. Probability Measure

## 1.1 Overview

## 1.2 Probability on a Field

■ **Definition 1.2.1** —  $\Omega$ . Non empty set.

■ **Definition 1.2.2** — **Paving**. A collection of a subset of  $\Omega$  is a paving.

■ **Definition 1.2.3** — **Field**. A field  $\mathcal{F}$  is a paving satisfying

- (i)  $\Omega \in \mathcal{F}$
- (ii)  $A \in \mathcal{F} \Rightarrow A^C \in \mathcal{F}$
- (iii)  $A, B \in \mathcal{F} \Rightarrow A \cup B \in \mathcal{F}$

### Derived Properties about a Field

- $\emptyset \in \mathcal{F}$  (by (i) and (ii):

$$\begin{aligned}\Omega \in \mathcal{F} &\Rightarrow \Omega^C \in \mathcal{F} \\ &\Rightarrow \emptyset \in \mathcal{F})\end{aligned}$$

- (i) can be replaced by " $\mathcal{R}$  is nonempty" because,  
Let  $A \in \mathcal{F}$ ,

$$\begin{aligned}&\Rightarrow A^C \in \mathcal{F} \\ &\Rightarrow A^C \cup A \in \mathcal{F} \\ &\Rightarrow \Omega \in \mathcal{F}\end{aligned}$$

- $A \in \mathcal{F}, B \in \mathcal{F} \Rightarrow A \cap B \in \mathcal{F}$  because,

$$\begin{aligned}(A \cap B)^C &= A^C \cup B^C \text{ (DeMorgan's Law)} \\ A \cap B &= (A^C \cup B^C)^C\end{aligned}$$

- $A_1, \dots, A_m \in \mathcal{F} \Rightarrow A_1 \cup \dots \cup A_m \in \mathcal{F}$  (mathematical induction)
- $A_1, \dots, A_m \in \mathcal{F} \Rightarrow A_1 \cap \dots \cap A_m \in \mathcal{F}$

**Definition 1.2.4 —  $\sigma$ -Field.** Similar to the definition of a field except for (iii). A paving satisfying

- (i)  $\Omega \in \mathcal{F}$
- (ii)  $A \in \mathcal{F} \Rightarrow A^c \in \mathcal{F}$
- (iii)  $A_1 \in \mathcal{F}, \dots, A_m \in \mathcal{F}$   
 $\bigcup_{k=1}^m A_k \in \mathcal{F}$  (finite additivity)

If we replace (iii) from before by (iii') here:

For  $A_1 \in \mathcal{F}, \dots, A_m \in \mathcal{F}$

$$\bigcup_{k=1}^{\infty} A_k \in \mathcal{F}$$

then  $\mathcal{F}$  is called a  **$\sigma$ -field**.

#### Derived Facts

- Again, (i) can be replaced by  $\mathcal{F}$  non empty, (iii) can be replaced  $A_1 \in \mathcal{F}, \dots, A_m \in \mathcal{F}$

■ **Example 1.1**  $\Omega = (0, 1]$  (from now on all intervals are left open, right closed)

**R** Recall that  $\sigma$ -fields are generated by fields. Fancy scripts denote a  $\sigma$ -field. Fancy scripts with a zero subscript denote a field.

$\mathcal{B}_0$  is the collection of all finite union of disjoint intervals. Asside: something that we can consider easily, but computers cannot. (e.g. think of all stars, etc)




Figure 1.1: Finite union of three disjoint intervals.

Then  $\mathcal{B}_0$  is a field.

- (i)  $(0, 1] \in \mathcal{B}_0$
- (ii)  $A \in \mathcal{B}_0 \Rightarrow A^c \in \mathcal{B}_0$
- (iii)  $A \in \mathcal{B}_0, B \in \mathcal{B}_0 \Rightarrow A \cup B \in \mathcal{B}_0$

■

Wednesday August 24







Figure 1.2: A and complement of A.

Figure 1.3: Union of A and B is still in  $\mathcal{B}_0$ 

$\mathcal{B}_0$  = collection of finite unions of disjoint subintervals of  $(0, 1]$ . Is a field.

**Definition 1.2.5 — Power Set.** A  $\sigma$ -field is generated by a paving of power set. Let  $\Omega$  be a set. The collection of all subsets of  $\Omega$  is the power set written as  $2^\Omega$ .

**R** Where does this notation come from?  
Consider the case where  $\Omega$  is finite

$$\Omega = \{\omega_1, \dots, \omega_n\}$$

Total number of subsets of  $\Omega$ .

$\emptyset$ , 1 element sets, 2-element sets, ..., n-element sets.

$$() + () + \dots + = (1 + 1)^n$$

$\#(\mathcal{F}) = 2^{\#\Omega}$ , so it seems reasonable to denote  $\mathcal{F} = 2^\Omega$ .

It is also easy to show that  $2^\Omega$  is a  $\sigma$ -field. (The largest, even. The smallest:  $\{\emptyset, \Omega\}$  which is also a  $\sigma$ -field.)

$$\{\emptyset, \Omega\} \subseteq \sigma\text{-field} \subseteq 2^\Omega$$

It turns out we can extend notion of length from  $\mathcal{B}_0$  to  $\sigma$ -field generated by  $\mathcal{B}_0$ .

Now, let  $\mathcal{A}$  be a nonempty paving of  $\Omega$ . We define

$$\sigma(\mathcal{A}) = \bigcap \{ \mathcal{B} \subseteq 2^\Omega : \mathcal{B} \text{ is a } \sigma\text{-field, } \mathcal{A} \subseteq \mathcal{B} \}$$

OR rather, the *intersection* of all  $\sigma$ -fields that contains  $\mathcal{A}$ .

Let


$$\mathbb{F}(\mathcal{A}) = \{ \mathcal{B} \subseteq 2^\Omega : \mathcal{B} \text{ is a } \sigma\text{-field, } \mathcal{B} \supseteq \mathcal{A} \}$$

Then,

$$\begin{aligned} \sigma(\mathcal{A}) &= \bigcap \mathcal{B} \\ \mathcal{B} &\in \mathbb{F}(\mathcal{A}) \end{aligned}$$

### Derived Facts

$\mathbb{F}(\mathcal{A})$  is nonempty. For example,  $2^\Omega$  is a  $\sigma$ -field and  $2^\Omega \supseteq \mathcal{A}$ .  
 $\bigcap \mathcal{B}$  is a  $\sigma$ -field. ( $\mathcal{B} \in \mathbb{F}(\mathcal{A})$ )

 Get notes about notation/levels.

*Proof.* We will prove that indeed  $\sigma(\mathcal{A})$  is a  $\sigma$ -field. Recall that we have three conditions above for  $\sigma$ -field.

(i)

$$\Omega \in \sigma(\mathcal{A})$$

$$\Omega \in \bigcap_{\mathcal{B} \in \mathbb{F}(\mathcal{A})} \mathcal{B}$$

Because:  $\mathcal{B}$  is  $\sigma$ -field,  $A \in \mathcal{B}, \forall \mathcal{B} \in \mathbb{F}(\mathcal{A})$ .

$$A \in \bigcap \mathcal{B} \Rightarrow A \in \mathcal{B} \forall \mathcal{B} \in \mathbb{F}(\mathcal{A})$$

(ii)  $\Rightarrow A^C \in \mathcal{B}, \forall \mathcal{B} \in \mathbb{F}(\mathcal{A})$

$$\Rightarrow A^C \in \bigcap_{\mathcal{B} \in \mathbb{F}(\mathcal{A})} \mathcal{B}$$

(iii)  $A_1, \dots, \in \bigcap_{\mathcal{B} \in \mathbb{F}(\mathcal{A})} \mathcal{B}, \forall \mathcal{B} \in \mathbb{F}(\mathcal{A})$

$$\Rightarrow \bigcup_{n=1}^{\infty} A_n \in \mathcal{B}, \forall \mathcal{B} \in \mathbb{F}(\mathcal{A})$$

So,  $\sigma(\mathcal{A})$  is a  $\sigma$ -field, we call it the  $\sigma$ -field, generated by  $\mathcal{B}_0$ . We know how to assign length to members of  $\mathcal{B}_0$ , we now show the assignment can be extended to  $\sigma(\mathcal{B}_0)$  ■

■ **Example 1.2** Let  $\mathcal{I}$  be the collection of *all* subintervals of  $(0,1]$ .

Note that  $\mathcal{I}$  is a smaller collection than  $\mathcal{B}_0$  since  $\mathcal{B}_0$  can have numerous different combinations of the sets.

Let

$$\mathcal{B} = \sigma(\mathcal{I})$$

This is a Borel- $\sigma$ -field. (a member of  $\mathcal{B}$  in Borel set.)

It turns out

$$\sigma(\mathcal{I}) = \sigma(\mathcal{B}_o)$$

This is because  $\sigma(\mathcal{I})$  is a  $\sigma$ -field.

So,

$$\sigma(\mathcal{I}) \supseteq \mathcal{B}_o$$

$$\sigma(\mathcal{I}) \supseteq \sigma(\mathcal{B}_o)$$

Also,

$$\mathcal{I} \subseteq \mathcal{B}_o$$

$$\sigma(\mathcal{I}) \subseteq \sigma(\mathcal{B}_o)$$

Thus,

$$\sigma(\mathcal{I}) = \sigma(\mathcal{B}_o)$$

■

**Definition 1.2.6 — Probability Measure.** Probability measures on field. Suppose  $\mathcal{F}$  is a field on a nonempty set  $\Omega$ . A probability measure is a function  $P : \mathcal{F} \rightarrow \mathbb{R}$ .

- (i)  $0 \leq P(A) \leq 1, \forall A \in \mathcal{F}$
- (ii)  $P(\emptyset) = 0, P(\Omega) = 1$
- (iii) If  $A_1, \dots$  are disjoint members of  $\mathcal{F}$  and  $\bigcup A_n \in \mathcal{F}$  then we have countable additivity:

$$P\left(\bigcup A_n\right) = \sum_{n=1}^{\infty} P(A_n)$$

**R** Note that (iii) also implies finite additivity. Prove by adding infinite empty sets on end.

If  $\Omega$  is nonempty set. And  $\mathcal{F}$  is a  $\sigma$ -field on  $\Omega$ . And  $P$  is a probability measure on  $\mathcal{F}$ . Then  $(\Omega, \mathcal{F}, P)$  is called a **probability space**. And  $(\Omega, \mathcal{F})$  is called a **measurable space**.

**R** If  $A \subseteq B$ , then  $P(A) \leq P(B)$ . This is because we may write  $B$  as

$$B = A \bigcup (B \setminus A)$$

**R**

$$P(A) + P(B) = P(A \bigcup B) + P(A \cap B)$$

**Friday August 26**

Recall,

Probability measure on a field,  $\mathcal{F}_0$ .

- $P(A) + P(B) = P(A \bigcup B) + P(A \cap B)$ 
  - $P(A) = P(AB^C) + P(AB)$
  - $P(B) = P(BA^C) + P(AB)$
  - $P(A) + P(B) = P(AB^C) + P(BA^C) + 2P(AB)$

- $P(A \cup B) = P(AB^C) + P(BA^C) + P(AB)$
- $P(A \cup B) = P(A) + P(B) - P(AB)$  By induction, we can prove if  $A_1, \dots, A_n$ ,

$$P\left(\bigcup_{k=1}^n A_k\right) = \sum_{k=1}^n P(A_k) - \sum_{i < j} P(A_i A_j) + \sum_{i < j < k} P(A_i A_j A_k) + \dots + (-1)^{n+1} P(A_1, \dots, A_n)$$

Inclusion- Exclusion Formula

- If  $A_1, \dots, A_n \in \mathcal{F}$ ,

$$B_1 = A_1$$

$$B_2 = A_2 \setminus A_1$$

$$\vdots$$

Then,

$$\bigcup_{k=1}^n A_k = \bigcup_{k=1}^n B_k$$

but the  $B_i$  are disjoint. Also  $A_k \subseteq B_k \forall k = 1, \dots, n$ .

$$P\left(\bigcup_{k=1}^n A_k\right) = P\left(\bigcup_{k=1}^n B_k\right) = \sum_{k=1}^n P(B_k) \leq \sum_{k=1}^n P(A_k)$$

Thus,  $P\left(\bigcup_{k=1}^n A_k\right) \leq \sum_{k=1}^n P(A_k)$ . Finite subadditivity.

Some conventions,

If  $A_1, \dots$  is a sequence of sets, we say  $A_n \uparrow A$  if

1.  $A_1 \subseteq A_2 \subseteq \dots$
2.  $\bigcup_{k=1}^{\infty} A_k = A$

If  $A_1, \dots$  is a sequence of sets, we say  $A_n \downarrow A$  if

1.  $A_1 \supseteq A_2 \supseteq \dots$
2.  $\bigcap_{k=1}^{\infty} A_k = A$

**Theorem 1.2.1** If  $P$  is a probability measure on a field  $\mathcal{F}$  Then,

1. Continuity from below.

If  $A_n \in \mathcal{F} \quad \forall n, A \in \mathcal{F}$

$$A_n \uparrow A$$

then

$$P(A_n) \uparrow P(A)$$

2. Continuity from above.

If  $A_n \in \mathcal{F} \quad \forall n, A \in \mathcal{F}$

$$A_n \downarrow A$$

then

$$P(A_n) \downarrow P(A)$$

3. Countable subadditivity.

If  $A_n \in \mathcal{F} \quad \forall n, \bigcup_{k=1}^{\infty} A_k \in \mathcal{F}$  then

$$P\left(\bigcup_{n=1}^{\infty} A_k\right) \leq \sum_{n=1}^{\infty} P(A_k)$$

*Proof.* 1. If  $A_1, \dots, A_n \in \mathcal{F}$ ,

$$B_1 = A_1$$

$$B_2 = A_2 \setminus A_1$$

$$B_3 = A_3 \setminus A_2$$

$$\vdots$$

then,  $B_1, \dots$  are disjoint.

$$\bigcup_{n=1}^{\infty} A_n = \bigcup_{n=1}^{\infty} B_n$$

$$\begin{aligned} P(A) &= P\left(\bigcup_{n=1}^{\infty} A_n\right) \\ &= P\left(\bigcup_{n=1}^{\infty} B_n\right) \\ &= \sum_{n=1}^{\infty} P(B_n) \\ &= \lim_{n \rightarrow \infty} \sum_{n=1}^{\infty} P(B_n) \\ &= \lim_{n \rightarrow \infty} P(A_n) \end{aligned}$$

$$2. A_n \downarrow A \Leftrightarrow A_n^C \uparrow A^C$$

But by (1),

$$P(A_n^C) \uparrow P(A^C)$$

$$1 - P(A_n) \uparrow 1 - P(A)$$

$$P(A_n) \downarrow P(A)$$

3. By finite subadditivity,

$$P\left(\bigcup_{k=1}^n A_k\right) \leq \sum_{k=1}^n P(A_k) \leq \sum_{n=1}^{\infty} P(A_n)$$

But since, by (1), because

$$\bigcup_{k=1}^n A_k \uparrow \bigcup_{n=1}^{\infty} A_n$$

$$P\left(\bigcup_{k=1}^n A_k\right) \uparrow P\left(\bigcup_{n=1}^{\infty} A_n\right)$$

So,

$$P\left(\bigcup_{n=1}^{\infty} A_n\right) \leq \sum_{n=1}^{\infty} P(A_n)$$

■

**R**  $A \in \mathcal{F} = \text{"A is F-set"}$ .

### 1.3 Extention of Probability Measure to a $\sigma$ -field

Let  $f$  be a function  $f : D \rightarrow R$ .

Let  $\tilde{D}$  be another set such that

$$D \subseteq \tilde{D}$$

An extantion of  $f$  onto  $\tilde{D}$  is

$$\tilde{f} : \tilde{D} \rightarrow R$$

Such that  $f(x) = \tilde{f}(x) \forall x \in D$

$\tilde{f}$  is an extention of  $f$  on  $D$ .

We say  $f$  has unique extention,  $\tilde{f}$  onto  $\tilde{D}$  if

1.  $\tilde{f}$  is an extension of  $f$  to  $\tilde{D}$ .
2. if  $g$  is another extension of  $f$  to  $\tilde{D}$  then  $\tilde{f} = g$  on  $D$ .

**Theorem 1.3.1** A probability measure on a field has a unique extension on the  $\sigma$ -field generated by this field.

This means that if  $\mathcal{F}_0$  is a field, and  $P$  is a probability measure on  $\mathcal{F}_0$ , then there exists a probability measure,  $Q$  on  $\sigma(\mathcal{F}_0)$  such that

$$Q(A) = P(A) \quad \forall A \in \mathcal{F}_0$$

Moreover, if  $\tilde{Q}$  is another probability measure on  $\sigma(\mathcal{F}_0)$  such that  $\tilde{Q} = P(A) \quad \forall A \in \mathcal{F}_0$  then

$$\tilde{Q} = Q$$

**R** The proof of this theorem will come after several definitions and lemmas.

**Outer Measure**  $P^* : 2^\Omega \rightarrow \mathbb{R}$

For any  $A \in 2^\Omega$  ( $A \subseteq \Omega$ )

$$P^*(A) = \inf \left\{ \sum_{n=1}^{\infty} P(A_n) : A_1, A_2, \dots \text{ is a sequence of } \mathcal{F}_0 \text{ sets, } A \subseteq \bigcup_{n=1}^{\infty} A_n \right\}$$

$P^*$  is a measure out until  $\mathcal{M}$ , but it is only a function beyond that on  $2^\Omega$ .

**Inner Measure**

$$P_*(A) = 1 - P^*(A)$$

Define the paving  $\mathcal{M}$  as follows

$$\mathcal{M} = \{A \in 2^\Omega : E \in 2^\Omega, P^*(E) = P^*(E \cap A) + P^*(E \cap A^C)\}$$

Idea: we came up with this  $\mathcal{M}$  such that  $P^*$  behaves as a measure. It will turn out to be that  $\mathcal{M}$  is a  $\sigma$ -field that contains  $\sigma(\mathcal{F}_0)$ .

### Monday August 29

$P^*$  satisfies the following probabilities:

- (i)  $P^*(\emptyset) = 0$
- (ii)  $P^*(A) \geq 0 \quad \forall A \in 2^\Omega$
- (iii)  $A \subseteq B \Rightarrow P^*(A) \leq P^*(B)$
- (iv)  $P^*(\bigcup_{n=1}^{\infty} A_n) \leq \sum_{n=1}^{\infty} P^*(A_n)$

*Proof.* (i) Take  $\{\emptyset, \emptyset, \dots\}$ .

$$\emptyset \in \mathcal{F}_0, \quad \emptyset \bigcup_{n=1}^{\infty} \emptyset$$

So,

$$P^*(\emptyset) \leq \sum_{n=1}^{\infty} P(\emptyset) = 0$$

Note,

$$P(A) \geq 0 \quad \forall A$$

So,

$$P^*(\emptyset) \geq 0$$

Thus,

$$P^*(\emptyset) = 0$$

(ii) Already done as part of (i).

(iii) Let  $A \subseteq B$

$$P^*(A) = \inf \left\{ \sum_{n=1}^{\infty} P(A_n), A_n \in \mathcal{F}_0, A \subseteq \bigcup A_n \right\}$$

Now, if  $B_1, \dots \in \mathcal{F}_0 \subseteq \bigcup B_n$

Then,

$$A \subseteq B \subseteq \bigcup_n B_n$$

If  $\{\{B_n\}_{n=1}^{\infty} : B_n \in \mathcal{F}_0, B \subseteq \bigcup_n B_n\} \subseteq \{\{A_n\}_{n=1}^{\infty} : A_n \in \mathcal{F}_0, A \subseteq \bigcup_n A_n\}$

Or in short, Collection 1  $\subseteq$  Collection 2.

So the inf of a larger set is smaller than (or equal to) the inf of a smaller set.

So,

$P^*(A) = \inf\{\sum_{n=1}^{\infty} P(A_n), A_n \in \text{collection \#1}\} \leq P^*(B) = \inf\{\sum_{n=1}^{\infty} P(B_n), A_n \in \text{collection \#2}\} = P^*(B)$   
 (iv) Want

$$P^*(\bigcup_n A_n) \leq \sum_n P^*(A_n)$$

$$P^*(A_n) = \inf\{\sum_{k=1}^{\infty} P(A_{nk}) : A_{nk} \in \mathcal{F}_0, A \subseteq \bigcup_k A_{nk}\}$$

Let  $\varepsilon > 0$ , by definition of there exists,

$$\{B_n\}_{n=1}^{\infty}$$

such that

$$\sum_{k=1}^{\infty} P(B_{nk}) \leq P^*(A_n) + \frac{\varepsilon}{2^n}$$

So,

$$\bigcup_n A_n \subseteq \bigcup_{n,k} B_{nk}$$

and,

$$\begin{aligned} P^*(\bigcup_n A_n) &\leq \sum_{n,k} P(B_{nk}) \\ &< \sum_n P^*(A_n) + \sum_n (\varepsilon 2^{-n}) \\ P^*(\bigcup_n A_n) &< \sum_n P^*(A_n) + \varepsilon \quad \forall \varepsilon > 0 \end{aligned}$$

Simply put,

$b$

So,

$$P^*(\bigcup_n A_n) \leq \sum_n P^*(A_n)$$

■

By definition,  $A \in \mathcal{M}$  if and only if  $P^*(EA) + P^*(EA^C) = P^*(E)$ .

We know that  $P^*$  is subadditive.

So, by subadditivity we know,

$$P^*(E) \leq P^*(AE) + P^*(A^C E)$$

Therefore, to show  $A \in \mathcal{M}$  we only need to show

$$P^*(E) \geq P^*(AE) + P^*(A^C E)$$

$\mathcal{M}$  is defined by  $P^*$  and  $P^*$  is defined using  $\mathcal{F}_0$  so  $\mathcal{M}$  is indirectly tied to  $\mathcal{F}_0$ .

**Lemma 1.**  $\mathcal{M}$  is a field.



*Proof.* (i)  $\Omega \in \mathcal{M}$

$$\begin{aligned} A &= \Omega \\ P^*(\emptyset) &= 0 \\ P^*(E) + P^*(\emptyset) &= P^*(E) \end{aligned}$$

(ii)  $A \in \mathcal{M} = A^C \in \mathcal{M}$

$$\begin{aligned} P^*(E) &= P^*(EA) + P^*(A^C E) \\ &= P^*(EA^C) + P^*(AE) \\ &= P^*(EA^C) + P^*((A^C)^C E) \end{aligned}$$

(iii)  $A, B \in \mathcal{M} \rightarrow A \cap B \in \mathcal{M}$

$$\begin{aligned} B \in \mathcal{M} &\Rightarrow P^*(E) = P^*(Eb) + P^*(B^C E) \quad \forall E \\ A \in \mathcal{M} &\Rightarrow P^*(BE) = P^*((BE)A) + P^*(A^C(BE)) \\ A \in \mathcal{M} &\Rightarrow P^*(B^C E) = P^*((B^C E)A) + P^*(A^C(B^C E)) \end{aligned}$$

Hence,

$$P^*(BE) + P^*(B^C E) = P^*((BE)A) + P^*(A^C(BE)) + P^*((B^C E)A) + P^*(A^C(B^C E))$$

$$\begin{aligned} P^*(A^C(BE)) + P^*((B^C E)A) + P^*(A^C(B^C E)) &\geq P^*((A^C BE) \cup (AB^C E) \cup (A^C B^C E)) \\ &= P^*(E \cap [A^C B \cup AB^C \cup A^C B^C]) \\ &= P^*(E \cap (AB)^C) \end{aligned}$$

$$\begin{aligned} P^*(E) &= P^*(BE) + P^*(B^C E) \\ &= P^*((BE)A) + P^*(A^C(BE)) + P^*((B^C E)A) + P^*(A^C(B^C E)) \\ &\geq P^*(ABE) + P^*(E(AB)^C) \end{aligned}$$

So,  $A, B \in \mathcal{M}$

■

**Lemma 2.** If  $A_1, A_2, \dots$  is a sequence of disjoint  $\mathcal{M}$ -sets then for each  $E \subseteq \Omega$ ,

$$P^*(E \cap (\bigcup_k A_k)) = \sum_k P^*(E \cap A_k)$$

*Proof.* First, prove this statement for finite sequence.

$$A_1, \dots, A_n$$

by mathematical induction.

If  $n = 1$  this is 'trivial',

$$P^*(E \cap A_1) = P^*(E \cap A_1)$$

If  $n = 2$  we need to show,

$$P^*(E(A_1 \cup A_2)) = P^*(EA_1) + P^*(EA_2)$$

Because  $A_1 \in \mathcal{M}$ ,

$$P^*(E(A_1 \cup A_2)) = P^*(E(A_1 \cup A_2)A_1) + P^*(E(A_1 \cup A_2)A_1^c)$$

$$E(A_1 \cup A_2)A_1 = E(A_1A_2 \cup A_1A_2^c) = EA_1$$

$$E(A_1 \cup A_2)A_1^c = E(A_1A_1^c \cup A_2A_1^c)$$

So,

$$P^*(E(A_1 \cup A_2)) = P^*(EA_1) + P^*(EA_2)$$

Suppose true for  $n = k$ . (induction hypothesis)

Now we must show for  $n = k + 1$ .

$$P^*(E \cap (\bigcup_{n=1}^{k+1} A_n)) = P^*([E \cap (\bigcup_{n=1}^k A_n)] \cup A_{k+1})$$

$(\bigcup_{n=1}^k A_n), A_{k+1}$  are two disjoint sets. Using the  $n=2$  case,

$$= \sum_{n=1}^k P^*(E \cap A_n) + P^*(E \cap A_{k+1}) = \sum_{n=1}^{k+1} P^*(E \cap A_n)$$

So this is now shown to be true for  $\{A_1, \dots, A_n\}$ . Next, show true for  $A_1, \dots$  in  $\mathcal{M}$  (disjoint).

Want:

$$P^*(E \cap (\bigcup_{n=1}^{\infty} A_n)) = \sum_{n=1}^{\infty} P^*(E \cap A_n)$$

Using countable subadditivity,

$$P^*(E \cap (\bigcup_{n=1}^{\infty} A_n)) = P^*(\bigcup_{n=1}^{\infty} E \cap A_n) \leq \sum_{n=1}^{\infty} P^*(E \cap A_n)$$

In the meantime, by the monotonicity of  $P^*$

$$P^*(E \cap (\bigcup_{n=1}^{\infty} A_n)) \geq P^*(E \cap (\bigcup_{n=1}^m A_n)) = \sum_{n=1}^m P^*(E \cap A_n)$$

So,

$$P^*(E \cap (\bigcup_{n=1}^{\infty} A_n)) \geq \lim_{m \rightarrow \infty} \sum_{n=1}^m P^*(E \cap A_n)$$

(\*), (\*\*) gives us,

$$P^*(E \cap (\bigcup_{n=1}^{\infty} A_n)) = \sum_{n=1}^{\infty} P^*(E \cap A_n)$$

■

**Wednesday August 31**

(finished proof)

**Lemma 3.**

1.  $\mathcal{M}$  is a  $\sigma$ -field
2.  $P^*$  restricted on  $\mathcal{M}$  is countably additive.

*Proof.* First we show if

1.  $\mathcal{M}$  is a field
2.  $\mathcal{M}$  is closed under countable disjoint union.

then  $\mathcal{M}$  is a  $\sigma$ -field.

Let's create disjoint sets,

$$A_n \in \mathcal{M}, n = 1, 2, \dots \quad B_1 = A_1 \quad B_2 = A_2 A_1^C \quad B_n = A_n A_1^C \dots A_{n-1}^C$$

$$B_1, \dots, B_n \in \mathcal{M} \text{ (disjoint)}$$

$$\bigcup_{n=1}^{\infty} B_n = \bigcup_{n=1}^{\infty} A_n$$

But we know that  $\bigcup_{n=1}^{\infty} B_n \in \mathcal{M}$  so  $\bigcup_{n=1}^{\infty} A_n \in \mathcal{M}$  and thus  $\mathcal{M}$  is a  $\sigma$ -field.

So it suffices to show that  $\mathcal{M}$  is closed under disjoint countable unions.

Let  $A_1, A_2, \dots$  are disjoint  $\mathcal{M}$ -sets.

Let  $A = \bigcup_{n=1}^{\infty} A_n$ .

Let  $F_n = \bigcup_{k=1}^n A_k$ .

Then  $F_n \in \mathcal{M}$ .

So,  $\forall E \in 2^{\Omega}$ ,

$$P^*(E) = P^*(EF_n) + P^*(EF_n^C)$$

$$\begin{aligned}
P^*(EF_n) &= P^*(E(\bigcup_{k=1}^n A_k)) \\
&= \sum_{k=1}^n P^*(EA_k) \\
P^*(EF_n^C) &\geq P^*(EA^C)(F_n \subseteq A, F_n^C \supseteq A^C) \\
\Rightarrow P^*(E) &\geq \lim_{n \rightarrow \infty} P^*(EA_k) + P^*(EA^C) \\
&= \sum_{k=1}^n P^*(EA_k) + P^*(EA^C) \\
&= P^*(EA) + P^*(EA^C)
\end{aligned}$$

■

So  $A \in \mathcal{M}$  and  $\mathcal{M}$  is a  $\sigma$ -field.

Now, let's show  $P^*$  is countably additive.

Let  $A_1, A_2, \dots$  be disjoint members of  $\mathcal{M}$ . Then  $\forall E \in 2^\Omega$ ,

$$P^*(E(\bigcup_{n=1}^{\infty} A_n)) = \sum_{n=1}^{\infty} P(EA_n)$$

Take  $E = \Omega$ .

$$P^*(\bigcup_{n=1}^{\infty} A_n) = \sum_{n=1}^{\infty} P(A_n)$$

**Lemma 4.**  $\mathcal{F}_0 \subseteq \mathcal{M}$

*Proof.* Let  $A \in \mathcal{F}$ .

Want:

$$A \in \mathcal{M}$$

$$P^*(E) = P^*(EA) + P^*(EA^C)$$

By definition, there exists  $E_n \in \mathcal{F}_0$  such that

$$\sum_{n=1}^{\infty} P^*(E_n) \leq P^*(E) + \varepsilon$$

$$\begin{aligned}
P^*(EA) &\leq P^*\left(\left(\bigcup_{n=1}^{\infty} E_n\right)A\right) \text{ (monotonocity)} \\
&= P^*\left(\bigcup_{n=1}^{\infty} E_n A\right) \\
&\leq \sum_{n=1}^{\infty} P^*(E_n A) \text{ (countably subadd)} \\
P^*(EA^C) &\leq \sum_{n=1}^{\infty} P^*(E_n A^C) \\
P^*(EA) + P^*(EA^C) &\leq \sum_{n=1}^{\infty} P^*(E_n A) + P^*(E_n A^C) \\
&= \sum_{n=1}^{\infty} P^*(E_n)
\end{aligned}$$

Recall,  $A, E_n \in \mathcal{F}_0$

$$\leq P^*(E) + \varepsilon$$

$$P^*(EA) + P^*(EA^C) \leq P^*(E) + \varepsilon \quad \forall \varepsilon$$

$$\Rightarrow P^*(EA) + P^*(EA^C) = P^*(E)$$

$$\Rightarrow A \in \mathcal{M}$$

$$\mathcal{F}_0 \in \mathcal{M}$$

■

**Lemma 5.**

$$P^*(A) = P(A) \quad \forall A \in \mathcal{F}_0$$

*Proof.* Let  $A \in \mathcal{F}_0$ .

Because,  $A, \emptyset, \emptyset, \dots, \in \mathcal{F}_0$ .

$$A \subseteq A \cup \emptyset \cup \emptyset \dots$$

$$P^*(A) \leq P(A) + P(\emptyset) + \dots$$

But, if

$$A_n \in \mathcal{F}_0$$

$$A \subseteq \bigcup_{n=1}^{\infty} A_n$$

$$P^*(A) \leq \sum_{n=1}^{\infty} P(A_n)$$

$$\Rightarrow P^*(A) \leq \inf \sum_{n=1}^{\infty} P(A_n)$$

$$= P^*(A)$$

■

**Friday September 2**



### 5 Lemma Recap

**Lemma 1.**  $\mathcal{M}$  is a field.

**Lemma 2.** If  $A_1, A_2, \dots$  is a sequence of disjoint  $\mathcal{M}$ -sets then for each  $E \subseteq \Omega$ ,

$$P^*(E \cap (\bigcup_k A_k)) = \sum_k P^*(E \cap A_k)$$

**Lemma 3.**

1.  $\mathcal{M}$  is a  $\sigma$ -field
2.  $P^*$  restricted on  $\mathcal{M}$  is countably additive.

**Lemma 4.**

$$\mathcal{F}_0 \subseteq \mathcal{M}$$

**Lemma 5.**

$$P^*(A) = P(A) \quad \forall A \in \mathcal{F}_0$$

Recall, Extension Theorem. That is, If  $\mathcal{F}$  is a field and  $P$  is a probability measure, then there exists a measure,  $Q$  such that

$$Q(A) = P(A) \quad \forall A \in \mathcal{F}_0$$

*Proof.* By Lemma 5,

$$P^*(\Omega) = P(\Omega) = 1$$

$$P^*(\emptyset) = P(\emptyset) = 0$$

Outline of what we need to show:

- $0 \leq M(A) \leq 1$
- $M(\emptyset) = 0, \quad M(\Omega) = 1$
- $M(\bigcup_n A_n) = \sum_n M(A_n)$

Since  $\forall A \in \mathcal{M}$ ,

$$\emptyset \subseteq A \subseteq \Omega$$

then

$$0 \leq P^*(\emptyset) \leq P^*(A) \leq P^*(\Omega) \leq 1$$

But, by Lemma 3,  $P^*$  is countably additive on  $\mathcal{M}$ . So  $P^*$  is probability measure on  $\mathcal{M}$  (which is a  $\sigma$ -field, by Lemma 3).

By Lemma 4,  $\mathcal{F}_0 \subseteq \mathcal{M} \Rightarrow \sigma(\mathcal{F}_0) \subseteq \mathcal{M}$ . So  $P^*$  is also probability measure on  $\sigma(\mathcal{F}_0)$ .

Finally, by Lemma 5, again  $P^*(A) = P(A)$ ,  $P^*$  is an extension of  $P$  from  $\mathcal{F}_0$  to  $\sigma(\mathcal{F}_0)$ . ■

Uniqueness of the extention,  $\pi - \lambda$  Theorem

Paving -  $\{\pi$ -system and  $\lambda$ -system.} (?)

**Definition 1.3.1 —  $\pi$ -System.** A class of subsets  $\mathcal{P}$  of  $\Omega$  is a  $\pi$  system, if

$$A, B \in \mathcal{P} \Rightarrow AB \in \mathcal{P}$$

**Definition 1.3.2 —  $\lambda$ -System.** A class  $\mathcal{L}$  is a  $\lambda$ -system if

- $\lambda$ (i)  $\Omega \in \mathcal{L}$
- $\lambda$ (ii)  $A \in \mathcal{L} \Rightarrow A^C \in \mathcal{L}$
- $\lambda$ (iii) If  $A_1, \dots \in \mathcal{L}$  are disjoint then  $\bigcup_{n=1}^{\infty} A_n \in \mathcal{L}$

So, the only difference is "disjoint". Weaker than a  $\sigma$ -field (i.e. A  $\sigma$ -field is always a  $\lambda$ -system). Note that  $(\lambda_2)$  can be replace by  $(\lambda_{2'})$  wherein

$$A, B \in \mathcal{F}, A \subseteq B, \Rightarrow B \setminus A \in \mathcal{L}$$

That is  $\lambda_1, \lambda_2, \lambda_3 \Leftrightarrow \lambda_1, \lambda_{2'}, \lambda_3$

**Lemma 6.** A class of sets that is both  $\pi$ -systema and  $\lambda$ -system is a  $\sigma$ -field.

*Proof.* Suppose  $\mathcal{F}$  is both  $\pi$ -system and  $\lambda$ -system.

By definition,

1.  $\Omega \in \mathcal{F}$
2.  $A \in \mathcal{F} \Rightarrow A^C \in \mathcal{F}$

Let  $A_1, A_2, \dots$  be  $\mathcal{F}$  sets.

Let's constructs disjoints sets, B

$$B_1 = A_1$$

$$B_2 = A_1 A_2^C$$

$$\vdots$$

Then  $B_n$  are  $\mathcal{F}$ -sets (by  $\lambda_{2'} - A_2^C = \Omega A_2^C \in \mathcal{F}$ , by  $\pi$ -system,  $A_1 A_2^C \in \mathcal{F}$  ).

By  $\lambda_3$ ,

$$\bigcup_n^{\infty} B_n \in \mathcal{F}$$

So,

$$\bigcup_n^{\infty} A_n \in \mathcal{F}$$

■

**Theorem 1.3.2 —  $\pi$ - $\lambda$  Theorem.** If  $\mathcal{P}$  is in a  $\pi$ -system,  $\mathcal{L}$  is in a  $\lambda$ -system, then

$$\mathcal{P} \subseteq \mathcal{L} \Rightarrow \sigma(\mathcal{P} \subseteq \mathcal{L})$$

*Proof.* Let  $\lambda(\mathcal{P})$  be the intersection of all  $\lambda$ -system that contains  $\mathcal{P}$ .

$$\lambda(\mathcal{P}) = \cap \{ \mathcal{L}' : \mathcal{L}' \supseteq \mathcal{P}, \mathcal{L}' \text{ is } \lambda\text{-set} \}$$

$\lambda(\mathcal{P})$  is a  $\lambda$ -system.

Goal: prove  $\lambda(\mathcal{P})$  is a  $\sigma$ -field. So we want to show that  $\lambda(\mathcal{P})$  is a  $\pi$ -system.

1.  $\Omega \in \lambda(\mathcal{P})$ ?

$$\Omega \in \mathcal{L}' \quad \forall \mathcal{L}'$$

$$\Omega \in \lambda(\mathcal{P})$$

2.  $A \in \lambda(\mathcal{P}) \Rightarrow A^C \in \lambda(\mathcal{P})$ ?

$$A \in \lambda(\mathcal{P}) \Rightarrow A \in \cap \{ \mathcal{L}' : \mathcal{L}' \supseteq \mathcal{P}, \mathcal{L}' \text{ is } \lambda\text{-set} \}$$

Then

$A \in \mathcal{L}'$  for any  $\mathcal{L}' \supseteq \mathcal{P}, \mathcal{L}'$  is  $\lambda$ -system.

$$\Rightarrow A^C \in \mathcal{L}'$$

$$\Rightarrow A^C \in \cap \{ \mathcal{L}' : \mathcal{L}' \supseteq \mathcal{P}, \mathcal{L}' \text{ is } \lambda\text{-set} \} = \lambda(\mathcal{P})$$

3.  $A_1, A_2, \dots \in \lambda(\mathcal{P})$  are disjoint then  $A_1, A_2, \dots \in \mathcal{L}' \quad \forall \mathcal{L}'$ .

Then  $\bigcup A_n \in \mathcal{L}'$  ( $\mathcal{L}'$   $\lambda$ -system)

So  $\bigcup A_n \in \lambda(\mathcal{P})$ .

We call  $\lambda(\mathcal{P})$  the  $\lambda$ -system generated by  $\mathcal{P}$ .

If we can say that  $\lambda(\mathcal{P})$  is also a  $\sigma$ -field, then  $\sigma(\mathcal{P}) \subseteq \lambda(\mathcal{P})$  because  $\sigma(\mathcal{P})$  is smallest.

So then,  $\sigma(\mathcal{P}) \subseteq \mathcal{L}$  because  $\lambda(\mathcal{P})$  is the small  $\lambda$ -system.

So it suffices to show that  $\lambda(\mathcal{P})$  is a  $\sigma$ -field. But we know if  $\lambda(\mathcal{P})$  is a system then  $\lambda(\mathcal{P})$  is  $\sigma$ -field. So it suffices to show that  $\lambda(\mathcal{P})$  is a  $\pi$ -system.

Construct again for any  $A \in 2^\Omega \quad (A \subseteq \Omega)$ , let

$$\mathcal{L}_A = \{ B : AB \in \lambda(\mathcal{P}) \}$$

Claim: If  $A \in \lambda(\mathcal{P})$  then  $\mathcal{L}_A$  is  $\lambda$ -system.

(a)  $\Omega \in \mathcal{L}_A$ ?

$$A\Omega = A \in \mathcal{L}_A$$

(b)  $(\lambda'_2) : B_1, B_2 \in \mathcal{L}_A, B_1 \subseteq B_2 \Rightarrow B_2 B_1^C \in \mathcal{L}_A$ ?

$$B_1 \in \mathcal{L}_A \Rightarrow AB_1 \in \lambda(\mathcal{P})$$

$$B_2 \in \mathcal{L}_A \Rightarrow AB_2 \in \lambda(\mathcal{P})$$

Since  $AB_1 \subseteq AB_2$ ,  $\lambda(\mathcal{P})$  is  $\lambda$ -system by  $(\lambda'_2)$  for  $\lambda(\mathcal{P})$



- (c) If  $B_n$  is disjoint,  $\mathcal{L}_A$ -sets.  
Want  $\bigcup_n B_n$  because

$$B_n \in \mathcal{L}_A$$

$$B_n A \in \lambda(\mathcal{P})$$

Because  $B_n$  disjoint we know that  $B_n A$  is also disjoint.  
Hence,

$$\bigcup_n (B_n A) \in \lambda(\mathcal{P})$$

Claim:  $\lambda(\mathcal{P})$  is  $\pi$ -system.

- (a) If  $A \in \mathcal{P}$ , then  $\mathcal{P} \subseteq \mathcal{L}_A$

Suppose  $A \in \mathcal{P}$ .

Let  $B \in \mathcal{P}$ , then  $AB \in \mathcal{P}$  ( $\pi$ -system), and  $AB \in \lambda(\mathcal{P}) \Rightarrow B \in \mathcal{L}_A$

- (b) If  $A \in \mathcal{P}$  then  $\lambda(\mathcal{P}) \subset \mathcal{L}_A$ .

- (c) If  $A \in \lambda(\mathcal{P})$ , then  $\mathcal{P} \in \mathcal{L}_A$

Suppose,  $A \in \lambda(\mathcal{P})$  and let  $B \in \mathcal{P}$ .

By step 2,

$$A \in \mathcal{L}_A$$

$$\Rightarrow AB \in \lambda(\mathcal{P})$$

$$\Rightarrow B \in \mathcal{L}_A$$

- (d) If  $A \in \lambda(\mathcal{P})$ , then  $\lambda(\mathcal{P}) \subseteq \mathcal{L}_A$ . This is because  $\lambda(\mathcal{P})$  is the smallest  $\lambda$ -system,  $\mathcal{L}_A$  is  $\lambda$ -system containing  $\mathcal{P}$  (by step 3).

Now show that  $\lambda(\mathcal{P})$  is  $\pi$ -system.

$A, B \in \lambda(\mathcal{P})$  because  $A \in \lambda(\mathcal{P})$ . We have that  $\lambda(\mathcal{P}) \in \mathcal{L}_A$ .

So

$$B \in \mathcal{L}_A$$

$$BA \in \lambda(\mathcal{P})$$

Thus  $\lambda(\mathcal{P})$  is  $\pi$ -system. ■

### Wednesday September 7

**Theorem 1.3.3** Suppose  $P_1$  and  $P_2$  are probability measures on  $\sigma(\mathcal{P})$  where  $\mathcal{P}$  is a  $\pi$ -system. If  $P_1$  and  $P_2$  agree on  $\mathcal{P}$  (that is,  $P_1(A) = P_2(A) \quad \forall A \in \mathcal{P}$ ) then they agree on  $\sigma(\mathcal{P})$ .

*Proof.* Let

$$\mathcal{L} = \{A \in \sigma(\mathcal{P}) : P_1(A) = P_2(A)\}$$

Then  $\mathcal{P} \subseteq \mathcal{L}$ .

It suffices to show that  $\mathcal{L}$  is a  $\lambda$ -system (because if so, then  $\sigma(\mathcal{P}) \subseteq \mathcal{L}$  - in fact,  $\sigma(\mathcal{P}) = \mathcal{L}$ ).

Show  $\mathcal{L}$  is a  $\lambda$ -system.

1.  $\Omega \in \mathcal{L}$ ?

$$P_1(\Omega) = P_2(\Omega) = 1, \quad \Omega \in \mathcal{P}$$

2.  $A \in \mathcal{L}$

$$P_1(A) = P_2(A) \Rightarrow P_1(A^C) = P_2(A^C), \quad A^C \in \mathcal{L}$$

3.  $A \in \mathcal{L}$ .  $A_n$  disjoint. Want  $\bigcup_{n=1}^{\infty} A_n \in \mathcal{L}$ .  
Since

$$A_n \in \mathcal{L}$$

$$P_1(A_n) = P_2(A_n) \quad \forall n$$

$$\sum_{n=1}^{\infty} P_1(A_n) = \sum_{n=1}^{\infty} P_2(A_n)$$

$$P_1\left(\bigcup_{n=1}^{\infty} A_n\right) = P_2\left(\bigcup_{n=1}^{\infty} A_n\right)$$

So,  $\bigcup A_n \in \mathcal{L}$ . ■

So our extension of (and uniqueness of the extension of)  $P$  on  $\mathcal{F}_0$  to  $\sigma(\mathcal{F}_0)$  is complete. We have shown the existence of  $Q$  on  $\mathcal{M}$ .

Since  $Q$  agrees with  $P$  on  $\mathcal{F}_0$  and  $\mathcal{F}_0$  is a field, this implies that this is a  $\pi$ -system.

If you have another extension, say  $\tilde{Q}$ , then  $\tilde{Q} = P$  on  $\mathcal{F}_0$ . That is,  $\tilde{Q} = Q$  on  $\mathcal{M}$ , where  $\mathcal{M}$  is a  $\sigma$ -field, which is a  $\pi$ -system.

So by Theorem 1.3.3,

$$\tilde{Q} = Q \text{ on } \sigma(\mathcal{P}).$$

So, Theorem 1.3.1, is completely proved.

Lemma 1 - Lemma 5 imply the extension.

$\pi - \lambda$  Theorem and Theorem 1.3.3 implies uniqueness.

This wraps up Theorem 1.3.1.

### Lebesgue measure on $(0,1]$

$$\Omega = (0,1]$$

Recall,  $\mathcal{B}_0$  is the finite disjoint unions of intervals in  $(0,1]$  and that  $\mathcal{B}_0$  is a field.

Let  $\mathcal{B} = \sigma(\mathcal{B}_0)$ .

For each  $A \in \mathcal{B}_0$ ,

$$A = \bigcup_{i=1}^n (a_i, b_i]$$

Let  $\lambda(A) = \sum_{i=1}^n (b_i - a_i)$ .

Question: Is  $\lambda$  a probability measure on  $\mathcal{B}_0$ ?

**Theorem 1.3.4 — Theorem 2.2 in Billingsly.** The set function  $\lambda$  on  $\mathcal{B}_0$  is a probability measure on  $\mathcal{B}_0$ .

*Proof.* 1.  $0 \leq \lambda(A) \leq 1$

2.

$$\lambda(\Omega) = \lambda((0, 1]) = 1 - 0 = 1$$

$$\lambda(\emptyset) = \lambda((0, 0]) = 0$$

3. This one requires Theorem 1.3 in Billingsly (proof omitted - calculus, blah).

Theorem 1.3 - If  $I$  is an interval in  $(0, 1]$  and  $\{I_k : k = 1, 2, \dots\}$  are disjoint intervals in  $(0, 1]$  such that

$$I = \bigcup_{k=1}^{\infty} I_k$$

then,

$$|I| = \sum_{k=1}^{\infty} |I_k|$$

where  $|a|$  means length of interval  $a$ .

Since  $\bigcup_{j=1}^{m_k} I_{kj} \in \mathcal{B}_0$  and  $\bigcup_{i=1}^m I_i = \bigcup_{k=1}^{\infty} \bigcup_{j=1}^{m_k} I_{kj}$ .

Then

$$\lambda(A) \lambda\left(\bigcup_{i=1}^m I_i\right) = \sum_{i=1}^m |I_i|$$

Since,  $I_i \subset \bigcup_{k=1}^{\infty} \bigcup_{j=1}^{m_k} I_{kj}$ , then

$$I_i = I_i \left( \bigcup_{k=1}^{\infty} \bigcup_{j=1}^{m_k} I_{kj} \right) = \bigcup_{k=1}^{\infty} \bigcup_{j=1}^{m_k} I_i I_{kj}$$

By Theorem 1.3,

$$|I| = \sum_{k=1}^{\infty} \sum_{j=1}^{m_k} |I_i I_{kj}|$$

$$\lambda(A) = \sum_{i=1}^m \sum_{k=1}^{\infty} \sum_{j=1}^{m_k} |I_i I_{kj}| = \sum_{k=1}^{\infty} \sum_{j=1}^{m_k} \sum_{i=1}^m |I_i I_{kj}|$$

Because  $I_{jk} \subseteq \bigcup_{i=1}^m I_i$ , we have that

$$I_{kj} = \bigcup_{i=1}^m I_{kj} I_i$$

Again by Theorem 1.3, (note that  $I_{kj} I_i$  are disjoint intervals)

$$|I_{kj}| = \sum_{i=1}^m |I_{kj} I_i|$$

$$\text{So, } \lambda(A) = \sum_{k=1}^{\infty} \sum_{j=1}^{m_k} |I_{kj}| = \sum_{k=1}^{\infty} \lambda(A_k)$$

■

### Friday September 9

Finished above proof.

So  $\lambda$  is a probability on  $\mathcal{B}_0$ . By Theorem 3.1, there exists a unique measure  $\tau$  on  $\sigma(\mathcal{B}_0) = \mathcal{B}$  such that

$$\tau(A) = \lambda(A) \quad \forall A \in \mathcal{B}_0$$

$\tau$  is called **Lebesgue Measure** on  $(0,1]$ . We may still write it as  $\lambda$ .

## 1.4 Probabilities Concerning Sequences of Events

### Set Limit

Let  $(\Omega, \mathcal{F})$  be a measurable space (i.e.  $\Omega$  is nonempty set and  $\mathcal{F}$  is  $\sigma$ -field).

let  $A_1, \dots \in \mathcal{F}$ . We define

$$\bigcap_{n=1}^{\infty} \bigcup_{k=n}^{\infty} A_k = \limsup_{n \rightarrow \infty} A_n$$

It is trivial to show that  $\limsup_{n \rightarrow \infty} A_n \in \mathcal{F}$ .

$$\bigcup_{n=1}^{\infty} \bigcap_{k=n}^{\infty} A_k = \liminf_{n \rightarrow \infty} A_n$$

We swapped intersection/union...what we are doing here?

$\omega$  (means outcome)  $\in \Omega$

$$\omega \in \limsup_{n \rightarrow \infty} A_n \Leftrightarrow \omega \in \bigcap_{n=1}^{\infty} \bigcup_{k=n}^{\infty} A_k$$

$$\Leftrightarrow \omega \in \bigcup_{k=1}^{\infty} A_k \quad \forall n = 1, 2, \dots$$

$$\Leftrightarrow \omega \in A_k \quad \text{for some } k \geq n, \quad \forall n = 1, 2, \dots$$

$\Leftrightarrow \omega$  is in infinitely many  $k$ .

Similarly,

$$\omega \in \liminf_{n \rightarrow \infty} A_n \Leftrightarrow \omega \in \bigcup_{n=1}^{\infty} \bigcap_{k=n}^{\infty} A_k$$

$$\Leftrightarrow \omega \in \bigcap_{k=1}^{\infty} A_k \quad \text{for some } n = 1, 2, \dots$$

$$\Leftrightarrow \omega \in A_k \quad \forall k \geq n, \quad \text{for some } n$$

$$\Leftrightarrow \omega \in \text{all but finitely many } A_k$$

So this is a much stronger requirement. Intuitively, if  $\omega$  is in all but finitely many  $A_k$ , then it must be in infinitely many  $A_k$  (i.e.  $\liminf_{n \rightarrow \infty} A_n \subseteq \limsup_{n \rightarrow \infty} A_n$ ).

For  $i > \max(n, m)$ ,

$$\begin{aligned} \cap_{k=m}^{\infty} A_k &\subseteq A_i \subseteq \bigcup_{k=n}^{\infty} A_k \\ \Rightarrow \bigcup_{m=1}^{\infty} \cap_{k=m}^{\infty} A_k &\subseteq \bigcup_{k=n}^{\infty} A_k \\ \Rightarrow \bigcup_{m=1}^{\infty} \cap_{k=m}^{\infty} A_k &\subseteq \cap_{n=1}^{\infty} \bigcup_{k=n}^{\infty} A_k \\ &\Rightarrow \liminf_{n \rightarrow \infty} A_n \subseteq \limsup_{n \rightarrow \infty} A_n \end{aligned}$$

$$\begin{aligned} \cap_{k=n}^{\infty} A_k &\uparrow \liminf_{n \rightarrow \infty} A_n \\ \bigcup_{k=n}^{\infty} A_k &\downarrow \limsup_{n \rightarrow \infty} A_n \end{aligned}$$

If  $\liminf_{n \rightarrow \infty} A_n = \limsup_{n \rightarrow \infty} A_n$ , then we say that the sequences  $\{A_n\}$  has a limit,

$$\begin{aligned} \lim_{n \rightarrow \infty} A_n &= \liminf_{n \rightarrow \infty} A_n = \limsup_{n \rightarrow \infty} A_n \\ \lim_{n \rightarrow \infty} A_n &\in \mathcal{F} \end{aligned}$$

Sometimes we write,

$$\limsup_{n \rightarrow \infty} A_n = [A_n \text{ i.o. }]$$

**Theorem 1.4.1** Suppose  $(\Omega, \mathcal{F}, P)$  is a probability space and  $A_n \in \mathcal{F} \quad n = 1, 2, \dots$

(i)

$$\limsup_{n \rightarrow \infty} P(A_n) \leq P(\limsup_{n \rightarrow \infty} A_n)$$

$$\liminf_{n \rightarrow \infty} P(A_n) \geq P(\liminf_{n \rightarrow \infty} A_n)$$

(ii)  $A_n \rightarrow A$  ( $A = \lim_{n \rightarrow \infty} A_n$ ), then we have continuity of probability of a set function:

$$\lim_{n \rightarrow \infty} P(A_n) = P(\lim_{n \rightarrow \infty} A_n)$$

**Monday September 12**

*Proof.* (i) Let  $B_n = \cap_{k=n}^{\infty} A_k$ .

$$B_n \uparrow \liminf_n A_n$$

By Theorem 2.1,

$$P(B_n) \uparrow P(\liminf_n A_n)$$

So,

$$P(B_n) \leq P(\liminf_n A_n) \quad \forall n$$

$$\lim_{n \rightarrow \infty} P(B_n) = P(\liminf_n A_n)$$

$$P(A_n) \geq P(B_n) \rightarrow P(\liminf_n A_n)$$

$$\liminf_n P(A_n) \geq P(\liminf_n A_n)$$

Similarly,

Let  $C_n = \bigcup_{k=n}^{\infty} A_k$ .

Then,

$$C_n \downarrow \bigcup_{k=n}^{\infty} A_k$$

$$P(A_n) \leq P(C_n) \rightarrow P(\limsup_n A_n)$$

$$\limsup_n P(A_n) \leq P(\limsup_n A_n)$$

(ii) If  $A_n$  has a limit (i.e.  $\limsup_n A_n = \liminf_n A_n = \lim A$ ) then,

$$\liminf_n P(A_n) \geq P(\liminf_n A_n) = P(\limsup_n A_n) \geq \limsup_n P(A_n)$$

So,  $\liminf_n P(A_n) = \limsup_n P(A_n)$ , thus

$$\lim_n P(A_n) = P(\lim_n A_n)$$

■

### Independent Events

$(\Omega, \mathcal{F}, P)$

Let  $A, B \in \mathcal{F}$ . They are independent if and only iff:

$$P(AB) = P(A)P(B)$$

$$A \perp\!\!\!\perp B$$

$A_1, \dots, A_n$  are independent if and only if for any  $\{k_1, \dots, k_j\} \subseteq \{1, \dots, n\}$ ,

$$P(A_{k_1} \dots A_{k_j}) = P(A_{k_1}) \dots P(A_{k_j})$$

In this case we write:  $A_1 \perp\!\!\!\perp \dots \perp\!\!\!\perp A_n$ .

Now let,  $\mathcal{A}_1, \dots, \mathcal{A}_n$  be pavings in  $\mathcal{F}$  (i.e.  $\mathcal{A}_k \subseteq \mathcal{F}$ ).

We say  $\mathcal{A}_1, \dots, \mathcal{A}_n$  are independent if and only if for any  $A_1 \in \mathcal{A}_1, \dots, A_n \in \mathcal{A}_n$  we have

$$A_1 \perp\!\!\!\perp \dots \perp\!\!\!\perp A_n$$

In this case we write:  $\mathcal{A}_1 \perp\!\!\!\perp \dots \perp\!\!\!\perp \mathcal{A}_n$ .

**Theorem 1.4.2** Suppose for  $(\Omega, \mathcal{F}, P)$  is a probability space if,

$$\mathcal{A}_1 \subseteq \mathcal{F} \dots \mathcal{A}_n \subseteq \mathcal{F}$$

are  $\pi$ -systems. Then,

$$\mathcal{A}_1 \perp\!\!\!\perp \dots \perp\!\!\!\perp \mathcal{A}_n \Rightarrow \sigma(\mathcal{A}_1) \perp\!\!\!\perp \dots \perp\!\!\!\perp \sigma(\mathcal{A}_n)$$

*Proof.* Let  $\mathcal{B}_i = \mathcal{A}_i \cup \{\Omega\}$ .

It is easy to show (in homework)

1.  $\mathcal{B}_i$  is still a  $\pi$ -system
2.  $\mathcal{B}_i$  are still independent

$$\mathcal{B}_1 \perp\!\!\!\perp \dots \perp\!\!\!\perp \mathcal{B}_n$$

For  $B_2 \in \mathcal{B}_2, \dots, B_n \in \mathcal{B}_n$  define,

$$\mathcal{L}(B_2, \dots, B_n) = \{B \in \mathcal{F} : B \perp\!\!\!\perp B_2 \perp\!\!\!\perp \dots \perp\!\!\!\perp B_n\}$$

1. First we show  $\mathcal{L}(B_2, \dots, B_n)$  is  $\lambda$ -system.

$$\Omega \in \mathcal{L}(B_2, \dots, B_n)$$

$$\Omega \perp\!\!\!\perp B_1 \perp\!\!\!\perp \dots \perp\!\!\!\perp B_n$$

This is true because  $P(\Omega B_2 \dots B_n) = P(B_2 \dots B_n) = P(B_2) \dots P(B_n) = P(\Omega)P(B_2) \dots P(B_n)$

2. Now  $A \in \mathcal{L}(B_2, \dots, B_n) \Rightarrow A^C \in \mathcal{L}(B_2, \dots, B_n)$

$$A \in \mathcal{L}(B_2, \dots, B_n)$$

$$\Rightarrow A \perp\!\!\!\perp B_2 \perp\!\!\!\perp \dots \perp\!\!\!\perp B_n$$

$$\Rightarrow P(AB_2 \dots B_n) = P(A)P(B_2) \dots P(B_n)$$

$$\Rightarrow P(A^C B_2 \dots B_n)$$

$$P(B_2 \dots B_n) \setminus P(AB_2 \dots B_n)$$

$$P(B_2 \dots B_n) - P(AB_2 \dots B_n)$$

$$P(B_2) \dots P(B_n) - P(A)P(B_2) \dots P(B_n)$$

$$(1 - P(A))P(B_2) \dots P(B_n) = P(A^C)P(B_2) \dots P(B_n)$$

Then we run this through all subadditives of  $A, B_2, \dots, B_n$ .

$$A^C \perp\!\!\!\perp B_2 \perp\!\!\!\perp \dots \perp\!\!\!\perp B_n$$

3. If  $C_1, C_2, \dots \in \mathcal{L}(B_2, \dots, B_n)$  they are disjoint. Want to show

$$\bigcup_{m=1}^i C_m \in \mathcal{L}(B_2, \dots, B_n)$$

$$C_m \in \mathcal{L}(B_2, \dots, B_n)$$

$$\Rightarrow C_m \perp\!\!\!\perp \dots \perp\!\!\!\perp B_n$$

$$\Rightarrow P(C_m B_2 \dots B_n) = P(C_m) \dots P(B_n) \quad \forall m = 1, 2, \dots$$

$$\sum_{m=1}^{\infty} P(C_m B_2 \dots B_n) = \left( \sum_{m=1}^{\infty} P(C_m) \right) P(B_2) \dots P(B_n)$$

But  $\{C_m, B_2, \dots, B_n, m = 1, 2, \dots\}$

So  $\bigcup_m C_m \in \mathcal{L}(B_2, \dots, B_n)$ .

And  $\mathcal{L}(B_2, \dots, B_n)$ .

Also,  $B_1 \in \mathcal{L}(B_2, \dots, B_n) \quad \forall B_1 \in \mathcal{B}_1$  therefore by definition,

$$\mathcal{B}_1 \subseteq \mathcal{L}(B_2, \dots, B_n)$$

So,  $\sigma(\mathcal{B}_1) \subseteq \mathcal{L}(B_2, \dots, B_n)$  and we have our  $\lambda - \pi$ -theorem.

This means that for all  $B_1 \in \sigma(\mathcal{B}_1)$

$$B_1 \perp\!\!\!\perp B_2 \perp\!\!\!\perp \dots \perp\!\!\!\perp B_n$$

Recall that  $B_i$  are arbitrary members of,

$$\sigma(\mathcal{B}_1) \perp\!\!\!\perp B_2 \perp\!\!\!\perp \dots \perp\!\!\!\perp B_n \Leftrightarrow \mathcal{B}_2 \perp\!\!\!\perp \sigma(\mathcal{B}_1) \perp\!\!\!\perp \dots \perp\!\!\!\perp \mathcal{B}_n$$

Run the previous argument repeatedly.

So

$$\sigma(\mathcal{B}_1) \perp\!\!\!\perp \sigma(\mathcal{B}_2) \perp\!\!\!\perp \dots \perp\!\!\!\perp \sigma(\mathcal{B}_n)$$

■

■ **Example 1.3** Let  $\mathcal{I}$  be the collection of all intervals, then its  $\pi$ -system. When we want to check  $X \perp\!\!\!\perp Y$ , we only need to check

$$P(X \in \text{interval}, Y \in \text{interval}) = P(X \in \text{interval})P(Y \in \text{interval})$$

■

**Wednesday September 14**

### Independence of Infinite Classes

Let  $\{\mathcal{A}_\theta : \theta \in \Theta\}$  where  $\theta$  is any infinite set (need not be countable) if and only if any (infinite)  $\{A_\theta : \theta \in \Theta\}$  where  $A_\theta \in \mathcal{A}_\theta$  are independent.

We already define independence of  $\{A_\theta : \theta \in \Theta\}$ ; that is for an infinite collection of sets is independent if and only if any finite subcollection  $\{A_{\theta_1}, \dots, A_{\theta_n}\}$  is independent.

With this device, we may make claims such as

$$\{X_t : t \in (0, 1]\}$$

are independent. Useful for stochastic process, functional data analysis.

It follows trivially,  $\{\mathcal{A}_\theta : \theta \in \Theta\}$  are independent if and only if any finite collection, say  $\{\mathcal{A}_{\theta_1}, \dots, \mathcal{A}_{\theta_n}\}$  are independent.

**Corollary 1.4.3 — To Theorem 4.2.** If  $(\Omega, \mathcal{F}, P)$ ,  $\mathcal{A}_\theta \subset \mathcal{F}$ ,  $\{\mathcal{A}_\theta : \theta \in \Theta\}$  is independent and each  $\mathcal{A}_\theta$  is a  $\pi$ -system, then

$$\{\sigma(\mathcal{A}_\theta) : \theta \in \Theta\}$$

are independent.



*Proof.*

$$\begin{aligned}\{\mathcal{A}_\theta : \theta \in \Theta\} \perp\!\!\!\perp &\Leftrightarrow \{\mathcal{A}_{\theta_1}, \dots, \mathcal{A}_{\theta_n}\} \perp\!\!\!\perp \\ &\Leftrightarrow \{\sigma(\mathcal{A}_{\theta_1}), \dots, \sigma(\mathcal{A}_{\theta_n})\} \perp\!\!\!\perp\end{aligned}$$

■

**Corollary 1.4.4** Suppose we have an array of sets,

$$\begin{array}{cccc} A_{11} & A_{12} & \dots & \dots \\ A_{21} & A_{22} & \dots & \dots \\ \vdots & \vdots & & \\ \vdots & \vdots & & \end{array} = \{A_{ij} : i, j = 1, \dots\} \subset \mathcal{F}$$

and this array is independent.

And let  $\mathcal{F}_i = \sigma(A_{i1}, A_{i2}, \dots)$ .

Then  $\mathcal{F}_1 \perp\!\!\!\perp \mathcal{F}_2$

*Proof.* Let  $\mathcal{A}_i$  be the class of all the finite intersections of

$$A_{i1}, A_{i2}, \dots$$

then  $\mathcal{A}_i$  is a  $\pi$ -system.

So,

$$\sigma(\mathcal{A}_i) = \mathcal{F}_i$$

because  $\{A_{i1}, A_{i2}, \dots\}$  are contained in  $\mathcal{A}_i$  which implies  $\mathcal{F}_i \subset \sigma(\mathcal{A}_i)$  and also  $\mathcal{A}_i \subset \mathcal{F}_i \Rightarrow \sigma(\mathcal{A}_i) \leq \mathcal{F}_i$ .

By Corollary 1, it suffices to show that  $\mathcal{A}_1, \mathcal{A}_2, \dots$  are independent. Further, it suffices to show that any finite subcollection is independent.

Let  $\{i_1, \dots, i_n\} \subseteq \{1, 2, \dots\}$ . We want to show that  $\mathcal{A}_{i_1} \perp\!\!\!\perp \dots \perp\!\!\!\perp \mathcal{A}_{i_n}$ . This would be implied by the following

$$\forall C_{i_1} \in \mathcal{A}_{i_1}, \dots, C_{i_n} \in \mathcal{A}_{i_n}$$

$$C_{i_1} \perp\!\!\!\perp \dots \perp\!\!\!\perp C_{i_n}$$

Because, and watch out with this notation here, for

$$C_{i_\alpha} \in \mathcal{A}_{i_\alpha}$$

there exists

$$A_{i_\alpha j_1}, A_{i_\alpha j_2}, \dots, A_{i_\alpha j_{m_\alpha}}$$

such that

$$C_{i_\alpha} = A_{i_\alpha j_1}, A_{i_\alpha j_2}, \dots, A_{i_\alpha j_{m_\alpha}}$$

We have

$$P(\cap_{\alpha=1}^n C_{i_\alpha}) = P(\cap_{\alpha=1}^n \bigcup_{\beta=1}^{m_\alpha} A_{i_\alpha j_\beta})$$

because

$$\{A_{i_\alpha j_\beta} : \alpha = 1, 2, \dots, n, \beta = 1, 2, \dots, m_\alpha\} \subseteq \{A_{ij} : i, j = 1, 2, \dots\}$$

$$\begin{aligned} P(\cap_{\alpha=1}^n \bigcup_{\beta=1}^{m_\alpha} A_{i_\alpha j_\beta}) &= \prod_{\alpha=1}^n \prod_{\beta=1}^{m_\alpha} P(A_{i_\alpha j_\beta}) \\ &= \prod_{\alpha=1}^n P(C_{i_\alpha}) \end{aligned}$$

■

### Borel-Cantelli Lemmas (that are actually Theorems)

**Theorem 1.4.5 — BC1.** For  $(\Omega, \mathcal{F}, P)$  probability space,

$$A_n \in \mathcal{F}, \quad n = 1, 2, \dots$$

If  $\sum_{n=1}^{\infty} P(A_n) < +\infty$  then

$$P(\limsup_{n \rightarrow \infty} A_n) = 0$$

*Proof.*  $\limsup_{n \rightarrow \infty} A_n = \cap_{n=1}^{\infty} \bigcup_{k=n}^{\infty} A_k \subset \bigcup_{k=1}^{\infty} P(A_k) \quad \forall n$   
So,

$$P(\limsup_{n \rightarrow \infty} A_n) \leq P(\bigcup_{k=1}^{\infty} P(A_k)) \leq \sum_{k=1}^{\infty} P(A_k)$$

■

**Theorem 1.4.6 — BC2.** If  $\{A_n\}$  are independent and  $\sum_{n=1}^{\infty} P(A_n) = \infty$ , then

$$P(\limsup_n A_n) = 1$$

*Proof.*  $P(\limsup_n A_n) = 1$

$$\Leftrightarrow P(\bigcap_{n=1}^{\infty} \bigcup_{k=n}^{\infty} A_k) = 1$$

$$\Leftrightarrow P(\bigcup_{n=1}^{\infty} \cap_{k=n}^{\infty} A_k^C) = 0 \quad (*)$$

because,

$$\Leftrightarrow P\left(\bigcup_{n=1}^{\infty} \bigcap_{k=n}^{\infty} A_k^C\right) \leq \sum_{k=n}^{\infty} P(\bigcap_{k=n}^{\infty} A_k^C)$$

$$\Leftrightarrow P(\bigcap_{k=n}^{\infty} A_k^C) = 0 \quad \forall n = 1, 2, \dots$$

but we need to prove this to imply (\*).

Shit, calculus.

$$1 - x \leq e^{-x} \quad \forall x \in \mathbb{R}$$

For any  $j = 1, 2, \dots$ ,

$$\begin{aligned} P\left(\bigcap_{k=n}^{n+j} A_k^C\right) &= \prod_{k=n}^{n+j} (1 - P(A_k)) \\ &\leq \prod_{k=n}^{n+j} e^{-P(A_k)} \\ &= e^{-\sum_{k=n}^{n+j} P(A_k)} \end{aligned}$$

Now,  $\sum_{k=1}^{\infty} P(A_k) = \infty$  and also

$$\sum_{k=n}^{\infty} P(A_k) \quad \forall n$$

So,

$$\lim_{j \rightarrow \infty} \sum_{k=n}^{n+j} P(A_k) \rightarrow \infty \quad \forall n$$

$$\lim_{j \rightarrow \infty} P\left(\bigcap_{k=n}^{n+j} A_k^C\right) = 0$$

Because,

$$\bigcap_{k=n}^{n+j} A_k^C \downarrow \bigcap_{k=n}^{\text{infy}} A_k^C \quad j \rightarrow \infty$$

By continuity of probability,

$$P\left(\bigcap_{k=n}^{n+j} A_k^C\right) \downarrow P\left(\bigcap_{k=n}^{\text{infy}} A_k^C\right) \quad j \rightarrow \infty$$

So,

$$P\left(\bigcap_{k=n}^{\text{infy}} A_k^C\right) = 0$$

■

Friday September 16

finished proof.

BC1 and BC@ say that  $P(\limsup_n A_n)$  is either 0 or 1.

This is a special case of a general phenomenon, the 0-1 Law.

Take  $\sigma$ -field,  $(\Omega, \mathcal{F}, P)$ ,

$$A_1, \dots \in \mathcal{F}$$

For each  $n$ ,

$$\sigma(A_n, A_{n+1}, \dots)$$

We have another  $\sigma$ -field called "tail of  $\sigma$ -field",

$$\mathcal{T} = \bigcap_{n=1}^{\infty} \sigma(A_n, A_{n+1}, \dots)$$

■ **Example 1.4 — 4.18 in Billingsly.**  $\limsup A_n \in \mathcal{T}$ ?

$$\bigcap_n \bigcup_{k=n}^{\infty} A_k$$

$$A_n, A_{n+1}, \dots \in \sigma(A_n, A_{n+1}, \dots) \Rightarrow \bigcup_{k=n}^{\infty} A_k \in \sigma(A_n, A_{n+1}, \dots)$$

$$\bigcap_n \bigcup_{k=n}^{\infty} A_k \in \bigcap_{n=1}^{\infty} \sigma(A_n, A_{n+1}, \dots)$$

$$\begin{aligned} \liminf A_n &= \left[ \bigcup_n \bigcap_{k=n}^{\infty} A_k \right]^C \\ &= \left[ \bigcup_n \bigcap_{k=n}^{\infty} A_k^C \right]^C \\ &= \left[ \limsup_n A_k^C \right]^C \in \mathcal{T} \end{aligned}$$

■

**Theorem 1.4.7** If  $A_1, A_2, \dots$  are independent, then for each  $A \in \mathcal{T}$  we have  $P(A) = 0$  or 1.

*Proof.* By Corollary 2,

$$\begin{aligned} &\sigma(A_1) \\ &\sigma(A_2) \\ &\vdots \\ &\vdots \\ &\sigma(A_{n-1}) \\ &\sigma(A_n, A_{n+1}, \dots) \end{aligned}$$

$$\sigma(A_1) \perp \dots \perp \sigma(A_n, A_{n+1}, \dots)$$

Let  $A \in \mathcal{F}$ , then,

$$A \in \sigma(A_n, A_{n+1}, \dots) \quad \forall n$$

So,  $A_1, \dots, A_{n-1}, A$  are independent. By taking  $n$  large enough, this implies that any finite subcollection of  $A, A_1, A_2, \dots$  is also independent.

This implies that the sequence  $\{A, A_1, A_2, \dots\}$  are independent.

But  $A \in \sigma(A_1, A_2, \dots)$  so,

$$A \perp\!\!\!\perp A$$

$$P(AA) = P(A)P(A) = P(A)^2$$

So  $P(A)$  must be zero or 1!

■

**R** We are now skipping a few sections (5 -9) in Billingsly. These are about special random variables, random walks, etc...

## 1.5 General Measure on a Field

### Borel Sets in $\mathbb{R}^k$

Two jumps, from  $(0, 1] \rightarrow \mathbb{R} \rightarrow \mathbb{R}^k$ .

$\mathcal{B}$  on  $(0, 1]$  is a  $\sigma$ -field generated by  $\mathcal{I}$  = collection of all intervals in  $(0, 1]$ .

$$\sigma(\mathcal{I}) = \mathcal{B} \text{ on } (0, 1]$$

When we work with  $\mathbb{R}$ ,

$$\mathcal{I}' = \text{collection of all intervals in } \mathbb{R}, (a, b)$$

$$\sigma(\mathcal{I}') = \mathcal{R}' \quad \text{linear Borel } \sigma\text{-field}$$

$\mathcal{I}^k$  is the collection of all rectangles in  $\mathbb{R}^k$ .

$$\mathcal{I}^k = \{(a_1, b_1] \times \dots \times (a_k, b_k] : a_1, b_1, \dots, a_k, b_k \in \mathbb{R}\}$$

$$\sigma(\mathcal{I}^k) = \mathcal{R}^k \quad \text{Borel } \sigma\text{-field in } \mathbb{R}^k$$

*Properties of  $\mathbb{R}^k$*

(\*) Any open set are in  $\mathcal{R}^k$ .

Let  $\mathbb{Q}$  be the set of all rational numbers. This is countable and dense subset of  $\mathbb{R}$ .

**Definition 1.5.1 — Dense.** Look up definition!

Class of rational rectangles:

$$\mathcal{J}_{\mathbb{Q}}^k = \{(a_1, b_1]x \dots x(a_k, b_k] : a_1, b_1, \dots, a_k, b_k \in \mathbb{Q}\}$$

Let  $G$  be an open set in  $\mathbb{R}^k$  and  $y \in G$ , then there exists

$$A_y \in \mathcal{J}_{\mathbb{Q}}^k$$

such that

$$y \in A_y \subset G$$

because  $\mathbb{Q}$  is dense in  $\mathbb{R}$ .

Note that,  $\bigcup_{y \in G} A_y = G$ .

But,  $\{A_y : y \in G\} \subseteq \mathcal{J}_{\mathbb{Q}}^k$ .

### Monday September 19

Note that the above union of  $A_y$  is countable.

Also,  $G = \bigcup_{y \in G} A_y$  and so  $G \in \sigma(\mathcal{J}_{\mathbb{Q}}^k) \subseteq \text{sigma}(\mathcal{J}_{\mathbb{Q}}) = \mathcal{R}^k$ .

Immediately, we see that

(\*) All closed sets  $F$  are in  $\mathcal{R}^k$ .

All sets we commonly see are in  $\mathcal{R}^k$ .

(\*)  $\mathcal{R}^k$  is in fact also the  $\sigma$ -field generated by the class of all open sets in  $\mathbb{R}^k$ ,  $\mathcal{G}^k$ .

*Proof.* Let

$$\mathcal{J}^k = \{(a_1, b_1)x \dots x(a_k, b_k) : a_1, b_1, \dots, a_k, b_k \in \mathbb{R}, k = 1, 2, \dots\}$$

Claim:  $\sigma(\mathcal{J}^k) \in \mathcal{R}^k$

Note that any

$$(a_1, b_1]x \dots x(a_k, b_k] \in \mathcal{J}^k$$

can be written as

$$\bigcap_{n=1}^{\infty} (a_1, b_1 + n^{-1}]x \dots x(a_k, b_k + n^{-1}]$$

The above statement is within  $\mathcal{J}^k$  with the intersection, otherwise it'd be in  $\mathcal{J}^k$ .

That means each  $A \in \mathcal{J}^k$  is in  $\sigma(\mathcal{J}^k)$ .

$$\mathcal{J}^k \subseteq \sigma(\mathcal{J}^k)$$

$$\sigma(\mathcal{I}^k) \subseteq \sigma(\mathcal{J}^k)$$

$$\mathcal{R}^k \subset \sigma(\mathcal{J}^k)$$

But since  $\mathcal{R}^k$  contains all open sets (previous (\*)), we have that  $\mathcal{R}^k \supseteq \mathcal{J}^k$  and  $\mathcal{R}^k \supseteq \sigma(\mathcal{J}^k)$ . So,

$$\mathcal{R}^k = \sigma(\mathcal{J}^k)$$

Our claim is proved.

Because  $\mathcal{R}^k \supseteq \mathcal{G}^k$ ,

$$\mathcal{R}^k \supseteq \sigma(\mathcal{G}^k)$$

But,  $\mathcal{J}^k \subset \mathcal{G}^k$

$$\mathcal{R}^k = \sigma(\mathcal{J}^k) \subset \sigma(\mathcal{G}^k)$$

So,

$$\mathcal{R}^k = \sigma(\mathcal{G}^k)$$

And this is the general definition of Borel  $\sigma$ -field, because open sets exists much more generally than rectangles.

In fact, Borel Sets in Hilbert spaces, Banach...etc. wherever you can define open sets, you can define Borel Sets.

■

### Borel Sets in Topological Space

**Definition 1.5.2 — Topology.** A paving  $\mathcal{T}$  is a **topology** on  $\Omega$  if it is closed under arbitrary union of finite intersections. That is, if you have an arbitrary index,

1. If  $\{A_\theta : \theta \in \Theta\} \subseteq \mathcal{T}$  then the

$$\bigcup_{\theta \in \Theta} A_\theta \in \mathcal{T}$$

2. If  $A, B \in \mathcal{T}, AB \in \mathcal{T}$  then any set  $A \in \mathcal{T}$  is called an open set with respect to  $\mathcal{T}$ .

$(\Omega, \mathcal{T}) \leftarrow$  Topological Space

$\sigma(\mathcal{T}) \leftarrow$  Borell  $\sigma$ -field generated by  $\mathcal{T}$ -open sets.

$(\Omega, \sigma(\mathcal{T}))$  is measureable.

Here is another question:

$$\mathcal{B}, \mathcal{R}', \mathcal{R}^*$$

Is it reasonalbe to conjecture to following?

$$\{A \subseteq \mathcal{R}' : A \subseteq (0, 1]\} = \mathcal{B}$$

### $\sigma$ -field Restricted on a Set

Let  $(\Omega, \mathcal{F})$  be a measure space.

$$\Omega_0 \subseteq \Omega$$

(otherwise arbitrary, especially  $\Omega_0$  need not be in  $\mathcal{F}$ )

Define (with some "lazy" notation),

$$\mathcal{F} \cap \Omega_0 = \{A \cap \Omega_0 : A \in \mathcal{F}\}$$

**Theorem 1.5.1** (i)  $\mathcal{F} \cap \Omega_0$  is a  $\sigma$ -field in  $\Omega_0$   
(ii) If  $\mathcal{A}$  generates  $\mathcal{F}$  then,  $A \cap \Omega_0$  generates  $\mathcal{F} \cap \Omega_0$

*Proof.* (i)  $\mathcal{F} \cap \Omega_0$  is a  $\sigma$ -field in  $\Omega_0$

Want to show:  $\Omega_0 \in \mathcal{F} \cap \Omega_0$

$$\begin{aligned} \Omega &\in \mathcal{F} \\ \Omega_0 &= \Omega \cap \Omega_0 \in \mathcal{F} \cap \Omega_0 \end{aligned}$$

Want to show:  $A \in \mathcal{F} \cap \Omega_0 \rightarrow A \in \mathcal{A}^C \in \mathcal{F} \cap \Omega_0$

$$\begin{aligned} A \in \mathcal{F} \cap \Omega_0 &\Rightarrow A = B \cap \Omega_0, B \in \mathcal{F} \\ B \in \mathcal{F} &\Rightarrow B^c \in \mathcal{F} \end{aligned}$$

So,

$$B^c \cap \Omega_0 \in \mathcal{F} \cap \Omega_0$$

But,

$$\begin{aligned} \Omega_0 \setminus A &= \Omega_0 \setminus (B \cap \Omega_0) \\ &= \Omega_0 \cap (B \cap \Omega_0)^c \\ &= \Omega_0 \cap (B^c \cup \Omega_0^c) \\ &= (\Omega_0 \cap B^c) \cup (\Omega_0 \cap \Omega_0^c) \\ &= (\Omega_0 \cap B^c) \cup \emptyset \\ &= (\Omega_0 \cap B^c) \in \mathcal{F} \cap \Omega_0 \end{aligned}$$

Want to show:  $A_1, \dots \in \mathcal{F} \cap \Omega_0 \Rightarrow \bigcup_{n=1}^{\infty} A_n \in \mathcal{F} \cap \Omega_0$

$$A_n \in \mathcal{F} \cap \Omega_0 \Rightarrow A_n = B_n \cap \Omega_0, B_n \in \mathcal{F}$$



$$\begin{aligned}
\bigcup_n B_n \in \mathcal{F} &\Rightarrow (\bigcup_n B_n)\Omega_0 \in \mathcal{F} \cap \Omega_0 \\
\text{But,} \quad &\Rightarrow \bigcup_n (B_n \Omega_0) \in \mathcal{F} \cap \Omega_0 \\
&\Rightarrow \bigcup_{n=1}^{\infty} A_n \in \mathcal{F} \cap \Omega_0
\end{aligned}$$

So the three things we've just shown gives us that  $\mathcal{F} \cap \Omega_0$  is in deed a  $\sigma$ -field.

(ii) If  $\mathcal{A}$  generates  $\mathcal{F}$  then,  $\mathcal{A} \cap \Omega_0$  generates  $\mathcal{F} \cap \Omega_0$

Let  $\mathcal{A} \subseteq \mathcal{F}$ ,  $\sigma(\mathcal{A}) = \mathcal{F}$ .

Let  $\mathcal{F}_0 = \sigma(\mathcal{A} \cap \Omega_0)$

Our goal:  $\mathcal{F} \cap \Omega_0 = \sigma(\mathcal{A})$

Step 1:  $\mathcal{F}_0 \subseteq \mathcal{F} \cap \Omega_0$

$$\mathcal{A} \cap \Omega_0 \subset \mathcal{F} \cap \Omega_0$$

$$\mathcal{F} \cap \Omega_0 \text{ is } \sigma\text{-field}$$

Step 2:  $\mathcal{F} \cap \Omega_0 \subset \mathcal{F}_0$  holds if  $[A \in \mathcal{F} \Rightarrow A\Omega_0 \in \mathcal{F}]$

$$\Rightarrow \bigcap \Omega_0 \subset \mathcal{F}_0$$

If bracket statement is true, then  $\mathcal{F} \cap \Omega_0 \subset \mathcal{F}$ .

Let  $A \in \mathcal{F} \cap \Omega_0$  then

$$A = B\Omega_0, B \in \mathcal{F}$$

But by bracket statement,

$$B \in \mathcal{F} \Rightarrow B\Omega_0 \in \mathcal{F} \Rightarrow A \in \mathcal{F}_0$$

So,  $\mathcal{F} \cap \Omega_0 \subset \mathcal{F}_0$ .

Step 3: Let  $\mathcal{G} = \{A \subset \Omega : A\Omega_0 \in \mathcal{F}_0\}$ . Then,

$$\mathcal{A} \subseteq \mathcal{G}$$

Pick  $A \in \mathcal{A}$ .

$$\Rightarrow A\Omega_0 \in \mathcal{A} \cap \Omega_0 \subset \mathcal{F}_0$$

$$\Rightarrow A \in \mathcal{G}$$

So,  $\mathcal{A} \subset \mathcal{G}$ .

Step 4:  $\mathcal{G}$  is  $\sigma$ -field in  $\Omega$ .

(a)

$$\Omega \in \mathcal{G} : \Omega\Omega_0 = \Omega_0 \in \mathcal{F}_0 = \sigma(\mathcal{A} \cap \Omega_0)$$

Generally, if  $\Omega$  is a set  $\mathcal{B}$  is a pvaing on  $\Omega$  then  $\sigma(\mathcal{B}) = \bigcap \{\mathcal{B}' : \mathcal{B}' \supseteq \mathcal{B}, \mathcal{B}' \text{ is a } \sigma\text{-field on } \Omega\}$ .

Which is true because  $\mathcal{A}$  generates  $\mathcal{F}$ ,  $\Omega \in \mathcal{A}$ .

This means that

$$\sigma(\mathcal{A} \cap \Omega_0) = \bigcap \{ \mathcal{B} : \mathcal{B} \supset \mathcal{A} \cap \Omega_0, \mathcal{B} \text{ is a } \sigma\text{-field on } \Omega \}$$

So,  $\Omega_0 \in \mathcal{F}_0$

(b)  $A \in \mathcal{G} \Rightarrow A^C \in \mathcal{G}$ .

$$A \in \mathcal{G} \Rightarrow A\Omega_0 \in \mathcal{F}_0$$

$$\Rightarrow \Omega_0 \setminus (A\Omega_0) \in \mathcal{F}_0$$

$$\Rightarrow \Omega_0 \cap (A\Omega_0)^C \in \mathcal{F}_0$$

$$\Rightarrow \Omega_0 \cap (A^C \Omega_0^C) \in \mathcal{F}_0$$

$$\Rightarrow (\Omega_0 A^C) \cup (\Omega_0 \Omega_0^C)$$

$$= \Omega_0 A^C \in \mathcal{F}_0$$

So,  $A^C \in \mathcal{F}$ .

(c)  $A_1, A_2, \dots \in \mathcal{G}$  are disjoint means  $A_n \Omega_0 \in \mathcal{F}_0$  and  $A_n \Omega_0$  disjoint.

$$\bigcup_{n=1}^{\infty} (A_n \Omega_0) \in \mathcal{F}$$

$$(\bigcup_{n=1}^{\infty} A_n) \Omega_0 \in \mathcal{F}$$

$$\bigcup_{n=1}^{\infty} A_n \in \mathcal{G}$$

Step 5:  $\mathcal{F} \cap \Omega_0 \subseteq \mathcal{F}_0$

By Step 3, we know that  $\mathcal{A} \subset \mathcal{G}$ .

By Step 4,  $\mathcal{G}$  is  $\sigma$ -field.

Together,  $\sigma(\mathcal{A}) = \mathcal{G}$ .

$$\sigma(\mathcal{A}) = \mathcal{F} \in \mathcal{G}$$

$$\Rightarrow [A \in \mathcal{F} \Rightarrow A\Omega_0 \in \mathcal{F}]$$

$$\Rightarrow \bigcap \Omega_0 \subset \mathcal{F}_0$$

■

**Wednesday September 21**

Worked on proof, but will need to go back to it in the future.

So we have now that  $\sigma(\mathcal{A} \cap \Omega_0) = \sigma(\mathcal{A} \setminus \bigcap \Omega)$

**Lemma 1.** If  $\Omega_0 \in \mathcal{F}$  then

$$\mathcal{F} \cap \Omega_0 = \{B \in \mathcal{F} : B \subset \Omega_0\}$$

*Proof.* Need to show that  $\{A\Omega_0 : A \in \mathcal{F}\} = \{B \in \mathcal{F} : B \subset \Omega_0\}$

Let  $C \in LHS$ .

$$C = A\Omega_0, A \in \mathcal{F}$$

then  $C \in \mathcal{F}, C \subset \Omega_0$ .

So  $C \in RHS$ .

Let  $B \in RHS$ ,

$$B \subset \Omega_0, B \in \mathcal{F}$$

$$B = B \bigcap \Omega_0, B \in \mathcal{F}$$

So,  $B \in LHS$ . ■

**Corollary 1.5.2**  $\mathcal{B} = \{A \subset (0, 1] : A \in \mathcal{R}'\} = \{A \in \mathcal{R}' : A \in (0, 1]\}$

*Proof.* Let  $\Omega \in \mathbb{R}$ .

$$\Omega_0 = (0, 1]$$

$$\mathcal{F} = \mathcal{R}'$$

$$\perp = \mathcal{I}'$$

By Theorem 10.1,

$$\sigma(\mathcal{A}) \bigcap \Omega_0 = \sigma(\mathcal{A} \bigcap \Omega)$$

$$\Leftrightarrow \sigma((\mathcal{I}') \bigcap (0, 1]) = \sigma(\mathcal{I}' \bigcap (0, 1]) = \sigma(\mathcal{I}) = \mathcal{B}$$

Now  $(0, 1] \in \mathcal{R}' \Leftrightarrow \Omega_0 \in \mathcal{F}$ .

By Lemma 1,

$$\Rightarrow \mathcal{F} \bigcap \Omega_0 = \{A \in \mathcal{F} : A \subset \Omega_0\}$$

$$\Rightarrow \mathcal{R} \bigcap (0, 1] = \{A \in \mathcal{R} : A \subset (0, 1]\}$$

So,

$$\mathcal{B} = \{A \in \mathcal{R} : A \subset (0, 1]\}$$
■

For general measure, need infinity convention.

For

$$x, y \in [0, \infty] = [0, \infty) \cup \{\infty\}$$

$x \leq y$  means that (either/or)

1.  $y = \infty$
  2.  $y < \infty, x < \infty, x \leq y$
- $x < y$  means that (either/or)

1.  $y = \infty, x < \infty$
2.  $x, y < \infty, x < y$

For a finite or infinite sequence,  $x, x_1, \dots \in [0, \infty]$ ,

$x = \sum_{k=1}^{\infty} x_k$  means that (either/or)

1.  $x = \infty, x_k = \infty$  for some  $k$
2.  $x = \infty, x < \infty \forall k$   
 $\sum_{k=1}^n x_k : n = 1, 2, \dots$  diverges.
3.  $x < \infty, x_k < \infty$

$$\lim_{n \rightarrow \infty} \sum_{k=1}^n x_k = x$$

**Friday September 23**

For any infinite sequence,  $x_1, x_2, \dots \in [0, \infty]$  and  $x \in [0, \infty]$

$x_k \uparrow x$  is true if and only if

1.  $x_k \leq x_{k+1} \leq x \quad \forall k$
2. either
  - (a)  $x < \infty$ , and  $x_k \uparrow x$  in usual sense.
  - (b)  $x_k = \infty$  for  $\forall k \geq m, x = \infty$
  - (c)  $x = \infty, x_k < \infty, x_k \uparrow \infty$

**Measures on Field**

Let  $\Omega \leftarrow$  nonempty and  $\mathcal{F}$  be a field on  $\Omega$ .

**Definition 1.5.3** A measure,  $\mu$  is a function on  $\mathcal{F}$  such that

1.  $\mu(A) \in [0, \infty]$
2.  $\mu(\emptyset) = 0$
3. If  $A_n \in \mathcal{F}$  are disjoint then

$$\mu\left(\bigcup_{n=1}^{\infty} A_n\right) = \sum_{n=1}^{\infty} \mu(A_n)$$

A measure is finite if  $\mu(\Omega) < \infty$ .

A measure is a probability measure or simply probability if  $\mu(\Omega) = 1$ .

A measure is  $\tau$ -finite if there exists  $\mathcal{F}$ -sets  $A_n$  such that

$$\bigcup_{n=1}^{\infty} A_n = \Omega, \mu(A_n) < \infty$$

If  $\mathcal{F}$  is a  $\sigma$ -field then  $(\Omega, \mathcal{F})$  is a measurable space.

If  $\mu$  is a measure on  $\mathcal{F}$  then  $(\Omega, \mathcal{F}, \mu)$  is a measure space.

**Definition 1.5.4 — Support.** If  $A \in \mathcal{F}, \mu(A^C) = 0$  then  $A$  is a support of  $\mu$ .

A measure,  $\mu$  on  $(\Omega, \mathcal{F})$  has the following properties (proof similar to probability case omitted).

*Finite additivity:*  $A_1, \dots, A_n$  are disjoint  $\mathcal{F}$ -sets implies

$$\mu\left(\bigcup_{i=1}^{\infty} A_i\right) = \sum_{i=1}^{\infty} \mu(A_i)$$

*Monotonicity:* If  $A, B \in \mathcal{F}, A \subseteq B$  then

$$\mu(A) \leq \mu(B)$$

*Inclusion-Exclusion Formula:*  $A_1, \dots, A_n \in \mathcal{F}_1$

$$\mu\left(\bigcup_{k=1}^{\infty} A_k\right) = \sum_{k=1}^{\infty} \mu(A_k) - \sum_{i=k < l=n} \mu(A_k A_l) + \dots + (-1)^{n+1} \mu(A_1 \dots A_k)$$

*Countable or Finite Subadditivity:*  $A_1, \dots \in \mathcal{F}$ , then

$$\mu\left(\bigcup_{n=1}^{\infty} A_n\right) \leq \sum_{n=1}^{\infty} \mu(A_n)$$

**Theorem 1.5.3** Let  $\mu$  be a measure on a  $\sigma$ -field,  $\mathcal{F}$ .

1. Continuity from below.

$$A_n, A \in \mathcal{F}, A_n \uparrow A \Rightarrow \mu(A_n) \uparrow \mu(A)$$

2. Continuity from above.

$$A_n, A \in \mathcal{F}, \mu(A_1) < \infty, A_n \downarrow A \Rightarrow \mu(A_n) \rightarrow \mu(A)$$

3. Countable subadditivity.

$$A_n \in \mathcal{F}, \bigcup_n \in \mathcal{F} \Rightarrow \mu\left(\bigcup_{n=1}^{\infty} A_n\right) \leq \sum_{n=1}^{\infty} \mu(A_n)$$

4. If  $\mu$  is  $\sigma$ -finite on  $\mathcal{F}$  then  $\mathcal{F}$  contain an uncountable, disjoint collection of sets with positive  $\mu$ -measure.

*Proof.* (i) and (iii) exactly the same as in probability case.

(ii) takes a little extra work.

If  $\mu(A_1) < \infty$  then

$$A_1 \setminus A_n \uparrow A_1 \setminus A$$

then by (i) we have

$$\mu(A_1 \setminus A_n) \uparrow \mu(A_1 \setminus A)$$

$$\mu(A_1) - \mu(A_n) \uparrow \mu(A_1) - \mu(A)$$

$$\mu(A_n) \downarrow \mu(A)$$

(iv) Let  $\{B_\theta : \theta \in \Theta\}$  be a disjoint collection of  $\mathcal{F}$ -sets such that  $\mu(B_\theta) > 0$ . We want to show it is countable.

Claim: If  $A \in \mathcal{F}$  and  $\mu(A) < \infty, \varepsilon > 0$ . then,

$\{\theta : \mu(AB_\theta) > \varepsilon\}$  is finite.

Show: If  $\{\theta : \mu(AB_\theta) > \varepsilon\}$  is infinite then there exists  $\theta_1, \theta_2, \dots$  such that  $\mu(AB_{\theta_i}) > \varepsilon \quad \forall \theta_i$ .

But, if this were so,

$$\sum_{i=1}^n \mu(AB_{\theta_i}) \geq n\varepsilon$$

$$\sum_{i=1}^n \mu(AB_{\theta_i}) \geq \mu(A)$$

for sufficiently large  $n$ . CONTRADICTION.

But,  $\{\theta : \mu(AB_\theta) > 0\} = \bigcup_{r \in \mathbb{Q}} \{\theta : \mu(AB_\theta) > r\}$  is a countable union of a finite set.

So  $\{\theta : \mu(AB_\theta) > 0\}$  is countable. Now we just need to show  $\{\theta : \mu(B_\theta) > 0\}$  is also countable.

But we know that  $\{\theta : \mu(\Omega B_\theta) > 0\}$  is countable because  $\mu$  is  $\sigma$ -finite so there exists  $A_n \in \mathcal{F}, n = 1, 2, \dots$  such that  $\mu(A_n) < \infty, \bigcup_{n=1}^{\infty} A_n = \Omega$ .

Want to show  $\{\theta : \mu(\bigcup_{n=1}^{\infty} A_n B_\theta) > 0\}$  is countable.

Note that

$$\mu\left(\bigcup_{n=1}^{\infty} A_n B_\theta\right) \leq \mu\left(\sum_{n=1}^{\infty} A_n B_\theta\right)$$

So,

$$\{\theta : \mu\left(\bigcup_{n=1}^{\infty} A_n B_\theta\right) > 0\} \subseteq \{\theta : \mu\left(\sum_{n=1}^{\infty} A_n B_\theta\right) > 0\}$$

The RHS may be rewritten as

$$\bigcup_{n=1}^{\infty} \{\theta : \mu\left(\bigcup_{n=1}^{\infty} A_n B_\theta\right) > 0\}$$

■

**Monday September 26**

**Uniqueness of Extension**

That is, if the extension exists (we'll explore later), then is it unique?

**Theorem 1.5.4** Suppose  $\mu_1, \mu_2$  are measures on  $\sigma(\mathcal{P})$ , where  $\mathcal{P}$  is a  $\pi$ -system of subsets of  $\Omega$ . Suppose  $\mu_1, \mu_2$  are  $\sigma$ -finite on  $\mathcal{P}$ . If  $\mu_1, \mu_2$  agree on  $\mathcal{P}$  (meaning  $\mu_1(A) = \mu_2(A) \forall A \in \mathcal{P}$ ) then they agree on  $\sigma(\mathcal{P})$ .

*$\sigma$ -Finite on  $\mathcal{P}$*

Let  $(\Omega, \mathcal{F}, \mu)$  be a measure space,  $\mathcal{A} \subseteq \mathcal{F}$ . We say  $\mu$  is  $\sigma$ -finite on  $\mathcal{A}$  if there exists  $\{A_n\}_{n=1}^\infty, A_n \in \mathcal{A}$  such that

$$\mu(A_n) < \infty, \Omega = \bigcup_{n=1}^\infty A_n$$

*Proof.* Idea: extend the parallel theorem in probability case.

Step 1: Prove that if  $B \in \mathcal{P}, \mu_1(B) = \mu_2(B) < \infty$  then

$$\mu_1(BA) = \mu_2(BA) \quad \forall A \in \sigma(\mathcal{P})$$

Let  $\mathcal{L}_B = \{A \in \sigma(\mathcal{P}) : \mu_1(BA) = \mu_2(BA) < \infty\}$ .

Need to prove  $\mathcal{L}_B$  is a  $\lambda$ -system.

1.  $\Omega \in \mathcal{L}_B$ .

We know that  $\Omega \in \sigma(\mathcal{P})$ , because  $\sigma(\mathcal{P})$  is itself a  $\sigma$ -field. Thus,

$$\mu_1(B\Omega) = \mu_1(B) = \mu_2(B) = \mu_2(B\Omega)$$

and so we have that  $\Omega \in \mathcal{L}_B$ .

2. Want  $A \in \mathcal{L}_B \Rightarrow A^C \in \mathcal{L}_B$ .

$$\begin{aligned} A \in \mathcal{L}_B &\Rightarrow \mu_1(AB) = \mu_2(AB) \\ &\Rightarrow \mu_1(B) - \mu_1(B \setminus (AB)) = \mu_2(B) - \mu_2(B \setminus (AB)) \\ &\Rightarrow \mu_1(B \setminus (AB)) = \mu_2(B \setminus (AB)) \\ &\Rightarrow \mu_1(BA^C) = \mu_2(BA^C) \end{aligned}$$

We have that  $A^C \in \mathcal{L}_B$ .

3. Want that if  $A_n$  are disjoint  $\mathcal{L}_B$ -sets  $\Rightarrow \bigcup_{n=1}^\infty A_n \in \mathcal{L}_B$ .

$$A_n \in \mathcal{L}_B : \mu_1(A_n B) = \mu_2(A_n B)$$

$$\sum_{n=1}^\infty \mu_1(A_n B) = \sum_{n=1}^\infty \mu_2(A_n B)$$

Because the  $A_n$  are disjoint, we have that  $BA_n$  are also disjoint.

$$\begin{aligned} \mu_1\left(\bigcup_{n=1}^\infty (A_n B)\right) &= \mu_2\left(\bigcup_{n=1}^\infty (A_n B)\right) \\ \mu_1\left(\left(\bigcup_{n=1}^\infty A_n\right) B\right) &= \mu_2\left(\left(\bigcup_{n=1}^\infty A_n\right) B\right) \end{aligned}$$

So,  $\bigcup_n A_n \in \mathcal{L}_B$ . And thus  $\mathcal{L}_B$  is a  $\lambda$ -system.

But,  $\mathcal{L}_B \supseteq \mathcal{P}$  by definition. So  $\mathcal{L}_B \supseteq \sigma(\mathcal{P})$  by the  $\pi - \lambda$  Theorem. So we have

$$\mu_1(AB) = \mu_2(AB) \quad \forall A \in \sigma(\mathcal{P})$$

and step 1 is finished.

Step 2: Want  $\mu_1(A) = \mu_2(A) \forall A \in \sigma(\mathcal{P})$ . Or rather  $\mu_1(A\Omega) = \mu_2(A\Omega) \forall A \in \sigma(\mathcal{P})$ , but  $\Omega$ , unlike  $B$ , may not have  $\mu(\Omega) < \infty$ .

Because  $\mu$  is  $\sigma$ -finite on  $\mathcal{P}$  there exists  $B_n \in \mathcal{P}, \mu(B_n) < \infty$  such that  $\bigcup_n B_n = \Omega$ .

So we need to show that

$$\begin{aligned} \mu_1 \left( \left( \bigcup_{i=1}^n B_i \right) A \right) &= \mu_2 \left( \left( \bigcup_{i=1}^n B_i \right) A \right) \\ \mu_1 \left( \left( \bigcup_{i=1}^n B_i \right) A \right) &= \mu_1 \left( \bigcup_{i=1}^n (B_i A) \right) \\ &= \sum_{1 \leq i \leq n} \mu_1(B_i A) - \sum_{1 \leq i < j \leq n} \mu_1(B_i B_j A) + \cdots + (-1)^{n+1} \mu_1(B_1 \dots B_n A) \\ &= \mu_2 \left( \left( \bigcup_{i=1}^n B_i \right) A \right) \end{aligned}$$

by the continuity shown below,

$$\begin{aligned} \mu_1 \left( \left( \bigcup_{i=1}^n B_i \right) A \right) &\uparrow \mu_1 \left( \left( \bigcup_n B_n \right) A \right) \\ \mu_2 \left( \left( \bigcup_{i=1}^n B_i \right) A \right) &\uparrow \mu_2 \left( \left( \bigcup_n B_n \right) A \right) \end{aligned}$$

So,  $\mu_1(\bigcup_n B_n A) = \mu_2(\bigcup_n B_n A)$ , and since the union of all of the  $B_n$  is  $\Omega$  we have,

$$\mu_1(A) = \mu_2(A)$$

■

This means, that if we can extend  $\mu$  from a field,  $\mathcal{F}_0$  to a  $\sigma$ -field,  $\sigma(\mathcal{F}_0)$  and we know that  $\mu$  is  $\sigma$ -finite on  $\mathcal{F}_0$ , then the extension is unique.

**Theorem 1.5.5** Suppose  $\mu_1$  and  $\mu_2$  are finite measures on  $\sigma(\mathcal{P})$  where  $\mathcal{P}$  is a  $\pi$ -system and  $\Omega$  is a countable union of  $\mathcal{P}$ -sets. then if  $\mu_1, \mu_2$  agree on  $\mathcal{P}$  then they will agree on  $\sigma(\mathcal{P})$ .

*Proof.* Be assumption, there exists  $B_1, B_2, \dots \in \mathcal{P}$  such that  $\Omega = \bigcup_n B_n$ . Because  $\mu_1(B_n) \leq \mu_1(\Omega) < \infty \forall n$  and  $\mu_2(B_n) \leq \mu_2(\Omega) < \infty \forall n$  then they are  $\sigma$ -finite on  $\mathcal{P}$ . Then the theorem follows from the previous theorem. ■

## 1.6 Extension of General Measure to $\sigma$ -Field

### Outer Measure

**Definition 1.6.1 — Outer Measure.**  $\Omega$  nonempty set

An outer measure is a function on  $2^\Omega$  such that



1.  $\mu^*(A) \in [0, \infty] \forall A \subseteq \Omega$
2.  $\mu^*(\emptyset) = 0$
3.  $A \subseteq B \Rightarrow \mu^*(A) \leq \mu^*(B)$
4.  $\forall \{A_n\} \subseteq 2^\Omega, \mu^*(\bigcup_n A_n) \leq \sum_n \mu^*(A_n)$

■ **Example 1.5 — 11.1 in Billingsly.**  $\Omega$ -set.

$\mathcal{A}$  is class of subsets of  $\Omega, \emptyset \in \mathcal{A}$

$$\rho : \mathcal{A} \rightarrow [0, \infty], \rho(\emptyset) = 0$$

We say that  $\{A_n\}$  is an  $\mathcal{A}$  covering of  $A \subseteq \Omega$  if  $A \subseteq \bigcup_n A_n, A_n \in \mathcal{A}$ . ■

**Wednesday September 28**

*Proof.* 1. Want  $\mu^*(A) \in [0, \infty]$

$$\mu^*(A) = \inf \left\{ \sum_n \rho(A_n), \{A_n\} \text{ is a } \mathcal{A}\text{-covering} \right\}$$

Here, we know that the probability is between  $[0, \infty]$  so the sum must be too, and thus the entire equation must be.

2. Want  $\mu^*(\emptyset) = 0$

Because,

$$\emptyset \in \mathcal{A}$$

So,

$$\emptyset \subseteq \emptyset \cup \emptyset \cup \emptyset \cup \dots$$

$$\mu^*(\emptyset) \leq \sum_n \rho(\emptyset) = 0$$

3. Want that if  $A \subseteq B$  and if  $\{A_n\}$  is a  $\mathcal{A}$ -covering of  $A$  then the collection of all  $\mathcal{A}$ -coverings of  $B$  is contained within the collection of all  $\mathcal{A}$ -coverings of  $\mathcal{A}$ .

So we can see that the  $\inf\{\text{statement 1}\} \geq \inf\{\text{statement 2}\}$ .

4. Let  $A_n$  be sets in  $2^\Omega$ , we want that  $\mu^*(\bigcup_n A_n) \leq \sum_n \mu^*(A_n)$ .

For each  $A_n$  let  $\{B_{nk}\}_{k=1}^\infty$  be an  $\mathcal{A}$ -covering of  $A_n$  such that

$$\sum_{k=1}^n \rho(B_{nk}) \leq \mu^*(A_n) + \frac{\epsilon}{2^n}$$

Also, we have  $\{B_{nk} : n = 1, 2, \dots, k = 1, 2, \dots\}$  this is an  $\mathcal{A}$ -covering,  $\bigcup_n A_n$ .

$$\begin{aligned}
\mu^*(\bigcup_n A_n) &\leq \sum_{n=1}^{\infty} \sum_{k=1}^{\infty} \rho(B_{nk}) \leq \mu^*(A_n) + \frac{\varepsilon}{2^n} \\
\mu^*(A_n) + \frac{\varepsilon}{2^n} &= \sum_{n=1}^{\infty} \mu^*(A_n) + \varepsilon \sum_{n=1}^{\infty} \frac{1}{2^n} \\
&= \sum_{n=1}^{\infty} \mu^*(A_n) + \varepsilon \\
\text{So } \mu^*(\bigcup_n A_n) &\leq \sum_n \mu^*(A_n) + \varepsilon \Rightarrow \mu^*(\bigcup_n A_n) \leq \sum_n \mu^*(A_n) \quad \forall \varepsilon > 0.
\end{aligned}$$

■

Now let  $\mathcal{M}(\mu^*)$  to be the paving,

$$\{A : \mu^*(E) = \mu^*(EA) + \mu^*(EA^C), \forall E \in 2^{\Omega}\}$$

**Theorem 1.6.1** If  $\mu^*$  is an outer measure, then  $\mathcal{M}(\mu^*)$  is a  $\sigma$ -field and  $\mu^*$  restricted on  $\mathcal{M}(\mu^*)$  is a measure.

*Proof.* The proof is the exact same as Lemma 3 of Section 3, except  $P^*$  is replace by  $\mu^*$ . It turns out that the  $\infty$  value of  $\mu^*$  does not cause any change in the proof once we invoke the three infinity conventions.

Thus, proof omitted.

■

### Extension of (General) Measure from a Field to a $\sigma$ -Field

**Theorem 1.6.2** A measure on a field has an extension to the generated  $\sigma$ -field.

We will use a different proof from section 3 (even though we could use it). We are going to prove a more general result. But first, we need to introduce the notion of Semiring.

**Definition 1.6.2 — Semiring.** A class of subsets of pavings on  $\Omega$  is called a **semiring** if

1.  $\emptyset \in \mathcal{A}$
2.  $A \in \mathcal{S}, B \in \mathcal{A} \Rightarrow AB \in \mathcal{S}$
3. If  $A \subset B$  then,

$$B \setminus A = \bigcup_{k=1}^n C_k$$

where  $C_1, \dots, C_k$  are disjoint intervals.

A semiring is also a  $\pi$ -system.

■ **Example 1.6** Let  $\mathcal{A} = \{(a, b] : a, b \in \mathcal{R}\}$

1.  $\emptyset \in \mathcal{A}$ ?  $(a, a] = \emptyset$
2. See photo.
3. See photo.

■

*Proof.* Suppose  $\mathcal{A}$  is a semiring.

$\mu$  is a set function such that  $\mu : \mathcal{A} \rightarrow [0, \infty)$ .

Assume  $\mu$  is finitely additive and countable subadditive.

Then  $\mu$  extends to a measure on  $\sigma(\mathcal{A})$  ■

More general than the previous theorem,

1.  $\mathcal{A}$  needs not be a field.
2.  $\mu$  need not be a measure.

*Proof.* 1.  $\mu^*(\emptyset) = 0$ .

2. By monotonicity, let  $A, B \in \mathcal{A}, A \subseteq B$ .

So, because  $\mathcal{A}$  is a semiring,

$$\mu(B \setminus A) = \mu\left(\bigcup_{k=1}^n C_k\right)$$

$$\mu(B) - \mu(A) = \sum_{k=1}^n \mu(C_k) \quad \text{finite additivity}$$

$$\mu(B) = \mu(A) + \sum_{k=1}^n \mu(C_k)$$

$$\geq \mu(A)$$

First, let us show that

$$\mathcal{A} \subseteq \mathcal{M}(\mu^*)$$

So we need to show that  $A \in \mathcal{A}$ .

$$\mu^*(E) = \mu^*(EA) + \mu^*(EA^C) \quad \forall E \in 2^\Omega$$

But we know  $\leq$ , so we need to show  $\geq$ .

If  $\mu^*(E) = \infty$  this is certainly true by  $\infty$ -convention. So let's do  $\mu^*(E) < \infty$ .

Fix a  $\varepsilon > 0$ . Let  $A_n$  be a  $\mathcal{A}$ -coving of  $E$  such that

$$\sum \mu(A_n) \leq \mu^*(E) + \varepsilon$$

Let  $A \in \mathcal{A}, A \setminus A_n$ , then we have

$$A = A \cap A_n \cup A \cap A_n^C = \bigcup_{k=1}^{m_n} C_{nk}$$

Note the  $C$ 's are disjoint  $\mathcal{A}$  sets.

So, for all  $A \in \mathcal{A}$  we may write  $A$  as the disjoint union of  $\mathcal{A}$  sets:

$$A = B_n \cup \left( \bigcup_{k=1}^{m_n} C_{nk} \right)$$

$$\mu^*(EA) = \mu^* \left( E \left( B_n \cup \bigcup_{k=1}^{m_n} C_{nk} \right) \right)$$

Want that  $\mu^*(EA) + \mu^*(EA^C) \leq \mu^*(E)$

$$= \mu^* \left( E B_n \cup \left( E \bigcup_{k=1}^{m_n} C_{nk} \right) \right)$$

■

Monday October 3

**Corollary 1.6.3 — To Theorem 11.1 & 11.3.** Suppose that  $\mathcal{A}$  is a semiring and that we have  $\mu$  such that  $\mathcal{A} \rightarrow [0, \infty]$  is set function such that:

1.  $\mu(\emptyset) = 0$
  2.  $\mu$  is finitely additive on  $\mathcal{A}$
  3.  $\mu$  is countably subadditive on  $\mathcal{A}$
  4.  $\mu$  is  $\sigma$ -finite on  $\mathcal{A}$
- Then,  $\mu$  is a unique extension on  $\sigma(\mathcal{A})$ .

■ **Example 1.7** Let  $\mathcal{A}$  be the collection of all in  $\mathbb{R}$ , that is,

$$\mathcal{A} = \{(a, b] : a, b, \in \mathbb{R}\}$$

Let  $\lambda_1 : \mathcal{A} \rightarrow \mathbb{R}, (a, b] = b - a$ .

By Theorem 1.3,  $\lambda_1$  is finitely additive, and indeed also countably subadditive. So it can be extended to  $\sigma(\mathcal{A}) = \mathcal{R}^1$ . But also we have that  $\lambda_1$  is  $\sigma$ -finite on  $\mathcal{A}$ .

$$\Omega = \mathbb{R} = \bigcup_{n=1}^{\infty} (-n, n], \lambda_1(-n, n] = 2n$$

Therefore the extension is unique. This is defined to be the Lebesgue measure on  $\mathcal{R}^1$  ■

### Approximation Theorem

Approximate  $\mu(A), A \in \sigma(\mathcal{A})$  by  $\mu(B), B \in \mathcal{A}$ .

**Lemma 1.** If  $\mathcal{A}$  is a semiring, and  $A, A_1, \dots, A_n \in \mathcal{A}$ , then we may write,

$$AA_1^C \dots A_n^C$$

as finite disjoint unions of  $\mathcal{A}$ -sets. That is there exists

$$C_1, \dots, C_m \in \mathcal{A}$$

that are disjoint such that

$$AA_1^C \dots A_n^C = C_1 \cup \dots \cup C_m$$

Think of this as a generalization of (iii) of Semiring.

*Proof.* Proof by Induction.

$n = 1$  Case

Want:  $AA_1^C \dots A_n^C = C_1 \cup \dots \cup C_m$

$$AA_1^C = A \setminus (AA_1)$$

$$(AA_1) \subseteq A$$

By (iii) of Semiring, the above is equal to  $C_1 \cup \dots \cup C_m$ .

Assume statement true for  $n$ .

$n + 1$  Case

Induction hypothesis gives us that

$$(AA_1^C \dots A_n^C)A_{n+1}^C = (C_1 \cup \dots \cup C_m)A_{n+1}^C$$

But when we use the  $n=1$  case we have that the following inner term are finitely disjoint unions of  $\mathcal{A}$ -sets,

$$\bigcup_{k=1}^{\infty} C_k A_{n+1}^C$$

So we have that all together (unioned) is a finite disjoint union of  $\mathcal{A}$ -sets. ■

### Symmetric Difference of Sets A, B

**Notation 1.1.**

$$A \triangle B = AB^C \cup BA^C$$

**Theorem 1.6.4** Suppose  $\mathcal{A}$  is a semiring,  $\mu$  is a measure on  $\sigma(\mathcal{A}) = \mathcal{F}$  and  $\mu$  is  $\sigma$ -finite on  $\mathcal{A}$ .

1. For  $B \in \mathcal{F}$ ,  $\varepsilon > 0$ , there exists a disjoint  $\mathcal{A}$ -sequence  $A_1, A_2, \dots$  such that

$$B \subseteq \bigcup_k A_k, \mu\left(\bigcup_k A_k \setminus B\right) < \varepsilon$$

2. If  $B \in \mathcal{F}$ ,  $\mu(B) < \infty$  then for any  $\varepsilon > 0$ , there exists a finite disjoint  $\mathcal{A}$ -sequence,  $A_1, \dots, A_n$  such that

$$\mu\left(B \triangle \bigcup_{k=1}^n A_k\right) < \varepsilon$$

*Proof.* 1. Let  $\mu^*$  be the outer measure,

$$\mu^*(A) = \inf\left\{\sum_n \mu(A_n) : \{A_n\} \text{ is a } \mathcal{A}\text{-covering}\right\}$$

then  $\mathcal{M}(\mu^*)$  is a  $\sigma$ -field,  $\mathcal{F} \subseteq \mathcal{M}(\mu^*)$ ,  $\mu^*$  is a measure,  $\mathcal{M}(\mu^*)$ ,  $\mu = \mu^*$  on  $\sigma(\mathcal{A}) = \mathcal{F}$ .

$$\mu(A) = \inf\left\{\sum_n \mu(A_n) : \{A_n\} \text{ is a } \mathcal{A}\text{-covering}\right\}$$

Let  $B \in \mathcal{F}$ ,  $\mu(B) < \infty$ . Then there exists an  $\mathcal{A}$ -covering  $\{A_k\}$  of  $B$  such that

$$\sum_n \mu(A_k) \leq \mu(B) + \varepsilon$$

So  $\mu(\bigcup_k A_k) \leq \sum_k \mu(A_k) \leq \mu(B) + \varepsilon$ .

And  $\mu(\cup_k A_k) - \mu(B) \leq \varepsilon$ .

$$\mu(\cup_k A_k \setminus B) \leq \varepsilon$$

Let

$$\begin{aligned} B_1 &= A_1 \\ B_2 &= A_2 A_1^C \\ &\vdots \\ B_k &= A_k A_1^C \dots A_{k-1}^C \\ &\vdots \end{aligned}$$

So by Lemma 1, the  $B_k$  are finite disjoint union of  $\mathcal{A}$ -sets. Also,

$$\bigcup_k A_k = \bigcup_k B_k$$

so we have that there exists  $C_1, C_2, \dots$  that are also disjoint  $\mathcal{A}$ -sets such that

$$\mu(\cup_k C_k \setminus B) \leq \varepsilon$$

Now suppose that  $B \in \mathcal{F}, \mu(B) = \infty$ . Because  $\mu$  is  $\sigma$ -finite on  $\mathcal{A}$  there exists  $C_1, \dots \in \mathcal{A}$  such that  $\mu(C)m \leq \infty$ .

$$\Omega = \bigcup_m C_m$$

So then

$$\begin{aligned} B &= B\Omega \\ &= B(\cup_m C_m) \\ &= \cup_m BC_m \end{aligned}$$

### Wednesday October 5

But  $\mu(BC_m) \leq \mu(C_m) < \infty$  so by the finite case there exists a disjoint  $\mathcal{A}$ -sequence  $\{A_{mk} : k = 1, 2, \dots\}$  such that

$$\bigcup_k A_{mk} \supseteq BC_m$$

$$\mu(\bigcup_k A_{mk} \setminus (BC_m)) < \frac{\varepsilon}{2^k}$$

So now,

$$B = \cup_m BC_m \subseteq \bigcup_m \bigcup_k A_{mk}$$

$$\begin{aligned}
\mu\left(\bigcup_m \bigcup_k A_{mk} \setminus B\right) &= \mu\left(\left(\bigcup_m \bigcup_k A_{mk}\right) \setminus (BC_m)\right) \\
&= \mu\left(\left(\bigcup_m \bigcup_k A_{mk}\right) \cap (BC_m)^C\right) \\
&= \mu\left(\bigcup_m \left(\bigcup_k A_{mk}\right) \cap (BC_m)^C\right) \\
&\leq \sum_m \mu\left(\bigcup_k A_{mk} \setminus (BC_m)\right) \\
&\leq \varepsilon
\end{aligned}$$

Since  $\bigcup_m (\bigcup_k A_{mk})$  is a countable union of  $\mathcal{A}$ -sets, we can write as

$$\bigcup_k D_k$$

and make them disjoint as before,

$$E_1 = D_1$$

$$E_2 = D_2 \setminus D_1^C$$

$$\vdots$$

By Lemma 1, each  $E_m$  is finite disjoint union of  $\mathcal{A}$ -sets.

$$\bigcup_m E_m = \bigcup_m F_m$$

where  $F_m$  are disjoint  $\mathcal{A}$ -sets.

Hence, part (i) is proved.

2. Recall for any  $B \in \mathcal{F}$ ,  $\mu(B) < \infty$  and  $\varepsilon > 0$ , there exists a finite  $\mathcal{A}$ -sequence,  $A_1, \dots, A_n$ , such that

$$\mu\left(B \triangle \bigcup_{k=1}^n A_k\right) < \varepsilon$$

$$\begin{aligned}
\mu\left[\left(\bigcup_{k=1}^n A_k\right) \triangle B\right] &= \mu\left(\left(\bigcup_{k=1}^n A_k\right)^C \cap B \cup \left(\bigcup_{k=1}^n A_k\right) \cap B^C\right) \\
&\leq \mu\left(\left(\bigcup_{k=1}^n A_k\right)^C \cap B\right) + \mu\left(\left(\bigcup_{k=1}^n A_k\right) \cap B^C\right)
\end{aligned}$$

By (i) there exists disjoint  $\mathcal{A}$ -sets  $\{A_n\}$  such that

$$\mu\left(\bigcup_n A_n \setminus B\right) < \varepsilon$$

Let  $A = \bigcup_n A_n$ .

Then,

$$A \setminus \bigcup_{k=1}^n A_k \downarrow \emptyset$$

Need,  $\mu(A \setminus A_1) < \infty$  and  $\mu(A) < \infty$ .

By continuity from above, we have  $\mu(A \setminus \bigcup_{k=1}^n A_k) \downarrow 0$ .

So, for sufficiently large  $n$  we have,

$$\mu(A \setminus \bigcup_{k=1}^n A_k) < \varepsilon$$

Now first take a look at  $\mu((\bigcup_{k=1}^n A_k)B^C)$ .

$$\begin{aligned} \mu((\bigcup_{k=1}^n A_k)B^C) &\leq \mu(AB^C) \\ &= \mu(A \setminus B) < \varepsilon \end{aligned}$$

$$\begin{aligned} \mu((\bigcup_{k=1}^n A_k)^C A) &\leq \\ &= \mu(A \setminus \bigcup_{k=1}^n A_k) < \varepsilon \end{aligned}$$

So we have that  $\mu(\bigcup_{k=1}^n A_k \triangle B) < 2\varepsilon$

■

The next lemma will be used in the next section, this is an extension of Theorem 1.3 in the textbook.

**Lemma 2.** Suppose  $\mathcal{A}$  is a semiring,  $A_1, \dots, A_n, A$  be  $\mathcal{A}$ -sets, and  $\mu$  is a non-negative, finitely additive set function on  $\mathcal{A}$ . Then

1. If  $\bigcup_{k=1}^n A_k \subset A$  and  $A_k$  are disjoint, then

$$\sum_{k=1}^n \mu(A_k) \leq \mu(A)$$

2. If  $A \subset \bigcup_{k=1}^n A_k$  ( $A_k$  don't have to be disjoint) then

$$\mu(A) \leq \sum_{k=1}^n \mu(A_k)$$

*Proof.* 1. By Lemma 1,

$$\begin{aligned} A \setminus \bigcup_{k=1}^n A_k &= A(\bigcup_{k=1}^n A_k)^C \\ &= AA_1^C \dots A_n^C \\ &= C_1 \cup \dots \cup C_n \end{aligned}$$

where the  $C_k$  are disjoint  $\mathcal{A}$ -sets.



So,

$$A = \left( \left( \bigcup_{k=1}^n A_k \right) \cup \left( \bigcup_{l=1}^m C_l \right) \right)$$

And thus we now have  $A_1, \dots, A_n, C_1, \dots, C_m$  disjoint  $\mathcal{A}$ -sets. By finite additivity of  $\mu$ ,

$$\mu(A) = \mu(A_1) + \dots + \mu(A_n) + \mu(C_1) + \dots + \mu(C_m) \geq \mu(A_1) + \dots + \mu(A_n)$$

2. Want  $A \subseteq \bigcup_{k=1}^n A_k \Rightarrow \mu(A) \leq \sum_{k=1}^n \mu(A_k)$ .

Let

$$B_1 = A_1$$

$$B_2 = A_2 A_1^C$$

$\vdots$

$$B_n = A_n A_1^C \dots A_{n-1}^C$$

Let

$$C_1 = AB_1$$

$\vdots$

$$C_n = AB_n$$

Then,  $C_1, \dots, C_n$  are disjoint.

$$A = \bigcup_{i=1}^n C_i$$

By Lemma 1,

$$C_i = AA_i A_1^C \dots A_{i-1}^C$$

which are finite disjoint union of  $\mathcal{A}$ -sets ( $AA_i$  are  $\pi$ -system  $\mathcal{A}$ -sets).

$$C_i = \bigcup_{j=1}^{m_i} D_{ij}$$

In the meantime,

$$C_i = AA_i A_1^C \dots A_{i-1}^C \subseteq A_i$$

Therefore,

$$\bigcup_{j=1}^{m_i} D_{ij} \subseteq A_i$$

By part (i)

$$\sum_{j=1}^{m_i} \mu(D_{ij}) \leq \mu(A_i)$$

$$\sum_{i=1}^n \sum_{j=1}^{m_i} \mu(D_{ij}) \leq \sum_{i=1}^n \mu(A_i)$$

But the  $D_{ij}$  are disjoint  $\mathcal{A}$ -sets.

$$\bigcup_{i=1}^m \bigcup_{j=1}^{m_i} D_{ij} = A$$

By finite additivity of  $\mu$  on  $\mathcal{A}$ ,

$$\mu(A) = \sum_{j=1}^{m_i} \mu(D_{ij})$$

Hence,

$$\mu(A) \leq \sum_{i=1}^n \mu(A_i)$$

■

## 1.7 Measure in Euclidean

### Extend Measure to $\mathcal{R}^k$

What we have done is to extend  $\mu$  from intervals to  $\mathcal{R}'$ .

**Friday October 5**

### Characterizing Measures in $\mathbb{R}$

The only measure we know so far is the Lebesgue measure. Let  $\mu$  be any measure that has finite value on bounded sets,  $\mu(A) < \infty$ .

A is ?.

Bounded set:  $\sup_{x \in A} ||x|| < \infty$

Then we can define

This is obviously nondecreasing.

For  $0 < a < b, a < 0 < b, a < b < 0$  we can show that  $F(b) \geq F(a)$ .

Also it is right continuous for  $x \geq 0$ . So if  $x_n \downarrow x$ ,

$$(0, x_n] \downarrow (0, x]$$

implies

$$\mu(0, x_n] \downarrow \mu(0, x]$$

and

$$F(x_n) \rightarrow F(x)$$

for  $x < 0$ , where  $x_n \downarrow x$ ,

$$(x_n, 0] \downarrow (x, 0]$$

implies

$$\mu(x_n, 0] \downarrow \mu(x, 0]$$

So,

$$-\mu(x_n, 0] \rightarrow -\mu(x, 0]$$

and

$$F(x_n) \rightarrow F(x)$$

So  $F$  is right continuous.

**Theorem 1.7.1 — Theorem 12.4.** If  $F$  is

1. nondecreasing
  2. right continuous
- then there exists a unique measure,  $\mu$  on  $\mathcal{R}$  such that

$$\mu(a, b] = F(b) - F(a) \quad \forall a, b \in \mathbb{R}$$

### Characterizing Measures in $\mathbb{R}^k$

Let  $\mathcal{R}^k$  be the Borel  $\sigma$ -field on  $\mathbb{R}^k$ . Note that  $\sigma(\mathcal{A}) = \mathcal{R}'$ .

So  $\mathcal{R}^k = \sigma\{(a_1, b_1]x \dots x(a_k, b_k] : a_1, b_1, \dots, a_k, b_k \in \mathbb{R}\}$ .

For each  $x \in \mathcal{R}^k$ , let

$$S_x = \text{"Southwest"} = (-\infty, x_1]x \dots x(-\infty, x_k]$$

Then we can show that

$$\mathcal{R}^k = \sigma\{S_x : x \in \mathbb{R}^k\}_{rs}$$

To see this, for any bounded rectangle  $A = (a_1, b_1]x \dots x(a_k, b_k]$  let  $V_A$  be the collection of all vertices of  $A$ , that is  $V_A = \{a_1, b_1\}x \dots x\{a_k, b_k\}$ . So we have that  $\#(V_A) = 2^k$ .

Now we can express

$$(a_1, b_1]x \dots x(a_k, b_k] = S_{b_1, \dots, b_k} \setminus \bigcup_{V_A \setminus \{b_1, \dots, b_k\}} S_{x_1, \dots, x_k}$$

So,  $\sigma\{\mathcal{A}\} = \mathcal{R}^k \subseteq \sigma\{S_x : x \in \mathbb{R}^k\}$ .

In the other direction, we have any

$$S_x = \bigcup_{n=1}^{\infty} (x_i - n, x_i]$$

Then  $\sigma\{S_x : x \in \mathbb{R}^k\} \subseteq \mathcal{R}^k$ .

So,

$$\sigma\{S_x : x \in \mathbb{R}^k\} = \mathcal{R}^k$$

**Notation 1.2 (Signum).**  $\text{sgn}_A(x)$  is a signum of a vertex in a rectangle. Signum means "sign" in Latin.

**Monday October 10**

Let  $F : \mathbb{R}^k \rightarrow \mathbb{R}$ . Also  $A \in \mathcal{J}^k$ ,  $\mathcal{J}^k$  is the collection of all bounded rectangles. That is,  $\{(a_1, b_1]x \dots x(a_k, b_k] : a_1, b_1, \dots, a_k, b_k \in \mathbb{R}\}$ .

Let

$$\triangle_A F = \sum_{x \in V_A} \text{sgn}_A(x) F(x)$$

where  $V_A$  is the collection of all vertices of  $A$ .

For illustration, assume  $\mu$  is a finite measure. We won't need this assumption.

Let  $F(x) = \mu(S_x)$ , where  $S_x$  is the southwest of  $x$ .

$$S_x = (-\infty, x_1]x \dots x(-\infty, x_k]$$

Then we show for any  $A \in \mathcal{J}^k$  we have

$$\mu(A) = \triangle_A F$$

To see this,

$$A = \bigcup_{x \in V_A \setminus \{b_1, \dots, b_k\}} S_{(x_1, \dots, x_k)}$$

For  $k = 2$ ,

$$\mu\left(\bigcup_{x \in V_A \setminus \{b_1, \dots, b_k\}} S_{(x_1, \dots, x_k)}\right) = \mu(S_{b_1 \dots b_k}) - \bigcup_{x \in V_A \setminus \{b_1, \dots, b_k\}} S_k$$

There are  $2^k - 1$  number of values in this range.

$$\bigcup_{i=1}^m B_i \quad m = 2^k - 1$$

$$\mu\left(\bigcup_{i=1}^m B_i\right) = \sum_{i=1}^m \mu(B_i) - \sum_{i < j} \mu(B_i B_j) + \dots + (-1)^{m+1} \mu(B_1 \dots B_m)$$

For  $k = 2$ ,

$$\sum_{i=1}^m \mu(B_i) = \mu(S_{a_1 a_2}) + \mu(S_{a_1 b_2}) + \mu(S_{b_1 a_2})$$

$$\sum_{i < j} \mu(B_i B_j) = \mu(B_1 B_2) + \mu(B_2 B_3) + \mu(B_1 B_3) = 3\mu(B_1)$$

$$\mu(B_1 B_2 B_3) = \mu(B_1)$$

All together,

$$\begin{aligned}
\mu\left(\bigcup_{i=1}^m B_i\right) &= \mu(S_{B_1 B_2}) - \mu(B_1) - \mu(B_2) - \mu(B_3) + 3\mu(B_1) - \mu(B_1) \\
&= \mu(S_{B_1 B_2}) + \mu(B_1) - \mu(B_2) - \mu(B_3) \\
&= \sum_{x \in V_A} \text{sgn}_A(x) F(x) \\
&= \triangle_A F
\end{aligned}$$

By induction, we may show that  $\mu(A) = \triangle_A F$ .

So,

$$\triangle_A F \geq 0 \quad \forall A \in \Omega^k$$

Also, if  $x^{(n)} \downarrow x$ , in the sense that

$$x_1^{(n)} \downarrow x, \dots, x_k^{(n)} \downarrow x$$

then,  $S_{x^{(n)}} \downarrow S_x$ .

So  $\mu(S_{x^{(n)}}) \downarrow \mu(S_x)$ .

$$F(x^{(n)}) \rightarrow F(x)$$

So,  $F(x)$  is continuous from above in the sense that  $x^{(n)} \downarrow x$ .

$$F(x^{(n)}) \downarrow F(x)$$

This shows that  $\mu \Rightarrow F$  such that  $\triangle_A F \geq 0$  continuous from above.

In fact, such an  $F$ , also uniquely determines a measure.

**Theorem 1.7.2** Suppose  $F : \mathbb{R}^k \rightarrow \mathbb{R}$  is continuous from above. That is,

$$\lim_{x^{(n)} \downarrow x} F(x^{(n)}) = F(x)$$

Also suppose that for all  $A \in \mathcal{J}^k$ ,

$$\triangle_A F \geq 0$$

corresponding to right continuous and nondecreasing in  $\mathbb{R}^k$ .

Then, there exists a unique measure  $\mu$  on  $(\mathbb{R}^k, \mathcal{B}^k)$  such that for all  $A \in \mathcal{J}^k$ ,

$$\mu(A) = \triangle_A F$$

The most important special case of this theorem is the case

$$F(x) = x_1 \dots x_k$$

$$\sum_{x \in V_A} \text{sgn}_A(x) F(x) = (b_1 - a_1) \dots (b_k - a_k)$$

So the  $\mu$  corresponding to this  $F$  is the Lebesgue measure.

This characterizes all measures in  $\mathbb{R}^k$ .

*Proof.* Note that  $\mu$  is defined on  $\mathcal{J}^k$ . So we need to show  $\mu$  can be uniquely extended to  $\sigma(\mathcal{J}^k) = \mathcal{R}^k$ .

*Uniqueness*

Want  $\mu$   $\sigma$ -finite on  $\mathcal{J}^k$ .

$$A_n = (-n, n] \times \dots \times (-n, n]$$

$$\cup_n A_n = \mathbb{R}^k$$

$$\mu(A_n) = \sum_{x \in V_{A_n}} s_{q_{A_n}}(x) F(x)$$

But  $F : \mathbb{R}^k \rightarrow \mathbb{R}$ ,  $F(x) < \infty \forall x$ , and  $\mu(A_n) < \infty$ .

So  $\mu$  is  $\sigma$ -finite on  $\mathcal{J}^k$ .

Still need existence of extension finitely additive, countably subadditive.

**Step 1** Finitely additive.

*Step 1 (a)*

Finitely additive on regular partition of  $A \in \mathcal{J}^k$ .

What's a regular partition? Irregular partition? Think disjoint vs overlapping.

It's easy to turn an irregular partition into a regular one.

**Wednesday October 12**

More explicitly,

$$A = I_1 \times \dots \times I_k$$

where,  $I_i = (a_i, b_i]$  for  $i = 1, \dots, k$ .

For each  $i$  let,

$$J_{i1}, \dots, J_{ik}$$

be a partition of  $I_i$  into subintervals.

For each  $(j_1, \dots, j_k) \in \{1, \dots, n_1\} \times \dots \times \{1, \dots, n_k\}$

Write

$$B_{j_1, \dots, j_k} = J_{1j_1} x \dots x J_{kj_k}$$

So we have

$$\mathcal{B} = \{B_{j_1 \dots j_k} : j_i \in \{1, \dots, n_i\} \forall i = 1, \dots, k\}$$

$$\#\mathcal{B} = n_1 \dots n_k$$

So  $\mathcal{B}$  is called a **regular decomposition of A**.

Obviously  $A = \bigcup_{B \in \mathcal{B}} B$  and  $\{B : B \in \mathcal{B}\}$  are disjoint and  $B \in \mathcal{J}^k$ .

Overall, Step 1 (a) claims that  $\mu(A) = \sum_{B \in \mathcal{B}} \mu(B)$ .

Recall that  $\mu(B) = \triangle_B F = \sum_{x \in V_B} \text{sgn}_B(x) F(x)$ .

Here we have,

$$\sum_{B \in \mathcal{B}} \mu(B) = \sum_{B \in \mathcal{B}} \sum_{x \in V_B} \text{sgn}_B(x) F(x)$$

We may change the order of summations,

$$\sum_{x \in V} \sum_{B \in W_x} \text{sgn}_B(x) F(x)$$

where  $W_x = \{B \in \mathcal{B} : x \in V_B\}$  and  $V = \bigcup_{B \in \mathcal{B}} V_B$ .

Now be separating into values of x in and not in  $V_A$ , we have,

$$\sum_{x \in V_A} \sum_{B \in W_x} \text{sgn}_B(x) F(x) + \sum_{x \notin V_A} \sum_{B \in W_x} \text{sgn}_B(x) F(x)$$

So the idea is that when  $x \notin V_1$  then  $\sum_{B \in W_x} \text{sgn}_B(x) = 0$  which means the second term above is also zero. But, if  $x \in V_A$ , then  $W_x$  is singleton and  $B \in W_x$  has same sign as A.

So ultimately,

$$\sum_{B \in \mathcal{B}} \mu(B) = \sum_{x \in V_A} \text{sgn}_A(x) F(x) = \mu(A)$$

*Step 1 (b)*

Now, consider, general situation. If we let  $A \in \mathcal{J}^k$  and suppose that  $A = \bigcup_{u=1}^n A_u, A_u \in \mathcal{J}^k$ . Because  $A_n \in \mathcal{J}^k$ , and  $A_u = I_{1u} x \dots x I_{ku} \in \mathcal{J}^1$  we have,

$$A = \bigcup_{u=1}^n (I_{1u} x \dots x I_{ku})$$

Meanwhile,  $A \in \mathcal{J}^k$ , so  $A = I_1 x \dots x I_k$ .

Claim:

$$I_1 x \dots x I_k = \bigcup_{u=1}^n (I_{1u} x \dots x I_{ku}) = \bigcup_{u=1}^n (I_{1u}) x \dots x \bigcup_{u=1}^n I_{ku}$$

But if  $(a_1, \dots, a)k \in \bigcup_{u=1}^n (I_{1u}x \dots x I_{ku})$ , then

$$(a_1, \dots, a)k \in I_{1u}x \dots x I_{ku}$$

for some  $u$ .

Then we have  $a_i \in I_{iu} \subseteq \bigcup_{u=1}^n I_{iu}$  which leads to

$$(a_1, \dots, a)k \subseteq \left( \bigcup_{u=1}^n I_{1u} \right) x \dots x \left( \bigcup_{u=1}^n I_{ku} \right)$$

So,

$$\bigcup_{u=1}^n (I_{1u}x \dots x I_{ku}) \subseteq \left( \bigcup_{u=1}^n I_{1u} \right) x \dots x \left( \bigcup_{u=1}^n I_{ku} \right)$$

On the other hand,

$$I_{iu} \subset I_u$$

$$\left( \bigcup_{u=1}^n I_{iu} \right) \subseteq I_u$$

$$\left( \bigcup_{u=1}^n I_{1u} \right) x \dots x \left( \bigcup_{u=1}^n I_{ku} \right) \subseteq I_1 x \dots x I_k$$

Thus, claim is proved. Hence,

$$\left( \bigcup_{u=1}^n I_{1u} \right) x \dots x \left( \bigcup_{u=1}^n I_{ku} \right) = I_1 x \dots x I_k$$

The union,  $(\bigcup_{u=1}^n I_{iu})$ , needs not be disjoint, but we may use all the endpoints of  $I_{iu}$  to make disjoint partitions.

So then

$$A = \left( \bigcup_{v=1}^{m_1} \tilde{I}_{1v} \right) x \dots x \left( \bigcup_{v=1}^{m_k} \tilde{I}_{kv} \right)$$

which by Step 1 (a) is a regular partition.

$$\mu(A) = \sum_{v=1}^{m_1} \dots \sum_{v=1}^{m_k} \mu(\tilde{I}_{1v_1} x \dots x \tilde{I}_{kv_k})$$

Let  $\tilde{\mathcal{B}} = \{\tilde{I}_{1v_1} x \dots x \tilde{I}_{kv_k} : v_i = 1, \dots, m_i \forall i = 1, \dots, k\}$ .

So,

$$\mu(A) = \sum_{\tilde{B} \in \tilde{\mathcal{B}}} \mu(\tilde{B}) = \sum_{u=1}^n \sum_{\tilde{B} \subseteq A_u} \mu(\tilde{B})$$

Thus finite additivity done.

**Step 2** Countably subadditive.



**Friday October 14**

First we want to show finite subadditive.

That is if  $A_1, \dots, A_n, A \in \mathcal{I}^k$  and  $A \subseteq \bigcup_{i=1}^n A_i$  then,

$$\mu(A) \leq \sum_{i=1}^n \mu(A_i)$$

But this is implied by Lemma 2 at end of section 11.

We need to extend this to countably subadditivity.

Recall compact set, in  $\mathbb{R}^N$  (definition applies to any topological space). A set is **compact** if any open covering has a finite subcovering. That is, if  $A \subseteq \bigcup_{G \in \mathcal{G}} G$  where  $G$  is open, then there exists a subcollection

$$\{G_1, \dots, G_n\} \subset \mathcal{G}$$

so that  $A \subset \bigcup_{i=1}^n G_i$  and we may call  $A$  compact. In the appendix of Billingsly, there is the Heine - Borel Theorem: A bounded and closed set in  $\mathbb{R}^k$  is compact.

Now we want to show that if  $A_1, \dots, A_n \in \mathcal{I}^k$  and  $A \subseteq \bigcup_n A_n$  then

$$\mu(A) \leq \sum_n \mu(A_n)$$

Because  $A \in \mathcal{I}^k$ , then  $A = (a_1, b_1] \times \dots \times (a_k, b_k]$ . Let

$$B(\delta) = (a_1 + \delta, b_1] \times \dots \times (a_k + \delta, b_k]$$

Then

$$\mu(B(\delta)) = \sum_{x \in V_{B(\delta)}} \text{sgn}_{B(\delta)}(x) F(x)$$


Note that  $x(\delta) \in V_{B(\delta)}$ .

So  $x(\delta)$  may be written as

$$x(\delta) = x + \begin{pmatrix} \delta \\ 0 \\ \delta \\ \vdots \\ \delta \end{pmatrix} = x + \begin{pmatrix} 1 \\ 0 \\ 1 \\ \vdots \\ 1 \end{pmatrix} \begin{pmatrix} \delta \\ \delta \\ \delta \\ \vdots \\ \delta \end{pmatrix}$$

■





# Part Two

2	Integration with Respect to a Measure	69
3	Random Variable .....	71
4	Convergence in Probability/Limit Theorem	73
5	Radon-Nikodym Derivative Theorem .	75
6	Special Topics .....	77
	Index .....	79





## 2. Integration with Respect to a Measure





### 3. Random Variable







## 4. Convergence in Probability/Limit Theorem





## 5. Radon-Nikodym Derivative Theorem





## 6. Special Topics





## Index

- Exten. Prob Measure to  $\sigma$ -field, 14
- Extension of General Measure to  $\sigma$ -Field, 48
- General Measure on a Field, 37
- Measure in Euclidean, 58
- Probabilities Concerning Sequences of Events,  
28
- Probability on a Field, 7