



Probability Theroy based on Measure Theory

STAT 517

Dr. John Smith



Copyright © 2013 John Smith

PUBLISHED BY PUBLISHER

BOOK-WEBSITE.COM

Licensed under the Creative Commons Attribution-NonCommercial 3.0 Unported License (the “License”). You may not use this file except in compliance with the License. You may obtain a copy of the License at <http://creativecommons.org/licenses/by-nc/3.0>. Unless required by applicable law or agreed to in writing, software distributed under the License is distributed on an “AS IS” BASIS, WITHOUT WARRANTIES OR CONDITIONS OF ANY KIND, either express or implied. See the License for the specific language governing permissions and limitations under the License.

First printing, March 2013

Contents

I	Part One	
1	Probability Measure	7
1.1	Overview	7
1.2	Probability on a Field	7
1.3	Extention of Probability Measure to a σ -field	14
2	General Measure	27
3	Integration with Respect to a Measure	29
4	Random Variable	31
5	Convergence in Probability/Limit Theorem	33
6	Radon-Nikodym Derivative Theorem	35
7	Special Topics	37
	Index	39

Part One

1	Probability Measure	7
1.1	Overview	
1.2	Probability on a Field	
1.3	Extention of Probability Measure to a σ -field	
2	General Measure	27
3	Integration with Respect to a Measure	29
4	Random Variable	31
5	Convergence in Probability/Limit Theorem	33
6	Radon-Nikodym Derivative Theorem .	35
7	Special Topics	37
	Index	39

1. Probability Measure

1.1 Overview

1.2 Probability on a Field

■ **Definition 1.2.1** — Ω . Non empty set.

■ **Definition 1.2.2** — **Paving**. A collection of a subset of Ω is a paving.

■ **Definition 1.2.3** — **Field**. A field \mathcal{F} is a paving satisfying

- (i) $\Omega \in \mathcal{F}$
- (ii) $A \in \mathcal{F} \Rightarrow A^C \in \mathcal{F}$
- (iii) $A, B \in \mathcal{F} \Rightarrow A \cup B \in \mathcal{F}$

Derived Properties about a Field

- $\emptyset \in \mathcal{F}$ (by (i) and (ii):

$$\begin{aligned}\Omega \in \mathcal{F} &\Rightarrow \Omega^C \in \mathcal{F} \\ &\Rightarrow \emptyset \in \mathcal{F})\end{aligned}$$

- (i) can be replaced by " \mathcal{R} is nonempty" because,
Let $A \in \mathcal{F}$,

$$\begin{aligned}&\Rightarrow A^C \in \mathcal{F} \\ &\Rightarrow A^C \cup A \in \mathcal{F} \\ &\Rightarrow \Omega \in \mathcal{F}\end{aligned}$$

- $A \in \mathcal{F}, B \in \mathcal{F} \Rightarrow A \cap B \in \mathcal{F}$ because,

$$\begin{aligned}(A \cap B)^C &= A^C \cup B^C \text{ (DeMorgan's Law)} \\ A \cap B &= (A^C \cup B^C)^C\end{aligned}$$

- $A_1, \dots, A_m \in \mathcal{F} \Rightarrow A_1 \cup \dots \cup A_m \in \mathcal{F}$ (mathematical induction)
- $A_1, \dots, A_m \in \mathcal{F} \Rightarrow A_1 \cap \dots \cap A_m \in \mathcal{F}$

Definition 1.2.4 — σ -Field. Similar to the definition of a field except for (iii). A paving satisfying

- (i) $\Omega \in \mathcal{F}$
- (ii) $A \in \mathcal{F} \Rightarrow A^c \in \mathcal{F}$
- (iii) $A_1 \in \mathcal{F}, \dots, A_m \in \mathcal{F}$
 $\bigcup_{k=1}^m A_k \in \mathcal{F}$ (finite additivity)

If we replace (iii) from before by (iii') here:

For $A_1 \in \mathcal{F}, \dots, A_m \in \mathcal{F}$

$$\bigcup_{k=1}^{\infty} A_k \in \mathcal{F}$$

then \mathcal{F} is called a **σ -field**.

Derived Facts

- Again, (i) can be replaced by \mathcal{F} not empty, (iii) can be replaced $A_1 \in \mathcal{F}, \dots, A_m \in \mathcal{F}$

■ **Example 1.1** $\Omega = (0, 1]$ (from now on all intervals are left open, right closed)

R Recall that σ -fields are generated by fields. Fancy scripts denote a σ -field. Fancy scripts with a zero subscript denote a field.

\mathcal{B}_0 is the collection of all finite union of disjoint intervals. Asside: something that we can consider easily, but computers cannot. (e.g. think of all stars, etc)




Figure 1.1: Finite union of three disjoint intervals.

Then \mathcal{B}_0 is a field.

- (i) $(0, 1] \in \mathcal{B}_0$
- (ii) $A \in \mathcal{B}_0 \Rightarrow A^c \in \mathcal{B}_0$
- (iii) $A \in \mathcal{B}_0, B \in \mathcal{B}_0 \Rightarrow A \cup B \in \mathcal{B}_0$

Wednesday August 24

\mathcal{B}_0 = collection of finite unions of disjoint subintervals of $(0, 1]$. Is a field.





Figure 1.2: A and complement of A.

Figure 1.3: Union of A and B is still in \mathcal{B}_0

Definition 1.2.5 — Power Set. A σ -field is generated by a paving of power set. Let Ω be a set. The collection of all subsets of Ω is the power set written as 2^Ω .

- R** Where does this notation come from?
Consider this case where Ω is finite

$$\Omega = \{\omega_1, \dots, \omega_n\}$$

Total number of subsets of Ω .

\emptyset , 1 element sets, 2-element sets, ..., n-element sets.

$$() + () + \dots + = (1 + 1)^n$$

$\#(\mathcal{F}) = 2^{\#\Omega}$, so it seems reasonable to denote $\mathcal{F} = 2^\Omega$.

It is also easy to show that 2^Ω is a σ -field. (The largest, even. The smallest: $\{\emptyset, \Omega\}$ which is also a σ -field.)

$$\{\emptyset, \Omega\} \subseteq \sigma\text{-field} \subseteq 2^\Omega$$

It turns out we can extend notion of length from \mathcal{B}_0 to σ -field generated by \mathcal{B}_0 .

Now, let \mathcal{A} be a nonempty paving of Ω . We define

$$\sigma(\mathcal{A}) = \cap \{ \mathcal{B} \subseteq 2^\Omega : \mathcal{B} \text{ is a } \sigma\text{-field, } \mathcal{A} \subseteq \mathcal{B} \}$$

OR rather, the *intersection* of all σ -fields that contains \mathcal{A} .

Let

$$\mathbb{F}(\mathcal{A}) = \{ \mathcal{B} \subseteq 2^\Omega : \mathcal{B} \text{ is a } \sigma\text{-field, } \mathcal{B} \supseteq \mathcal{A} \}$$

Then,


$$\sigma(\mathcal{A}) = \cap \mathcal{B}$$

$$\mathcal{B} \in \mathbb{F}(\mathcal{A})$$

Derived Facts

$\mathbb{F}(\mathcal{A})$ is nonempty. For example, 2^Ω is a σ -field and $2^\Omega \supseteq \mathcal{A}$.

$\cap \mathcal{B}$ is a σ -field. ($\mathcal{B} \in \mathbb{F}(\mathcal{A})$)

 Get notes about notation/levels.

Proof. We will prove that indeed $\sigma(\mathcal{A})$ is a σ -field. Recall that we have three conditions above for σ -field.

(i)

$$\Omega \in \sigma(\mathcal{A})$$

$$\Omega \in \cap_{\mathcal{B} \in \mathbb{F}(\mathcal{A})} \mathcal{B}$$

Because: \mathcal{B} is σ -field, $\Omega \in \mathcal{B}, \forall \mathcal{B} \in \mathbb{F}(\mathcal{A})$.

(ii)

$$(iii) \quad A_1, \dots, \in \cap_{\mathcal{B} \in \mathbb{F}(\mathcal{A})} \mathcal{B}, \forall \mathcal{B} \in \mathbb{F}(\mathcal{A})$$

$$\Rightarrow \cup_{n=1}^{\infty} A_n \in \mathcal{B}, \forall \mathcal{B} \in \mathbb{F}(\mathcal{A})$$

So, $\sigma(\mathcal{A})$ is a σ -field, we call it the σ -field, generated by \mathcal{B}_o . We know how to assign length to members of \mathcal{B}_o , we now show the assignment can be extended to $\sigma(\mathcal{B}_o)$ ■

■ **Example 1.2** Let \mathcal{I} be the collection of *all* subintervals of $(0,1]$.

Note that \mathcal{I} is a smaller collection than \mathcal{B}_0 since \mathcal{B}_0 can have numerous different combinations of the sets.

Let

$$\mathcal{B} = \sigma(\mathcal{I})$$

This is a Borel- σ -field. (a member of \mathcal{B} in Borel set.)

It turns out

$$\sigma(\mathcal{I}) = \sigma(\mathcal{B}_o)$$

This is because $\sigma(\mathcal{I})$ is a σ -field.

So,

$$\begin{aligned}\sigma(\mathcal{I}) &\supseteq \mathcal{B}_o \\ \sigma(\mathcal{I}) &\supseteq \sigma(\mathcal{B}_o)\end{aligned}$$

Also,

$$\begin{aligned}\mathcal{I} &\subseteq \mathcal{B}_o \\ \sigma(\mathcal{I}) &\subseteq \sigma(\mathcal{B}_o)\end{aligned}$$

Thus,

$$\sigma(\mathcal{I}) = \sigma(\mathcal{B}_o)$$

■

Definition 1.2.6 — Probability Measure. Probability measures on field. Suppose \mathcal{F} is a field on a nonempty set Ω . A probability measure is a function $P : \mathcal{F} \rightarrow \mathbb{R}$.

- (i) $0 \leq P(A) \leq 1, \forall A \in \mathcal{F}$
- (ii) $P(\emptyset) = 0, P(\Omega) = 1$
- (iii) If A_1, \dots are disjoint members of \mathcal{F} and $\cup A_n \in \mathcal{F}$ then we have countable additivity:

$$P(\cup A_n) = \sum_{n=1}^{\infty} P(A_n)$$

R Note that (iii) also implies finite additivity. Prove by adding infinite empty sets on end.

If Ω is nonempty set. And \mathcal{F} is a σ -field on Ω . And P is a probability measure on \mathcal{F} . Then (Ω, \mathcal{F}, P) is called a **probability space**. And (Ω, \mathcal{F}) is called a **measurable space**.

R If $A \subseteq B$, then $P(A) \leq P(B)$. This is because we may write B as

$$B = A \cup (B \setminus A)$$

R

$$P(A) + P(B) = P(A \cup B) + P(A \cap B)$$

Friday August 26

Recall,

Probability measure on a field, \mathcal{F}_0 .

- $P(A) + P(B) = P(A \cup B) + P(A \cap B)$
 - $P(A) = P(AB^C) + P(AB)$
 - $P(B) = P(BA^C) + P(AB)$
 - $P(A) + P(B) = P(AB^C) + P(BA^C) + 2P(AB)$
 - $P(A \cup B) = P(AB^C) + P(BA^C) + P(AB)$

- $P(A \cup B) = P(A) + P(B) - P(AB)$ By induction, we can prove if A_1, \dots, A_n ,

$$P(\cup_{k=1}^n A_k) = \sum_{k=1}^n P(A_k) - \sum_{i < j} P(A_i A_j) + \sum_{i < j < k} P(A_i A_j A_k) + \dots + (-1)^{n+1} P(A_1, \dots, A_n)$$

Inclusion- Exclusion Formula

- If $A_1, \dots, A_n \in \mathcal{F}$,

$$B_1 = A_1$$

$$B_2 = A_2 \setminus A_1$$

$$\vdots$$

Then,

$$\cup_{k=1}^n A_k = \cup_{k=1}^n B_k$$

but the B_i are disjoint. Also $A_k \subseteq B_k \forall k = 1, \dots, n$.

$$P(\cup_{k=1}^n A_k) = P(\cup_{k=1}^n B_k) = \sum_{k=1}^n P(B_k) \leq \sum_{k=1}^n P(A_k)$$

Thus, $P(\cup_{k=1}^n A_k) \leq \sum_{k=1}^n P(A_k)$. Finite subadditivity.

Some conventions,

If A_1, \dots is a sequence of sets, we say $A_n \uparrow A$ if

1. $A_1 \subseteq A_2 \subseteq \dots$
2. $\cup_{k=1}^{\infty} A_k = A$

If A_1, \dots is a sequence of sets, we say $A_n \downarrow A$ if

1. $A_1 \supseteq A_2 \supseteq \dots$
2. $\cap_{k=1}^{\infty} A_k = A$

Theorem 1.2.1 If P is a probability measure on a field \mathcal{F} Then,

1. Continuity from below.

If $A_n \in \mathcal{F} \quad \forall n, A \in \mathcal{F}$

$$A_n \uparrow A$$

then

$$P(A_n) \uparrow P(A)$$

2. Continuity from above.

If $A_n \in \mathcal{F} \quad \forall n, A \in \mathcal{F}$

$$A_n \downarrow A$$

then

$$P(A_n) \downarrow P(A)$$

3. Countable subadditivity.

If $A_n \in \mathcal{F} \quad \forall n, \cup_{k=1}^{\infty} A_k \in \mathcal{F}$ then

$$P(\cup_{n=1}^{\infty} A_k) \leq \sum_{n=1}^{\infty} P(A_k)$$

Proof. 1. If $A_1, \dots, A_n \in \mathcal{F}$,

$$B_1 = A_1$$

$$B_2 = A_2 \setminus A_1$$

$$B_3 = A_3 \setminus A_2$$

$$\vdots$$

then, B_1, \dots are disjoint.

$$\cup_{n=1}^{\infty} A_n = \cup_{n=1}^{\infty} B_n$$

$$\begin{aligned} P(A) &= P(\cup_{n=1}^{\infty} A_n) \\ &= P(\cup_{n=1}^{\infty} B_n) \\ &= \sum_{n=1}^{\infty} P(B_n) \\ &= \lim_{n \rightarrow \infty} \sum_{n=1}^{\infty} P(B_n) \\ &= \lim_{n \rightarrow \infty} P(A_n) \end{aligned}$$

$$2. A_n \downarrow A \Leftrightarrow A_n^C \uparrow A^C$$

But by (1),

$$P(A_n^C) \uparrow P(A^C)$$

$$1 - P(A_n) \uparrow 1 - P(A)$$

$$P(A_n) \downarrow P(A)$$

3. By finite subadditivity,

$$P(\cup_{k=1}^n A_k) \leq \sum_{k=1}^n P(A_k) \leq \sum_{n=1}^{\infty} P(A_n)$$

But since, by (1), because


$$\cup_{k=1}^n A_k \uparrow \cup_{n=1}^{\infty} A_n$$

$$P(\cup_{k=1}^n A_k) \uparrow P(\cup_{n=1}^{\infty} A_n)$$

So,

$$P(\cup_{n=1}^{\infty} A_n) \leq \sum_{n=1}^{\infty} P(A_n)$$

■

 $A \in \mathcal{F} = \text{"A is F-set"}$.

1.3 Extention of Probability Measure to a σ -field

Let f be a function $f : D \rightarrow R$.

Let \tilde{D} be another set such that

$$D \subseteq \tilde{D}$$

An extantion of f onto \tilde{D} is

$$\tilde{f} : \tilde{D} \rightarrow R$$

Such that $f(x) = \tilde{f}(x) \forall x \in D$

\tilde{f} is an extension of f on D .

We say f has unique extension, \tilde{f} onto \tilde{D} if

1. \tilde{f} is an extension of f to \tilde{D} .
2. if g is another extension of f to \tilde{D} then $\tilde{f} = g$ on D .

Theorem 1.3.1 A probability measure on a field has a unique extension on the σ -field generated by this field.

This means that if \mathcal{F}_0 is a field, and P is a probability measure on \mathcal{F}_0 , then there exists a probability measure, Q on $\sigma(\mathcal{F}_0)$ such that

$$Q(A) = P(A) \quad \forall A \in \mathcal{F}_0$$

Moreover, if \tilde{Q} is another probability measure on $\sigma(\mathcal{F}_0)$ such that $\tilde{Q} = P(A) \quad \forall A \in \mathcal{F}_0$ then

$$\tilde{Q} = Q$$



The proof of this theorem will come after several definitions and lemmas.

Outer Measure $P^* : 2^\Omega \rightarrow \mathbb{R}$

For any $A \in 2^\Omega$ ($A \subseteq \Omega$)

$$P^*(A) = \inf \left\{ \sum_{n=1}^{\infty} P(A_n) : A_1, A_2, \dots \text{ is a sequence of } \mathcal{F}_0 \text{ sets, } A \subseteq \bigcup_{n=1}^{\infty} A_n \right\}$$

P^* is a measure out until \mathcal{M} , but it is only a function beyond that on 2^Ω .

Inner Measure

$$P_*(A) = 1 - P^*(A)$$

Define the paving \mathcal{M} as follows

$$\mathcal{M} = \{A \in 2^\Omega : E \in 2^\Omega, P^*(E) = P^*(E \cap A) + P^*(E \cap A^C)\}$$

Idea: we came up with this \mathcal{M} such that P^* behaves as a measure. It will turn out to be that \mathcal{M} is a σ -field that contains $\sigma(\mathcal{F}_0)$.

Monday August 29

P^* satisfies the following probabilities:

- (i) $P^*(\emptyset) = 0$
- (ii) $P^*(A) \geq 0 \quad \forall A \in 2^\Omega$
- (iii) $A \subseteq B \Rightarrow P^*(A) \leq P^*(B)$
- (iv) $P^*(\bigcup_{n=1}^{\infty} A_n) \leq \sum_{n=1}^{\infty} P^*(A_n)$

Proof. (i) Take $\{\emptyset, \emptyset, \dots\}$.

$$\emptyset \in \mathcal{F}_0, \quad \emptyset \cup_{n=1}^{\infty} \emptyset$$

So,

$$P^*(\emptyset) \leq \sum_{n=1}^{\infty} P(\emptyset) = 0$$

Note,

$$P(A) \geq 0 \quad \forall A$$

So,

$$P^*(\emptyset) \geq 0$$

Thus,

$$P^*(\emptyset) = 0$$

(ii) Already done as part of (i).

(iii) Let $A \subseteq B$

$$P^*(A) = \inf\left\{\sum_{n=1}^{\infty} P(A_n), A_n \in \mathcal{F}_0, A \subseteq \bigcup A_n\right\}$$

Now, if $B_1, \dots \in \mathcal{F}_0 \subseteq \bigcup B_n$

Then,

$$A \subseteq B \subseteq \bigcup B_n$$

If $\{\{B_n\}_{n=1}^{\infty} : B_n \in \mathcal{F}_0, B \subseteq \bigcup B_n\} \subseteq \{\{A_n\}_{n=1}^{\infty} : A_n \in \mathcal{F}_0, A \subseteq \bigcup A_n\}$

Or in short, Collection 1 \subseteq Collection 2.

So the inf of a larger set is smaller than (or equal to) the inf of a smaller set.

So,

$$P^*(A) = \inf\left\{\sum_{n=1}^{\infty} P(A_n), A_n \in \text{collection \#1}\right\} \leq P^*(B) = \inf\left\{\sum_{n=1}^{\infty} P(B_n), A_n \in \text{collection \#2}\right\} = P^*(B)$$

(iv) Want

$$P^*(\bigcup_n A_n) \leq \sum_n P^*(A_n)$$

$$P^*(A_n) = \inf\left\{\sum_{k=1}^{\infty} P(A_{nk}) : A_{nk} \in \mathcal{F}_0, A \subseteq \bigcup_k A_{nk}\right\}$$

Let $\varepsilon > 0$, by definition of there exists,

$$\{B_n\}_{n=1}^{\infty}$$

such that

$$\sum_{k=1}^{\infty} P(B_{nk}) \leq P^*(A_n) + \frac{\varepsilon}{2^n}$$

So,

$$\cup_n A_n \subseteq \cup_{n,k} B_{nk}$$

and,

$$\begin{aligned} P^*(\cup_n A_n) &\leq \sum_{n,k} P(B_{nk}) \\ &< \sum_n P^*(A_n) + \sum_n (\varepsilon 2^{-n}) \\ P^*(\cup_n A_n) &< \sum_n P^*(A_n) + \varepsilon \quad \forall \varepsilon > 0 \end{aligned}$$

Simply put,

b

So,

$$P^*(\cup_n A_n) \leq \sum_n P^*(A_n)$$

■

By definition, $A \in \mathcal{M}$ if and only if $P^*(EA) + P^*(EA^C) = P^*(E)$.

We know that P^* is subadditive.

So, by subadditivity we know,

$$P^*(E) \leq P^*(AE) + P^*(A^C E)$$

Therefore, to show $A \in \mathcal{M}$ we only need to show

$$P^*(E) \geq P^*(AE) + P^*(A^C E)$$

\mathcal{M} is defined by P^* and P^* is defined using \mathcal{F}_0 so \mathcal{M} is indirectly tied to \mathcal{F}_0 .

Lemma 1. \mathcal{M} is a field.

Proof. (i) $\Omega \in \mathcal{M}$

$$A = \Omega$$

$$P^*(\emptyset) = 0$$

$$P^*(E) + P^*(\emptyset) = P^*(E)$$

(ii) $A \in \mathcal{M} = A^C \in \mathcal{M}$

$$\begin{aligned}
P^*(E) &= P^*(EA) + P^*(A^C E) \\
&= P^*(EA^C) + P^*(AE) \\
&= P^*(EA^C) + P^*((A^C)^C E)
\end{aligned}$$

(iii) $A, B \in \mathcal{M} \rightarrow A \cap B \in \mathcal{M}$

$$\begin{aligned}
B \in \mathcal{M} &\Rightarrow P^*(E) = P^*(Eb) + P^*(B^C E) \quad \forall E \\
A \in \mathcal{M} &\Rightarrow P^*(BE) = P^*((BE)A) + P^*(A^C (BE)) \\
A \in \mathcal{M} &\Rightarrow P^*(B^C E) = P^*((B^C E)A) + P^*(A^C (B^C E))
\end{aligned}$$

Hence,

$$\begin{aligned}
P^*(BE) + P^*(B^C E) &= P^*((BE)A) + P^*(A^C (BE)) + P^*((B^C E)A) + P^*(A^C (B^C E)) \\
P^*(A^C (BE)) + P^*((B^C E)A) + P^*(A^C (B^C E)) &\geq P^*((A^C BE) \cup (AB^C E) \cup (A^C B^C E)) \\
&= P^*(E \cap [A^C B \cup AB^C \cup A^C B^C]) \\
&= P^*(E \cap (AB)^C)
\end{aligned}$$

$$\begin{aligned}
P^*(E) &= P^*(BE) + P^*(B^C E) \\
&= P^*((BE)A) + (P^*(A^C (BE)) + P^*((B^C E)A) + P^*(A^C (B^C E))) \\
&\geq P^*(ABE) + P^*(E(AB)^C)
\end{aligned}$$

So, $A, B \in \mathcal{M}$

■

Lemma 2. If A_1, A_2, \dots is a sequence of disjoint \mathcal{M} -sets then for each $E \subseteq \Omega$,

$$P^*(E \cap (\cup_k A_k)) = \sum_k P^*(E \cap A_k)$$

Proof. First, prove this statement for finite sequence.

$$A_1, \dots, A_n$$

by mathematical induction.

If $n = 1$ this is 'trivial',

$$P^*(E \cap A_1) = P^*(E \cap A_1)$$

If $n = 2$ we need to show,

$$P^*(E(A_1 \cup A_2)) = P^*(EA_1) + P^*(EA_2)$$

Because $A_1 \in \mathcal{M}$,

$$P^*(E(A_1 \cup A_2)) = P^*(E(A_1 \cup A_2))A_1 + P^*(E(A_1 \cup A_2)A_1^C)$$

$$E(A_1 \cup A_2) = E(A_1 A_2 \cup A_1 A_2^C) = EA_1$$

$$E(A_1 \cup A_2)A_1^C = E(A_1A_1^C \cup A_2A_2^C)$$

So,

$$P^*(E(A_1 \cup A_2)) = P^*(EA_1) + P^*(EA_2)$$

Suppose true for $n = k$. (induction hypothesis)

Now we must show for $n = k + 1$.

$$P^*(E \cap (\cup_{n=1}^{k+1} A_n)) = P^*([E \cap (\cup_{n=1}^k A_n)] \cup A_{k+1})$$

$(\cup_{n=1}^k A_n), A_{k+1}$ are two disjoint sets. Using the $n=2$ case,

$$= \sum_{n=1}^k P^*(E \cap A_n) + P(E \cap A_{k+1}) = \sum_{n=1}^{k+1} P^*(E \cap A_n)$$

So this is now shown to be true for $\{A_1, \dots, A_n\}$. Next, show true for A_1, \dots in \mathcal{M} (disjoint).
Want:

$$P^*(E \cap (\cup_{n=1}^{\infty} A_n)) = \sum_{n=1}^{\infty} P^*(E \cap A_n)$$

Using countable subadditivity,

$$P^*(E \cap (\cup_{n=1}^{\infty} A_n)) = P^*(\cup_{n=1}^{\infty} E \cap A_n) \leq \sum_{n=1}^{\infty} P^*(E \cap A_n)$$

In the meantime, by the monotonicity of P^*

$$P^*(E \cap (\cup_{n=1}^{\infty} A_n)) \geq P^*(E \cap (\cup_{n=1}^m A_n)) = \sum_{n=1}^m P^*(E \cap A_n)$$

So,

$$P^*(E \cap (\cup_{n=1}^{\infty} A_n)) \geq \lim_{m \rightarrow \infty} \sum_{n=1}^m P^*(E \cap A_n)$$

(*), (**) gives us,

$$P^*(E \cap (\cup_{n=1}^{\infty} A_n)) = \sum_{n=1}^{\infty} P^*(E \cap A_n)$$

■

Wednesday August 31

(finished proof)

Lemma 3.

1. \mathcal{M} is a σ -field
2. P^* restricted on \mathcal{M} is countably additive.

Proof. First we show if

1. \mathcal{M} is a field
 2. \mathcal{M} is closed under countable disjoint union.
- then \mathcal{M} is a σ -field.

Let's create disjoint sets,

$$A_n \in \mathcal{M}, n = 1, 2, \dots \quad B_1 = A_1 \quad B_2 = A_2 A_1^C : B_n = A_n A_1^C \dots A_{n-1}^C$$

$$B_1, \dots, B_n \in \mathcal{M} \text{ (disjoint)}$$

$$\bigcup_{n=1}^{\infty} B_n = \bigcup_{n=1}^{\infty} A_n$$

But we know that $\bigcup_{n=1}^{\infty} B_n \in \mathcal{M}$ so $\bigcup_{n=1}^{\infty} A_n \in \mathcal{M}$ and thus \mathcal{M} is a σ -field.

So it suffices to show that \mathcal{M} is closed under disjoint countable unions.

Let A_1, A_2, \dots are disjoint \mathcal{M} -sets.

Let $A = \bigcup_{n=1}^{\infty} A_n$.

Let $F_n = \bigcup_{k=1}^n A_k$.

Then $F_n \in \mathcal{M}$.

So, $\forall E \in 2^{\Omega}$,

$$P^*(E) = P^*(EF_n) + P^*(EF_n^C)$$

$$P^*(EF_n) = P^*(E(\bigcup_{k=1}^n A_k))$$

$$= \sum_{k=1}^n P^*(EA_k)$$

$$P^*(EF_n^C) \geq P^*(EA^C)(F_n \subseteq A, F_n^C \supseteq A^C)$$

$$\Rightarrow P^*(E) \geq \lim_{n \rightarrow \infty} P^*(EA_k) + P^*(EA^C)$$

$$= \sum_{k=1}^{\infty} P^*(EA_k) + P^*(EA^C)$$

$$= P^*(EA) + P^*(EA^C)$$

■

So $A \in \mathcal{M}$ and \mathcal{M} is a σ -field.

Now, let's show P^* is countably additive.

Let A_1, A_2, \dots be disjoint members of \mathcal{M} . Then $\forall E \in 2^{\Omega}$,

$$P^*(E(\bigcup_{n=1}^{\infty} A_n)) = \sum_{n=1}^{\infty} P^*(EA_n)$$

Take $E = \Omega$.

$$P^*(\bigcup_{n=1}^{\infty} A_n) = \sum_{n=1}^{\infty} P(A_n)$$

Lemma 4. $\mathcal{F}_0 \subseteq \mathcal{M}$

Proof. Let $A \in \mathcal{F}$.

Want:

$$A \in \mathcal{M}$$

$$P^*(E) = P^*(EA) + P^*(EA^C)$$

By definition, there exists $E_n \in \mathcal{F}_0$ such that

$$\sum_{n=1}^{\infty} P^*(E_n) \leq P^*(E) + \varepsilon$$

■

Lemma 5.

$$P^*(A) = P(A) \quad \forall A \in \mathcal{F}_0$$

Proof. Let $A \in \mathcal{F}_0$.

Because, $A, \emptyset, \emptyset, \dots, \in \mathcal{F}_0$.

$$A \subseteq A \cup \emptyset \cup \emptyset \dots$$

$$P^*(A) \leq P(A) + P(\emptyset) + \dots$$

But, if

$$A_n \in \mathcal{F}_0$$

$$A \subseteq \bigcup_{n=1}^{\infty} A_n$$

$$P^*(A) \leq \sum_{n=1}^{\infty} P(A_n)$$

$$\Rightarrow P^*(A) \leq \inf \sum_{n=1}^{\infty} P(A_n)$$

$$= P^*(A)$$

■

Friday September 2

Recall, Extension Theorem. That is, If \mathcal{F} is a field and P is a probability measure, then there exists a measure, Q such that

$$Q(A) = P(A) \quad \forall A \in \mathcal{F}_0$$

Proof. By Lemma 5,

$$P^*(\Omega) = P(\Omega) = 1 \quad P^*(\emptyset) = P(\emptyset) = 0$$

Outline of what we need to show:

- $0 \leq M(A) \leq 1$
- $M(\emptyset) = 0, \quad M(\Omega) = 1$

$$\bullet M(\cup_n A_n) = \sum_n M(A_n)$$

Since $\forall A \in \mathcal{M}$,

$$\emptyset \subseteq A \subset \Omega$$

then

$$0 \leq P^*(\emptyset) \leq P^*(A) \leq P^*(\Omega) \leq 1$$

But, by Lemma 3, P^* is contably additive on \mathcal{M} . So P^* is probability measure on \mathcal{M} (which is a σ -field, by Lemma 3).

By Lemma 4, $\mathcal{F}_0 \subset \mathcal{M} \Rightarrow \sigma(\mathcal{F}_0) \subseteq \mathcal{M}$. So P^* is also probability measure on $\sigma(\mathcal{F}_0)$.

Finally, by Lemma 5, again $P^*(A) = P(A)$, P^* is an extension of P from \mathcal{F}_0 to $\sigma(\mathcal{F}_0)$. ■

Uniqueness of the extension, $\pi - \lambda$ Theorem.

Paving - $\{\pi$ -system and λ -system. (?)

A clan of subsets \mathcal{P} of Ω is a π system, if $A, B \in \mathcal{P} \Rightarrow AB \in \mathcal{P}$.

A class \mathcal{L} is a λ -system if

1. $\Omega \in \mathcal{L}$
2. $A \in \mathcal{L} \Rightarrow A^C \in \mathcal{L}$
3. If $A_1, \dots \in \mathcal{L}$ are disjoint then $\cup_{n=1}^{\infty} A_n \in \mathcal{L}$

So, the only difference is "disjoint". So, weaker than a σ -field (i.e. A σ -field is always a λ -system).

Note that (λ_2) can be replace by $(\lambda_{2'})$ wherein

$$A, B \in \mathcal{F}, A \subseteq B, \Rightarrow B \setminus A \in \mathcal{L}$$

That is $\lambda_1, \lambda_2, \lambda_3 \Leftrightarrow \lambda_1, \lambda_{2'}, \lambda_3$

Lemma 6. A class of sets that is both π -systema and λ -system is a σ -field.

Proof. Suppose \mathcal{F} is both π -systema and λ -system.

By definition,

1. $\Omega \in \mathcal{F}$
2. $A \in \mathcal{F} \Rightarrow A^C \in \mathcal{F}$

Let A_1, A_2, \dots be \mathcal{F} sets.

Let's constructs disjoint sets, B

$$B_1 = A_1$$

$$B_2 = A_1 A_2^C$$

$$\vdots$$

Then B_n are \mathcal{F} -sets (by $\lambda_{2'} - A_2^C = \Omega A_2^C \in \mathcal{F}$, by π -system, $A_1 A_2^C \in \mathcal{F}$).

By λ_3 ,

$$\cup_n^{\infty} B_n \in \mathcal{F}$$

So,

$$\cup_n^{\infty} A_n \in \mathcal{F}$$

■

Theorem 1.3.2 — π - λ Theorem. If \mathcal{P} is in a π -system, \mathcal{L} is in a λ -system, then

$$\mathcal{P} \subseteq \mathcal{L} \Rightarrow \sigma(\mathcal{P} \subseteq \mathcal{L})$$

Proof. Let $\lambda(\mathcal{P})$ be the intersection of all λ -system that contains \mathcal{P} .

$$\lambda(\mathcal{P}) = \cap \{ \mathcal{L}' : \mathcal{L}' \supseteq \mathcal{P}, \mathcal{L}' \text{ is } \lambda\text{-set} \}$$

$\lambda(\mathcal{P})$ is a λ -system.

Goal: prove $\lambda(\mathcal{P})$ is a σ -field. So we want to show that $\lambda(\mathcal{P})$ is a π -system.

1. $\Omega \in \lambda(\mathcal{P})$?

$$\Omega \in \mathcal{L}' \quad \forall \mathcal{L}'$$

$$\Omega \in \lambda(\mathcal{P})$$

2. $A \in \lambda(\mathcal{P}) \Rightarrow A^C \in \lambda(\mathcal{P})$?

$$A \in \lambda(\mathcal{P}) \Rightarrow A \in \cap \{ \mathcal{L}' : \mathcal{L}' \supseteq \mathcal{P}, \mathcal{L}' \text{ is } \lambda\text{-set} \}$$

Then

$A \in \mathcal{L}'$ for any $\mathcal{L}' \supseteq \mathcal{P}, \mathcal{L}'$ is λ -system.

$$\Rightarrow A^C \in \mathcal{L}'$$

$$\Rightarrow A^C \in \cap \{ \mathcal{L}' : \mathcal{L}' \supseteq \mathcal{P}, \mathcal{L}' \text{ is } \lambda\text{-set} \} = \lambda(\mathcal{P})$$

3. $A_1, A_2, \dots \in \lambda(\mathcal{P})$ are disjoint then $A_1, A_2, \dots \in \mathcal{L}' \quad \forall \mathcal{L}'$.

Then $\cup A_n \in \mathcal{L}'$ (\mathcal{L}' λ -system)

So $\cup_n A_n \in \lambda(\mathcal{P})$.

We call $\lambda(\mathcal{P})$ the λ -system generated by \mathcal{P} .

If we can say that $\lambda(\mathcal{P})$ is also a σ -field, then $\sigma(\mathcal{P}) \subseteq \lambda(\mathcal{P})$ because $\sigma(\mathcal{P})$ is smallest.

So then, $\sigma(\mathcal{P}) \subseteq \mathcal{L}$ because $\lambda(\mathcal{P})$ is the small λ -system.

So it suffices to show that $\lambda(\mathcal{P})$ is a σ -field. But we know if $\lambda(\mathcal{P})$ is a system then $\lambda(\mathcal{P})$ is σ -field. So it suffices to show that $\lambda(\mathcal{P})$ is a π -system.

Construct again for any $A \in 2^\Omega \quad (A \subseteq \Omega)$, let

$$\mathcal{L}_A = \{ B : AB \in \lambda(\mathcal{P}) \}$$

Claim: If $A \in \lambda(\mathcal{P})$ then \mathcal{L}_A is λ -system.

(a) $\Omega \in \mathcal{L}_A$?

$$A\Omega = A \in \mathcal{L}_A$$

(b) $(\lambda'_2) : B_1, B_2 \in \mathcal{L}_A, B_1 \subseteq B_2 \Rightarrow B_2 B_1^C \in \mathcal{L}_A$?

$$B_1 \in \mathcal{L}_A \Rightarrow AB_1 \in \lambda(\mathcal{P})$$

$$B_2 \in \mathcal{L}_A \Rightarrow AB_2 \in \lambda(\mathcal{P})$$

Since $AB_1 \subseteq AB_2$, $\lambda(\mathcal{P})$ is λ -system by (λ'_2) for $\lambda(\mathcal{P})$

(c) If B_n is disjoint, \mathcal{L}_A -sets.

Want $\cup_n B_n$ because

$$B_n \in \mathcal{L}_A$$

$$B_n A \in \lambda(\mathcal{P})$$

Because B_n disjoint we know that $B_n A$ is also disjoint.

Hence,

$$\cup_n (B_n A) \in \lambda(\mathcal{P})$$

Claim: $\lambda(\mathcal{P})$ is π -system.

(a) If $A \in \mathcal{P}$, then $\mathcal{P} \subseteq \mathcal{L}_A$

Suppose $A \in \mathcal{P}$.

Let $B \in \mathcal{P}$, then $AB \in \mathcal{P}$ (π -system), and $AB \in \lambda(\mathcal{P}) \Rightarrow B \in \mathcal{L}_A$

(b) If $A \in \mathcal{P}$ then $\lambda(\mathcal{P}) \subset \mathcal{L}_A$.

(c) If $A \in \lambda(\mathcal{P})$, then $\mathcal{P} \in \mathcal{L}_A$

Suppose, $A \in \lambda(\mathcal{P})$ and let $B \in \mathcal{P}$.

By step 2,

$$A \in \mathcal{L}_A$$

$$\Rightarrow AB \in \lambda(\mathcal{P})$$

$$\Rightarrow B \in \mathcal{L}_A$$

(d) If $A \in \lambda(\mathcal{P})$, then $\lambda(\mathcal{P}) \subseteq \mathcal{L}_A$. This is because $\lambda(\mathcal{P})$ is the smallest λ -system, \mathcal{L}_A is λ -system containing \mathcal{P} (by step 3).

Now show that $\lambda(\mathcal{P})$ is π -system.

$A, B \in \lambda(\mathcal{P})$ because $A \in \lambda(\mathcal{P})$. We have that $\lambda(\mathcal{P}) \in \mathcal{L}_A$.

So

$$B \in \mathcal{L}_A$$

$$BA \in \lambda(\mathcal{P})$$

Thus $\lambda(\mathcal{P})$ is π -system. ■

Wednesday September 7

Theorem 1.3.3 Suppose P_1 and P_2 are probability measures on $\sigma(\mathcal{P})$ where \mathcal{P} is a π -system. If P_1 and P_2 agree on \mathcal{P} (that is, $P_1(A) = P_2(A) \quad \forall A \in \mathcal{P}$) then they agree on $\sigma(\mathcal{P})$.

Proof. Let

$$\mathcal{L} = \{A \in \sigma(\mathcal{P}) : P_1(A) = P_2(A)\}$$

Then $\mathcal{P} \subseteq \mathcal{L}$.

It suffices to show that \mathcal{L} is a λ -system (because if so, then $\sigma(\mathcal{P}) \subseteq \mathcal{L}$ - in fact, $\sigma(\mathcal{P}) = \mathcal{L}$).

Show \mathcal{L} is a λ -system.

1. $\Omega \in \mathcal{L}$?

$$P_1(\Omega) = P_2(\Omega) = 1, \quad \Omega \in \mathcal{P}$$

2. $A \in \mathcal{L}$

$$P_1(A) = P_2(A) \Rightarrow P_1(A^C) = P_2(A^C), \quad A^C \in \mathcal{L}$$

3. $A \in \mathcal{L}$. A_n disjoint. Want $\cup_{n=1}^{\infty} A_n \in \mathcal{L}$.

Since

$$A_n \in \mathcal{L}$$

$$P_1(A_n) = P_2(A_n) \quad \forall n$$

$$\sum_{n=1}^{\infty} P_1(A_n) = \sum_{n=1}^{\infty} P_2(A_n)$$

$$P_1(\cup_{n=1}^{\infty} A_n) = P_2(\cup_{n=1}^{\infty} A_n)$$

So, $\cup A_n \in \mathcal{L}$.

■

So our extension of (and uniqueness of the extension of) P on \mathcal{F}_0 to $\sigma(\mathcal{F}_0)$ is complete. We have shown the existence of Q on \mathcal{M} .

Since Q agrees with P on \mathcal{F}_0 and \mathcal{F}_0 is a field, this implies that this is a π -system.

If you have another extension, say \tilde{Q} , then $\tilde{Q} = P$ on \mathcal{F}_0 . That is, $\tilde{Q} = Q$ on \mathcal{M} , where \mathcal{M} is a σ -field, which is a π -system.

So by Theorem 1.3.3,

$$\tilde{Q} = Q \text{ on } \sigma(\mathcal{P}).$$

So, Theorem 1.3.1, is completely proved.

Lemma 1 - Lemma 5 imply the extension.

$\pi - \lambda$ Theorem and Theorem 1.3.3 implies uniqueness.

This wraps up Theorem 1.3.1.

Lebesgue measure on $(0,1]$

$$\Omega = (0, 1]$$

Recall, \mathcal{B}_0 is the finite disjoint unions of intervals in $(0,1]$ and that \mathcal{B}_0 is a field.

Let $\mathcal{B} = \sigma(\mathcal{B}_0)$.

For each $A \in \mathcal{B}_0$,

$$A = \cup_{i=1}^n (a_i, b_i]$$

$$\text{Let } \lambda(A) = \sum_{i=1}^n (b_i - a_i).$$

Question: Is λ a probability measure on \mathcal{B}_0 ?

Theorem 1.3.4 — Theorem 2.2 in Billingsly. The set function λ on \mathcal{B}_0 is a probability measure on \mathcal{B}_0 .

Proof. 1. $0 \leq \lambda(A) \leq 1$
 2.

$$\lambda(\Omega) = \lambda((0, 1]) = 1 - 0 = 1$$

$$\lambda(\emptyset) = \lambda((0, 0]) = 0$$

3. This one requires Theorem 1.3 in Billingsly (proof omitted - calculus, blah).
 Theorem 1.3 - If I is an interval in $(0, 1]$ and $\{I_k : k = 1, 2, \dots\}$ are disjoint intervals in $(0, 1]$ such that

$$I = \bigcup_{k=1}^{\infty} I_k$$

then,

$$|I| = \sum_{k=1}^{\infty} |I_k|$$

where $|a|$ means length of interval a .

■



2. General Measure



3. Integration with Respect to a Measure



4. Random Variable



5. Convergence in Probability/Limit Theorem



6. Radon-Nikodym Derivative Theorem



7. Special Topics



Index

Exten. Prob Measure to σ -field, 14

Probability on a Field, 7