



# Probability Theroy based on Measure Theory

STAT 517

Dr. John Smith



Copyright © 2013 John Smith

PUBLISHED BY PUBLISHER

BOOK-WEBSITE.COM

Licensed under the Creative Commons Attribution-NonCommercial 3.0 Unported License (the “License”). You may not use this file except in compliance with the License. You may obtain a copy of the License at <http://creativecommons.org/licenses/by-nc/3.0>. Unless required by applicable law or agreed to in writing, software distributed under the License is distributed on an “AS IS” BASIS, WITHOUT WARRANTIES OR CONDITIONS OF ANY KIND, either express or implied. See the License for the specific language governing permissions and limitations under the License.

*First printing, March 2013*

# Contents

I	Part One	
1	Probability Measure .....	7
1.1	Probability on a Field .....	7
1.2	Extention of Probability Measure to a $\sigma$ -field .....	13
2	General Measure .....	19
3	Integration with Respect to a Measure .....	21
4	Random Variable .....	23
5	Convergence in Probability/Limit Theorem .....	25
6	Radon-Nikodym Derivative Theorem .....	27
7	Special Topics .....	29
	Index .....	31



# Part One

<b>1</b>	<b>Probability Measure</b> .....	<b>7</b>
1.1	Probability on a Field	
1.2	Extention of Probability Measure to a $\sigma$ -field	
<b>2</b>	<b>General Measure</b> .....	<b>19</b>
<b>3</b>	<b>Integration with Respect to a Measure</b>	<b>21</b>
<b>4</b>	<b>Random Variable</b> .....	<b>23</b>
<b>5</b>	<b>Convergence in Probability/Limit Theorem</b>	<b>25</b>
<b>6</b>	<b>Radon-Nikodym Derivative Theorem</b> .	<b>27</b>
<b>7</b>	<b>Special Topics</b> .....	<b>29</b>
	<b>Index</b> .....	<b>31</b>



# 1. Probability Measure

## 1.1 Probability on a Field

■ **Definition 1.1.1** —  $\Omega$ . Non empty set.

■ **Definition 1.1.2** — **Paving**. A collection of a subset of  $\Omega$  is a paving.

■ **Definition 1.1.3** — **Field**. A field  $\mathcal{F}$  is a paving satisfying

- (i)  $\Omega \in \mathcal{F}$
- (ii)  $A \in \mathcal{F} \Rightarrow A^C \in \mathcal{F}$
- (iii)  $A, B \in \mathcal{F} \Rightarrow A \cup B \in \mathcal{F}$

### Derived Properties about a Field

- $\emptyset \in \mathcal{F}$  (by (i) and (ii):

$$\begin{aligned}\Omega \in \mathcal{F} &\Rightarrow \Omega^C \in \mathcal{F} \\ &\Rightarrow \emptyset \in \mathcal{F})\end{aligned}$$

- (i) can be replaced by " $\mathcal{B}$  is nonempty" because,  
Let  $A \in \mathcal{F}$ ,

$$\begin{aligned}&\Rightarrow A^C \in \mathcal{F} \\ &\Rightarrow A^C \cup A \in \mathcal{F} \\ &\Rightarrow \Omega \in \mathcal{F}\end{aligned}$$

- $A \in \mathcal{F}, B \in \mathcal{F} \Rightarrow A \cap B \in \mathcal{F}$  because,

$$\begin{aligned}(A \cap B)^C &= A^C \cup B^C \text{ (DeMorgan's Law)} \\ A \cap B &= (A^C \cup B^C)^C\end{aligned}$$

- $A_1, \dots, A_m \in \mathcal{F} \Rightarrow A_1 \cup \dots \cup A_m \in \mathcal{F}$  (mathematical induction)
- $A_1, \dots, A_m \in \mathcal{F} \Rightarrow A_1 \cap \dots \cap A_m \in \mathcal{F}$

**Definition 1.1.4 —  $\sigma$ -Field.** Similar to the definition of a field except for (iii). A paving satisfying

- (i)  $\Omega \in \mathcal{F}$
- (ii)  $A \in \mathcal{F} \Rightarrow A^c \in \mathcal{F}$
- (iii)  $A_1 \in \mathcal{F}, \dots, A_m \in \mathcal{F}$   
 $\bigcup_{k=1}^m A_k \in \mathcal{F}$  (finite additivity)

If we replace (iii) from before by (iii') here:

For  $A_1 \in \mathcal{F}, \dots, A_m \in \mathcal{F}$

$$\bigcup_{k=1}^{\infty} A_k \in \mathcal{F}$$

then  $\mathcal{F}$  is called a  **$\sigma$ -field**.

### Derived Facts

- Again, (i) can be replaced by  $\mathcal{F}$  not empty, (iii) can be replaced by  $A_1 \in \mathcal{F}, \dots, A_m \in \mathcal{F}$

■ **Example 1.1**  $\Omega = (0, 1]$  (from now on all intervals are left open, right closed)

**R** Recall that  $\sigma$ -fields are generated by fields. Fancy scripts denote a  $\sigma$ -field. Fancy scripts with a zero subscript denote a field.

$\mathcal{B}_0$  is the collection of all finite union of disjoint intervals. Asside: something that we can consider easily, but computers cannot. (e.g. think of all stars, etc)




Figure 1.1: Finite union of three disjoint intervals.

Then  $\mathcal{B}_0$  is a field.

- (i)  $(0, 1] \in \mathcal{B}_0$
- (ii)  $A \in \mathcal{B}_0 \Rightarrow A^c \in \mathcal{B}_0$
- (iii)  $A \in \mathcal{B}_0, B \in \mathcal{B}_0 \Rightarrow A \cup B \in \mathcal{B}_0$

**Wednesday August 24**

$\mathcal{B}_0$  = collection of finite unions of disjoint subintervals of  $(0, 1]$ . Is a field.







Figure 1.2: A and complement of A.

Figure 1.3: Union of A and B is still in  $\mathcal{B}_0$ 

**Definition 1.1.5 — Power Set.** A  $\sigma$ -field is generated by a paving of power set. Let  $\Omega$  be a set. The collection of all subsets of  $\Omega$  is the power set written as  $2^\Omega$ .

- R** Where does this notation come from?  
Consider this case where  $\Omega$  is finite

$$\Omega = \{\omega_1, \dots, \omega_n\}$$

Total number of subsets of  $\Omega$ .

$\emptyset$ , 1 element sets, 2-element sets, ..., n-element sets.

$$() + () + \dots + = (1 + 1)^n$$

$\#(\mathcal{F}) = 2^{\#\Omega}$ , so it seems reasonable to denote  $\mathcal{F} = 2^\Omega$ .

It is also easy to show that  $2^\Omega$  is a  $\sigma$ -field. (The largest, even. The smallest:  $\{\emptyset, \Omega\}$  which is also a  $\sigma$ -field.)

$$\{\emptyset, \Omega\} \subseteq \sigma\text{-field} \subseteq 2^\Omega$$

It turns out we can extend notion of length from  $\mathcal{B}_0$  to  $\sigma$ -field generated by  $\mathcal{B}_0$ .

Now, let  $\mathcal{A}$  be a nonempty paving of  $\Omega$ . We define

$$\sigma(\mathcal{A}) = \cap \{ \mathcal{B} \subseteq 2^\Omega : \mathcal{B} \text{ is a } \sigma\text{-field, } \mathcal{A} \subseteq \mathcal{B} \}$$

OR rather, the *intersection* of all  $\sigma$ -fields that contains  $\mathcal{A}$ .

Let

$$\mathbb{F}(\mathcal{A}) = \{ \mathcal{B} \subseteq 2^\Omega : \mathcal{B} \text{ is a } \sigma\text{-field, } \mathcal{B} \supseteq \mathcal{A} \}$$

Then,


$$\sigma(\mathcal{A}) = \cap \mathcal{B}$$

$$\mathcal{B} \in \mathbb{F}(\mathcal{A})$$

### Derived Facts

$\mathbb{F}(\mathcal{A})$  is nonempty. For example,  $2^\Omega$  is a  $\sigma$ -field and  $2^\Omega \supseteq \mathcal{A}$ .

$\cap \mathcal{B}$  is a  $\sigma$ -field. ( $\mathcal{B} \in \mathbb{F}(\mathcal{A})$ )

 Get notes about notation/levels.

*Proof.* We will prove that indeed  $\sigma(\mathcal{A})$  is a  $\sigma$ -field. Recall that we have three conditions above for  $\sigma$ -field.

(i)

$$\Omega \in \sigma(\mathcal{A})$$

$$\Omega \in \cap_{\mathcal{B} \in \mathbb{F}(\mathcal{A})} \mathcal{B}$$

Because:  $\mathcal{B}$  is  $\sigma$ -field,  $\Omega \in \mathcal{B}, \forall \mathcal{B} \in \mathbb{F}(\mathcal{A})$ .

(ii)

$$(iii) \quad A_1, \dots, \in \cap_{\mathcal{B} \in \mathbb{F}(\mathcal{A})} \mathcal{B}, \forall \mathcal{B} \in \mathbb{F}(\mathcal{A})$$

$$\Rightarrow \cup_{n=1}^{\infty} A_n \in \mathcal{B}, \forall \mathcal{B} \in \mathbb{F}(\mathcal{A})$$

So,  $\sigma(\mathcal{A})$  is a  $\sigma$ -field, we call it the  $\sigma$ -field, generated by  $\mathcal{B}_0$ . We know how to assign length to members of  $\mathcal{B}_0$ , we now show the assignment can be extended to  $\sigma(\mathcal{B}_0)$  ■

■ **Example 1.2** Let  $\mathcal{I}$  be the collection of *all* subintervals of  $(0,1]$ .

Note that  $\mathcal{I}$  is a smaller collection than  $\mathcal{B}_0$  since  $\mathcal{B}_0$  can have numerous different combinations of the sets.

Let

$$\mathcal{B} = \sigma(\mathcal{I})$$

This is a Borel- $\sigma$ -field. (a member of  $\mathcal{B}$  is Borel set.)

It turns out

$$\sigma(\mathcal{I}) = \sigma(\mathcal{B}_0)$$

This is because  $\sigma(\mathcal{I})$  is a  $\sigma$ -field.

So,

$$\begin{aligned}\sigma(\mathcal{I}) &\supseteq \mathcal{B}_o \\ \sigma(\mathcal{I}) &\supseteq \sigma(\mathcal{B}_o)\end{aligned}$$

Also,

$$\begin{aligned}\mathcal{I} &\subseteq \mathcal{B}_o \\ \sigma(\mathcal{I}) &\subseteq \sigma(\mathcal{B}_o)\end{aligned}$$

Thus,

$$\sigma(\mathcal{I}) = \sigma(\mathcal{B}_o)$$

■

**Definition 1.1.6 — Probability Measure.** Probability measures on field. Suppose  $\mathcal{F}$  is a field on a nonempty set  $\Omega$ . A probability measure is a function  $P : \mathcal{F} \rightarrow \mathbb{R}$ .

- (i)  $0 \leq P(A) \leq 1, \forall A \in \mathcal{F}$
- (ii)  $P(\emptyset) = 0, P(\Omega) = 1$
- (iii) If  $A_1, \dots$  are disjoint members of  $\mathcal{F}$  and  $\cup A_n \in \mathcal{F}$  then we have countable additivity:

$$P(\cup A_n) = \sum_{n=1}^{\infty} P(A_n)$$

**R** Note that (iii) also implies finite additivity. Prove by adding infinite empty sets on end.

If  $\Omega$  is nonempty set. And  $\mathcal{F}$  is a  $\sigma$ -field on  $\Omega$ . And  $P$  is a probability measure on  $\mathcal{F}$ . Then  $(\Omega, \mathcal{F}, P)$  is called a **probability space**. And  $(\Omega, \mathcal{F})$  is called a **measurable space**.

**R** If  $A \subseteq B$ , then  $P(A) \leq P(B)$ . This is because we may write  $B$  as

$$B = A \cup (B \setminus A)$$

**R**

$$P(A) + P(B) = P(A \cup B) + P(A \cap B)$$

### Friday August 26

Recall,

Probability measure on a field,  $\mathcal{F}_0$ .

- $P(A) + P(B) = P(A \cup B) + P(A \cap B)$ 
  - $P(A) = P(AB^C) + P(AB)$
  - $P(B) = P(BA^C) + P(AB)$
  - $P(A) + P(B) = P(AB^C) + P(BA^C) + 2P(AB)$
  - $P(A \cup B) = P(AB^C) + P(BA^C) + P(AB)$

- $P(A \cup B) = P(A) + P(B) - P(AB)$  By induction, we can prove if  $A_1, \dots, A_n$ ,

$$P(\cup_{k=1}^n A_k) = \sum_{k=1}^n P(A_k) - \sum_{i < j} P(A_i A_j) + \sum_{i < j < k} P(A_i A_j A_k) + \dots + (-1)^{n+1} P(A_1, \dots, A_n)$$

Inclusion- Exclusion Formula

- If  $A_1, \dots, A_n \in \mathcal{F}$ ,

$$B_1 = A_1$$

$$B_2 = A_2 \setminus A_1$$

$$\vdots$$

Then,

$$\cup_{k=1}^n A_k = \cup_{k=1}^n B_k$$

but the  $B_i$  are disjoint. Also  $A_k \subseteq B_k \forall k = 1, \dots, n$ .

$$P(\cup_{k=1}^n A_k) = P(\cup_{k=1}^n B_k) = \sum_{k=1}^n P(B_k) \leq \sum_{k=1}^n P(A_k)$$

Thus,  $P(\cup_{k=1}^n A_k) \leq \sum_{k=1}^n P(A_k)$ . Finite subadditivity.

Some conventions,

If  $A_1, \dots$  is a sequence of sets, we say  $A_n \uparrow A$  if

1.  $A_1 \subseteq A_2 \subseteq \dots$
2.  $\cup_{k=1}^{\infty} A_k = A$

If  $A_1, \dots$  is a sequence of sets, we say  $A_n \downarrow A$  if

1.  $A_1 \supseteq A_2 \supseteq \dots$
2.  $\cap_{k=1}^{\infty} A_k = A$

**Theorem 1.1.1** If  $P$  is a probability measure on a field  $\mathcal{F}$  Then,

1. Continuity from below.

If  $A_n \in \mathcal{F} \forall n, A \in \mathcal{F}$

$$A_n \uparrow A$$

then

$$P(A_n) \uparrow P(A)$$

2. Continuity from above.

If  $A_n \in \mathcal{F} \forall n, A \in \mathcal{F} A_n \downarrow A$ , then  $P(A_n) \downarrow P(A)$

3. Countable subadditivity.

If  $A_n \in \mathcal{F} \forall n, \cup_{k=1}^{\infty} A_k \in \mathcal{F}$  then

$$P(\cup_{n=1}^{\infty} A_k) \leq \sum_{n=1}^{\infty} P(A_k)$$

*Proof.* 1. If  $A_1, \dots, A_n \in \mathcal{F}$ ,

$$B_1 = A_1$$

$$B_2 = A_2 \setminus A_1$$

$$B_3 = A_3 \setminus A_2$$

$$\vdots$$

then,  $B_1, \dots$  are disjoint.

$$\cup_{n=1}^{\infty} A_n = \cup_{n=1}^{\infty} B_n$$

$$P(A) = P(\cup_{n=1}^{\infty} A_n) = P(\cup_{n=1}^{\infty} B_n) = \sum_{n=1}^{\infty} P(B_n) = \lim_{n \rightarrow \infty} \sum_{n=1}^n P(B_n) = \lim_{n \rightarrow \infty} P(A_n)$$

$$2. A_n \downarrow A \Leftrightarrow A_n^C \uparrow A^C$$

But by (1),

$$P(A_n^C) \uparrow P(A^C) \quad 1 - P(A_n) \uparrow 1 - P(A) \quad P(A_n) \downarrow P(A)$$

3. By finite subadditivity,

$$P(\cup_{k=1}^n A_k) \leq \sum_{k=1}^n P(A_k) = n P(A_k) \leq \sum_{k=1}^{\infty} P(A_k)$$

But since, by (1), because

$$\cup_{k=1}^n A_k \uparrow \cup_{n=1}^{\infty} A_n$$

$$P(\cup_{k=1}^n A_k) \uparrow P(\cup_{n=1}^{\infty} A_n)$$

So,

$$P(\cup_{n=1}^{\infty} A_n) \leq \sum_{n=1}^{\infty} P(A_n)$$

■

**R**  $A \in \mathcal{F} = \text{"A is F-set"}$ .

## 1.2 Extention of Probability Measure to a $\sigma$ -field

Let  $f$  be a function  $f : D \rightarrow R$ .

Let  $\tilde{D}$  be another set such that

$$D \subseteq \tilde{D}$$

An extantion of  $f$  onto  $\tilde{D}$  is

$$\tilde{f} : \tilde{D} \rightarrow R$$

Such that  $f(x) = \tilde{f}(x) \forall x \in D$

$\tilde{f}$  is an extention of  $f$  on  $D$ .

We say  $f$  has unique extention,  $\tilde{f}$  onto  $\tilde{D}$  if

1.  $\tilde{f}$  is an extension of  $f$  to  $\tilde{D}$ .
2. if  $g$  is another extension of  $f$  to  $\tilde{D}$  then  $\tilde{f} = g$  on  $D$ .

**Theorem 1.2.1** A probability measure on a field has a unique extension on the  $\sigma$ -field generated by this field.

Means:  $\mathcal{F}_0$  is a field

$P$  is a probability measure on  $\mathcal{F}_0$

Then there exists a probability measure,  $Q$  on  $\sigma(\mathcal{F})$  such that  $Q(A) = P(A) \forall A \in \mathcal{F}_0$

Moreover, if  $\tilde{Q}$  is another probability measure on  $\sigma(\mathcal{F}_0)$  such that  $\tilde{Q} = P(A) \forall A \in \mathcal{F}$  then  $\tilde{Q} = Q$ .

**Outer Measure**  $P^* : 2^\Omega \rightarrow \mathbb{R}$   
 For any  $A \in 2^\Omega$  ( $A \subseteq \Omega$ )

$$P^*(A) = \inf \left\{ \sum_{n=1}^{\infty} P(A_n) : A_1, A_2, \dots \text{ is a sequence of } \mathcal{F}_0 \text{ sets, } A \subseteq \bigcup_{n=1}^{\infty} A_n \right\}$$

**Inner Measure**

$$P_*(A) = 1 - P^*(A)$$

Define the paving  $\mathcal{M}$  as follows

$$\mathcal{M} = \{A \in 2^\Omega : E \in 2^\Omega, P^*(E) = P^*(E \cap A) + P^*(E \cap A^C)\}$$

**Monday August 29**

$P^*$  satisfies the following probabilities:

- (i)  $P^*(\emptyset) = 0$
- (ii)  $P^*(A) \geq 0 \quad \forall A \in 2^\Omega$
- (iii)  $A \subseteq B \Rightarrow P^*(A) \leq P^*(B)$
- (iv)  $P^*(\bigcup_{n=1}^{\infty} A_n) \leq \sum_{n=1}^{\infty} P^*(A_n)$

*Proof.* (i) Take  $\{\emptyset, \emptyset, \dots\}$ .

$$\emptyset \in \mathcal{F}_0, \quad \emptyset \subseteq \bigcup_{n=1}^{\infty} \emptyset$$

So,

$$P^*(\emptyset) \leq \sum_{n=1}^{\infty} P(\emptyset) = 0$$

Note,

$$P(A) \geq 0 \quad \forall A$$

So,

$$P^*(\emptyset) \geq 0$$

Thus,

$$P^*(\emptyset) = 0$$

(ii) Already done as part of (i).

(iii) Let  $A \subseteq B$

$$P^*(A) = \inf \left\{ \sum_{n=1}^{\infty} P(A_n), A_n \in \mathcal{F}_0, A \subseteq \bigcup_{n=1}^{\infty} A_n \right\}$$

Now, if  $B_1, \dots \in \mathcal{F}_0 \subseteq \bigcup B_n$

Then,

$$A \subseteq B \subseteq \bigcup_n B_n$$

If  $\{\{B_n\}_{n=1}^\infty : B_n \in \mathcal{F}_0, B \subseteq \cup_n B_n\} \subseteq \{\{A_n\}_{n=1}^\infty : A_n \in \mathcal{F}_0, A \subseteq \cup_n A_n\}$   
 Or in short, Collection 1  $\subseteq$  Collection 2.

So the inf of a larger set is smaller than (or equal to) the inf of a smaller set.

So,

$$P^*(A) = \inf\left\{\sum_{n=1}^\infty P(A_n), A_n \in \text{collection \#1}\right\} \leq P^*(B) = \inf\left\{\sum_{n=1}^\infty P(B_n), A_n \in \text{collection \#2}\right\} = P^*(B)$$

(iv) Want

$$P^*(\cup_n A_n) \leq \sum_n P^*(A_n)$$

$$P^*(A_n) = \inf\left\{\sum_{k=1}^\infty P(A_{nk}) : A_{nk} \in \mathcal{F}_0, A \subseteq \cup_k A_{nk}\right\}$$

Let  $\varepsilon > 0$ , by definition of there exists,

$$\{B_n\}_{n=1}^\infty$$

such that

$$\sum_{k=1}^\infty P(B_{nk}) \leq P^*(A_n) + \frac{\varepsilon}{2^n}$$

So,

$$\cup_n A_n \subseteq \cup_{n,k} B_{nk}$$

and,

$$\begin{aligned} P^*(\cup_n A_n) &\leq \sum_{n,k} P(B_{nk}) \\ &< \sum_n P^*(A_n) + \sum_n (\varepsilon 2^{-n}) \\ P^*(\cup_n A_n) &< \sum_n P^*(A_n) + \varepsilon \quad \forall \varepsilon > 0 \end{aligned}$$

Simply put,

$b$

So,

$$P^*(\cup_n A_n) \leq \sum_n P^*(A_n)$$

■

By definition,  $A \in \mathcal{M}$  if and only if  $P^*(EA) + P^*(EA^C) = P^*(E)$ .

We know that  $P^*$  is subadditive.

So, by subadditivity we know,

$$P^*(E) \leq P^*(AE) + P^*(A^C E)$$

Therefore, to show  $A \in \mathcal{M}$  we only need to show

$$P^*(E) \geq P^*(AE) + P^*(A^C E)$$

$\mathcal{M}$  is defined by  $P^*$  and  $P^*$  is defined using  $\mathcal{F}_0$  so  $\mathcal{M}$  is indirectly tied to  $\mathcal{F}_0$ .

**Lemma 1.**  $\mathcal{M}$  is a field.

*Proof.* (i)  $\Omega \in \mathcal{M}$

$$\begin{aligned} A &= \Omega \\ P^*(\emptyset) &= 0 \\ P^*(E) + P^*(\emptyset) &= P^*(E) \end{aligned}$$

(ii)  $A \in \mathcal{M} = A^C \in \mathcal{M}$

$$\begin{aligned} P^*(E) &= P^*(EA) + P^*(A^C E) \\ &= P^*(EA^C) + P^*(AE) \\ &= P^*(EA^C) + P^*((A^C)^C E) \end{aligned}$$

(iii)  $A, B \in \mathcal{M} \rightarrow A \cap B \in \mathcal{M}$

$$\begin{aligned} B \in \mathcal{M} &\Rightarrow P^*(E) = P^*(Eb) + P^*(B^C E) \quad \forall E \\ A \in \mathcal{M} &\Rightarrow P^*(BE) = P^*((BE)A) + P^*(A^C(BE)) \\ A \in \mathcal{M} &\Rightarrow P^*(B^C E) = P^*((B^C E)A) + P^*(A^C(B^C E)) \end{aligned}$$

Hence,

$$\begin{aligned} P^*(BE) + P^*(B^C E) &= P^*((BE)A) + P^*(A^C(BE)) + P^*((B^C E)A) + P^*(A^C(B^C E)) \\ P^*(A^C(BE)) + P^*((B^C E)A) + P^*(A^C(B^C E)) &\geq P^*((A^C BE) \cup (AB^C E) \cup (A^C B^C E)) \\ &= P^*(E \cap [A^C B \cup AB^C \cup A^C B^C]) \\ &= P^*(E \cap (AB)^C) \end{aligned}$$

$$\begin{aligned} P^*(E) &= P^*(BE) + P^*(B^C E) \\ &= P^*((BE)A) + (P^*(A^C(BE)) + P^*((B^C E)A) + P^*(A^C(B^C E))) \\ &\geq P^*(ABE) + P^*(E(AB)^C) \end{aligned}$$

So,  $A, B \in \mathcal{M}$

■

**Lemma 2.** If  $A_1, A_2, \dots$  is a sequence of disjoint  $\mathcal{M}$ -sets then for each  $E \subseteq \Omega$ ,

$$P^*(E \cap (\cup_k A_k)) = \sum_k P^*(E \cap A_k)$$

*Proof.* First, prove this statement for finite sequence.

$$A_1, \dots, A_n$$



by mathematical induction.

If  $n = 1$  this is 'trivial',

$$P^*(E \cap A_1) = P^*(E \cap A_1)$$

If  $n = 2$  we need to show,

$$P^*(E(A_1 \cup A_2)) = P^*(EA_1) + P^*(EA_2)$$

Because  $A_1 \in \mathcal{M}$ ,

$$P^*(E(A_1 \cup A_2)) = P^*(E(A_1 \cup A_2))A_1 + P^*(E(A_1 \cup A_2)A_1^C)$$

$$E(A_1 \cup A_2) = E(A_1 A_2 \cup A_1^C A_2) = EA_1$$

$$E(A_1 \cup A_2)A_1^C = E(A_1 A_1^C \cup A_2 A_1^C)$$

So,

$$P^*(E(A_1 \cup A_2)) = P^*(EA_1) + P^*(EA_2)$$

■





## 2. General Measure





### 3. Integration with Respect to a Measure





## 4. Random Variable







## 5. Convergence in Probability/Limit Theorem





## 6. Radon-Nikodym Derivative Theorem





## 7. Special Topics





## Index

Exten. Prob Measure to  $\sigma$ -field, 13

Probability on a Field, 7