

# Étale Fundamental Groups

Organised by Matt Byrne

October 9, 2026

## Contents

|                                                                 |          |
|-----------------------------------------------------------------|----------|
| <b>1 Motivation and examples (Mark)</b>                         | <b>2</b> |
| 1.1 Fundamental groups and covers: the classical case . . . . . | 2        |
| 1.2 An interesting coincidence . . . . .                        | 3        |
| 1.2.1 A little bit of number theory . . . . .                   | 3        |
| 1.2.2 A little bit of complex geometry . . . . .                | 4        |
| 1.2.3 Weil's Rosetta stone . . . . .                            | 4        |
| 1.2.4 A bonus "rad" fact . . . . .                              | 4        |

## Boring admin stuff

We're meeting every Thursday at 13:00-14:00 in J11 Hicks, beginning 8<sup>th</sup> October. Notes will be updated (hopefully) shortly after each talk, and will be emailed out once they're finished. The main reference is Section 5 of Tamás Szamuley's *Galois Groups and Fundamental Groups*.

# 1 Motivation and examples (Mark)

## 1.1 Fundamental groups and covers: the classical case

Let  $X$  be a topological space. One historically important problem is that of classifying the *covering spaces* of  $X$ .

**Definition 1.1.** Suppose  $X$  and  $Y$  are topological spaces. A continuous map  $p : Y \rightarrow X$  is called a *cover* if for all  $x \in X$ , there is an open neighbourhood  $U_x \subseteq X$  of  $x$  whose preimage decomposes as a disjoint union of homeomorphic copies of  $U_x$ , that is, we may decompose

$$p^{-1}[U_x] = \coprod_{i \in I} V_{x,i}$$

with each  $V_{x,i} \subseteq Y$  open and  $p|_{V_{x,i}} : V_{x,i} \xrightarrow{\sim} U_x$  a homeomorphism. We say that  $Y$  is a *covering space* for  $X$ . If  $X$  and  $Y$  are additionally pointed (i.e. have distinguished basepoints  $x_0 \in X, y_0 \in Y$ ), and the cover  $p : (Y, y_0) \rightarrow (X, x_0)$  satisfies  $p(y_0) = x_0$ , we may say that  $(Y, y_0)$  is a *pointed covering space* of  $(X, x_0)$ .

These covering spaces of  $X$  define a category  $\mathbf{Cov}(X)$ , whose morphisms are continuous maps  $Y_1 \rightarrow Y_2$  commuting with the covers  $p_i : Y_i \rightarrow X$ . We will write  $\mathbf{FinCov}(X)$  for the full subcategory of finite covers, and  $\mathbf{Cov}_*(X, x_0)$  for the subcategory of pointed covers (with pointed morphisms).

One notable property of covering spaces is the *homotopy lifting property*: that given a homotopy  $\{f_t : Z \rightarrow X\}_{t \in [0,1]}$  of maps to  $X$ , and a lift  $g_0 : Z \rightarrow Y$  of  $f_0$  through  $p : Y \rightarrow X$ , there exists a unique homotopy  $\{\hat{f}_t : Z \rightarrow Y\}_{t \in [0,1]}$  lifting  $\{f_t\}_t$  through  $p : Y \rightarrow X$  such that  $\hat{f}_0 = g_0$ .

As a consequence of the homotopy lifting property, their fibres admit a *monodromy action*: given a cover  $p : Y \rightarrow X$  and some  $x_0 \in X$ , the fibre  $p^{-1}(x_0) \subseteq Y$  is a discrete subspace. Given a point  $y \in p^{-1}(x_0)$  and some class  $[\alpha] \in \pi_1(X, x_0)$ , which we recall is a strong homotopy equivalence class of loops in  $X$  based at  $x_0$ , any representative  $\alpha : [0, 1] \rightarrow X$  lifts to a unique  $\hat{\alpha}_y : [0, 1] \rightarrow Y$  starting at  $y$  whose strong homotopy class depends only on the strong homotopy class  $[\alpha]$ . Thus the end-point  $\hat{\alpha}_y(1)$  does not depend on the choice of lift, and we define  $[\alpha] \cdot y := \hat{\alpha}_y(1)$ .

So we can think of the taking the fibre at  $x_0$  as a functor

$$\text{Fib}_{x_0} : \mathbf{Cov}(X) \rightarrow \pi_1(X, x_0)\text{-Set}$$

from the category of covering spaces of  $X$  to the category of  $\pi_1(X, x_0)$ -sets, i.e. discrete sets with an action of  $\pi_1(X, x_0)$ . As it turns out, when  $X$  is especially nice (path-connected, locally path-connected, and semilocally simply connected) this is a *representable functor*, inasmuch as

$$\text{Fib}_{x_0}(Y) \cong \text{Hom}_{\mathbf{Cov}(X)}(\tilde{X}, Y)$$

for some  $\tilde{X} \in \mathbf{Cov}(X)$  unique up to isomorphism. Such  $\tilde{X}$  is called a *universal covering space* for  $X$ , and may be constructed explicitly by choosing a basepoint  $x_0 \in X$  and defining

$$\tilde{X}_{x_0} := \frac{\{\gamma : [0, 1] \xrightarrow{\text{cts}} X \mid \gamma(0) = x_0\}}{\text{homotopy}}, \quad \tilde{p} : \tilde{X}_{x_0} \rightarrow X; \gamma \mapsto \gamma(1).$$

By definition, we see that  $\text{Fib}_{x_0}(\tilde{X}_{x_0}) = \pi_1(X, x_0)$ , and thus by universality we see

$$\pi_1(X, x_0) \cong \text{Hom}_{\mathbf{Cov}(X)}(\tilde{X}_{x_0}, \tilde{X}_{x_0}) = \text{Aut}_{\mathbf{Cov}(X)}(\tilde{X}_{x_0})$$

This final isomorphism is sometimes referred to as the *deck transformation construction*.

**Theorem 1.2.** *Pointed covers of a path-connected, locally path-connected, and semilocally simply connected pointed space  $(X, x_0)$  are in bijection with subgroups of  $\pi_1(X, x_0)$ . To each  $H \leq \pi_1(X, x_0)$  we associate  $(\tilde{X}_{x_0}/H, Hx_0)$ , and to each  $p : (Y, y_0) \rightarrow (X, x_0)$  we associate  $p_*\pi_1(Y, y_0) \leq \pi_1(X, x_0)$ .*

The main idea of this reading group will be to define a sort of covering space theory for schemes, by way of the *étale fundamental group*. But how do we do this?

| Topological spaces                                                    | Schemes                                                                          |
|-----------------------------------------------------------------------|----------------------------------------------------------------------------------|
| $X$ a nice topological space                                          | $S$ a nice scheme                                                                |
| $p : Y \rightarrow X$ a topological cover                             | $\phi : X \rightarrow S$ a <i>finite étale cover</i>                             |
| $x_0 \in X$ a point                                                   | $\overline{s_0} : \text{Spec}(\Omega) \rightarrow S$ a <i>geometric point</i>    |
| $\pi_1(X, x_0)$ defined in terms of loops/paths                       | Paths are meaningless in Zariski!                                                |
| $\pi_1(X, x_0) = \text{Aut}(\text{Fib}_{x_0})$                        | $\pi_1^{\text{ét}}(S, \overline{s_0}) = \text{Aut}(\text{Fib}_{\overline{s_0}})$ |
| $\tilde{X} \rightarrow X$ universal cover (generally infinite degree) | $\mathfrak{X} \in \text{pro}(\mathbf{Fét}_S)$ a <i>universal pro-object</i>      |
| $\pi_1(X, x_0) \cong \text{Aut}(\tilde{X})$                           | $\pi_1^{\text{ét}}(S, s_0) \cong \varprojlim_{X \in \mathfrak{X}} \text{Aut}(X)$ |

In fact for complex affine varieties  $X_{/\mathbb{C}}$  we will be able to define both the classical fundamental group  $\pi_1(X^{\text{an}}, x_0)$  (considering its complex points  $X(\mathbb{C})$  canonically retopologised as a complex manifold  $X^{\text{an}}$ ) and the étale fundamental group  $\pi_1^{\text{ét}}(X, \overline{x_0})$  (considering it as a scheme, where  $\overline{x_0}$  is the geometric point induced by  $x_0$ ), and in this case the two *almost* coincide:

$$\pi_1^{\text{ét}}(X, \overline{x_0}) \cong \widehat{\pi_1(X^{\text{an}}, x_0)},$$

where the only difference is a profinite completion on the right, appearing since we are only considering *finite* covers as we will see later.

The approach we will take is to use this recontextualisation of  $\pi_1(X, x_0)$  as an automorphism group as the basis for our definition of  $\pi_1^{\text{ét}}$ . After defining all of the scheme theory we will need, we will try to reformulate the theory of covering spaces independent of loops so we can do it for schemes. There will be some interesting differences on

## 1.2 An interesting coincidence

One slightly more specific example which might motivate the need for a theory of étale covering space is the famous *Weil Rosetta stone*. This is a set of strange coincidences between the theories of Riemann surfaces and of number fields that seem to suggest that both are offshoots of a more general theory (which will turn out to be our theory of the étale fundamental group). To state this, we first must define some things. We will start with the number fields setting.

### 1.2.1 A little bit of number theory

**Definition 1.3.** A *number field*  $K$  is a finite field extension of  $\mathbb{Q}$ . That is, it is a field containing  $\mathbb{Q}$  which is finite-dimensional as a  $\mathbb{Q}$ -vector space. Each number field  $K$  admits a unique ring  $\mathcal{O}_K$  called its *ring of integers*, which is the integral closure of  $\mathbb{Z}$  in  $K$ .

In each extension  $L/K$  of number fields, we can look at prime ideals  $\mathfrak{p}$  of the ring of integers  $\mathcal{O}_K$  of  $K$  (which we be called ‘primes of  $K$ ’ for short). Each such prime must factorise in  $\mathcal{O}_L$  as a unique product

$$\mathfrak{p}\mathcal{O}_L = \mathfrak{P}_1^{e_1} \cdots \mathfrak{P}_r^{e_r}$$

of powers of primes  $\mathfrak{P}_i$  of  $L$ . We say that  $\mathfrak{p}$  *ramifies* in  $L/K$  if any of the  $e_i > 1$ , otherwise we say it is *unramified*.

**Example 1.4.** In the extension  $\mathbb{Q}(i)/\mathbb{Q}$ , the ideal  $5\mathbb{Z}[i]$  factorises as  $\langle 2+i \rangle \langle 2-i \rangle$ , so 5 is unramified. In contrast, the ideal  $2\mathbb{Z}[i]$  factorises as  $\langle 1+i \rangle^2$  since  $(1+i)(1-i) = 2$ , so 2 ramifies.  $\diamond$

If a set  $S$  of primes of  $K$  contains all of the primes at which  $L/K$  ramifies, we say that  $L/K$  is *unramified away from  $S$* , or just  *$S$ -unramified*.

Given an algebraic closure  $\overline{K}/K$ , we can form the *maximal  $S$ -unramified extension*  $K_S/K$  by taking the union of all finite subextensions  $L/K$  of  $\overline{K}/K$  which are  $S$ -unramified. Then we define the  *$S$ -unramified absolute Galois group*  $G_{K,S}$  to be topological group

$$G_{K,S} := \text{Gal}(K_S/K) := \varprojlim_{L/K} \text{Aut}_K(L)$$

where  $L/K$  ranges over all finite,  $S$ -unramified extensions.

**Theorem 1.5.** *The open subgroups of  $G_{K,S}$  are in bijection with the set of finite,  $S$ -unramified extensions of  $K$  contained in  $\bar{K}$ . An open subgroup  $H$  is sent to its fixed field  $K_S^H$ , and a finite  $S$ -unramified extension  $L/K$  is sent to the open subgroup  $\text{Gal}(K_S/L)$ .*

There is an underlying geometric object here, called  $\text{Spec}(\mathcal{O}_{K,S})$  (see next week's talk). We will see that finite  $S$ -unramified extensions correspond to finite étale covers  $X \twoheadrightarrow \text{Spec}(\mathcal{O}_{K,S})$ , and a choice of algebraic closure  $\bar{K}$  corresponds to a choice of *geometric basepoint*  $\text{Spec}(\bar{K})$ .

### 1.2.2 A little bit of complex geometry

**Definition 1.6.** A *Riemann surface* is a smooth complex manifold of dimension 1.

We are interested in holomorphic maps between compact Riemann surfaces, so fix a surjective holomorphic map  $\phi : Y \rightarrow X$ . Interestingly, any such map is a finite topological cover away from a finite set of points of  $X$ , which are called the *ramification points* of  $\phi$ . If a discrete subset  $S \subset X$  contains all of the ramification points of  $\phi$ , we say that  $\phi$  is *unramified away from  $S$* .

Since any finite cover of Riemann surface is itself a Riemann surface, Theorem 1.2 specialises in this case to give a specific correspondence between pointed,  $S$ -unramified holomorphic covers of  $(X, x_0)$  and finite-index subgroups of  $\pi_1(X \setminus S, x_0)$ , which is equivalent to the following statement about the profinite completion  $\widehat{\pi}_1$  of the fundamental group.

**Theorem 1.7.** *The open subgroups of  $\widehat{\pi}_1(X \setminus S, x_0)$  are in bijection with finite,  $S$ -unramified, pointed holomorphic covers of  $(X, x_0)$ . An open subgroup  $H$  is sent to the compactification of the orbit space  $\widehat{X \setminus S}_{x_0} / H$  with basepoint  $Hx_0$ , and a finite,  $S$ -unramified, pointed holomorphic cover  $\phi : (Y, y_0) \rightarrow (X, x_0)$  is sent to the open subgroup*

$$\phi_* \widehat{\pi}_1(Y \setminus \phi^{-1}[S], y_0) \subseteq \widehat{\pi}_1(X \setminus S, x_0).$$

There is an underlying algebraic object here, that being the field  $K(X)$  of meromorphic functions on  $X$ , and the subring  $\mathcal{O}_{X,S}$  of meromorphic functions which are holomorphic away from  $S$ . Any holomorphic cover  $\phi : Y \rightarrow X$  realises a field extension  $\phi^* : K(X) \hookrightarrow K(Y)$ , and it sends  $\mathcal{O}_{X,S}$  to  $\mathcal{O}_{Y, \phi^{-1}[S]}$ .

### 1.2.3 Weil's Rosetta stone

We can formalise this correspondence later, but for now there's a long list of things that seem to be similar. We will write the objects of interest, as well as what the 'étale equivalent' will be.

| Compact Riemann surface $X$                     | Number field $K$                               | Étale equivalent                                                                                   |
|-------------------------------------------------|------------------------------------------------|----------------------------------------------------------------------------------------------------|
| $X$                                             |                                                | $\text{Spec}(\mathcal{O}_K)$                                                                       |
| $X'$                                            |                                                | $\text{Spec}(\mathcal{O}_{K,S})$                                                                   |
| $\phi : Y \twoheadrightarrow X$ holomorphic     | $\iota : K \hookrightarrow L$ finite extension | $\iota^* : \text{Spec}(\mathcal{O}_L) \twoheadrightarrow \text{Spec}(\mathcal{O}_K)$ finite smooth |
| $S$ ramification points                         | $S$ ramified primes                            | $S$ ramification points                                                                            |
| $\phi$ is $S$ -unramified                       | $L/K$ is $S$ -unramified                       | $\iota^*$ is $S$ -unramified                                                                       |
| $\phi _{Y'} \in \mathbf{FinCov}(X \setminus S)$ |                                                | $\iota^* _{\text{Spec}(\mathcal{O}_{L,S})} \in \mathbf{Fét}(\text{Spec}(\mathcal{O}_{K,S}))$       |
| $x_0 \in X'$ basepoint                          | $\bar{K}/K_S/K$ algebraic closure              | $\bar{s}_0 : \text{Spec}(\bar{K}) \rightarrow \text{Spec}(\mathcal{O}_{K,S})$ geom. point          |
| $\widehat{\pi}_1(X', x_0)$                      | $G_{K,S} := \text{Gal}(K_S/K)$                 | $\pi_1^{\text{ét}}(\text{Spec}(\mathcal{O}_{K,S}), \bar{s}_0)$                                     |
| $\phi^{-1}(x)$                                  | Primes above $\mathfrak{p} \notin S$           | $\text{Spec}(\mathcal{O}_{L,S}) \times_{\text{Spec}(\mathcal{O}_{K,S})} \text{Spec}(\Omega)$       |
| Classification of hol. covers                   | Galois theory                                  | Still to come...                                                                                   |

Here we write  $X' := X \setminus S$  and  $Y' := Y \setminus \phi^{-1}[S]$  for brevity's sake.

### 1.2.4 A bonus "rad" fact

Though not entirely in the spirit of finite étale covers, the analogy between compact Riemann surfaces and number fields is even deeper than this. This subsection is only included for fun and is very informal, so feel free to skip it!

One of the greatest achievements of early 20<sup>th</sup> Century number theory was the development of *class field theory*. The point of class field theory is to classify every so-called abelian number field, i.e. number fields with abelian Galois group over  $\mathbb{Q}$ . Along with the development of a theory of analysis over  $\mathbb{Q}$  (i.e. analysis over the ring  $\mathbb{A}_{\mathbb{Q}}$  of adèles of  $\mathbb{Q}$ ), it is known for some fairly mysterious isomorphisms. One of the most notable being:

$$G_{K,S}^{\text{ab}} \cong K^{\times} \backslash \mathbb{A}_{K,\text{fin}}^{\times} / U_S, \quad U_S := \prod_{v \notin S} U_v.$$

The double quotient group here may be thought of as the Picard group of  $\text{Spec}(\mathcal{O}_{K,S})$ , i.e. we may think of this as a canonical isomorphism

$$G_{K,S}^{\text{ab}} \cong \text{Pic}(\text{Spec } \mathcal{O}_{K,S}).$$

Simultaneously on the other side, if we take the (punctured) compact Riemann surface  $X' := X \setminus S$  we may consider its *Jacobian variety*

$$\text{Jac}(X') := H^0(X', \Omega_{X'}^1)^* / H_1(X')$$

where a 1-chain  $\gamma$  is seen as a functional on global holomorphic differentials via the usual pairing  $(\gamma, \omega) \mapsto \int_{\gamma} \omega$ . Then *geometric class field theory* gives another mysterious canonical isomorphism

$$\pi_1(X', x_0)^{\text{ab}} \otimes \mathbb{R}/\mathbb{Z} \cong \text{Jac}(X').$$

Since  $\text{Jac}(X')$  may be seen as the identity component of the Picard variety, Weil's Rosetta stone suggests that these two statements are somewhat analogous. Writing  $Z' := \text{Spec}(\mathcal{O}_{K,S})$  and  $\overline{z_0} : \text{Spec}(\overline{K}) \rightarrow Z'$ , the statements even kind of looks the same:

$$\pi_1^{\text{ét}}(Z', \overline{z_0})^{\text{ab}} \cong \text{Pic}(Z') \iff \pi_1(X', x_0)^{\text{ab}} \otimes \mathbb{R}/\mathbb{Z} \cong \text{Pic}^{\circ}(X').$$

Perhaps the interesting most interesting thing here, however, is why these statements are different. Why the extra  $\mathbb{R}/\mathbb{Z}$ ? Why  $\pi_1$  instead of  $\widehat{\pi_1} = \pi_1^{\text{ét}}$ ? Why  $\text{Jac} = \text{Pic}^{\circ}$  instead of  $\text{Pic}$ ? The answers to these questions are simple... I don't know. But we may find out in one of the bonus lectures!