# GeoNews — Week 45/2026

**Weekly digest on Quaternary palaeoclimate research, East African Rift dynamics, palaeohydrology and time-series analysis · 2 November 2026**

> **What is GeoNews?** An **automatically generated weekly digest**. A research workflow scans publications from the past one to two weeks as well as funding, policy and community sources, scores the hits against a fixed set of criteria and assembles this issue from them.
>
> The thematic profile is **aligned with the research interests of Markus L. Fischer** (Palaeoclimate Dynamics group, University of Potsdam): Quaternary palaeoclimate and palaeoenvironmental reconstruction, the East African Rift, the African Humid Period and human–environment coupling, lake-level and water-balance modelling, palaeofloods and fluvial geomorphology, ICDP drill cores, and linear and nonlinear time-series analysis. Researchers in these fields will probably find something useful here — it is explicitly not a complete overview of Quaternary science.
>
> Selection, scoring and summaries are generated automatically. Entries may contain errors; when in doubt, the linked original source prevails.

*Each paper receives a score of 0–100 from five criteria: topical fit (40), methodological fit (25), impact/venue (15), connectivity (10) and recency (10). Categories: 🔴 75+ must-read · 🟡 50–74 relevant · 🟢 30–49 radar. The score measures proximity to this newsletter's profile and prioritises reading — it is not a quality judgement on the individual paper.*

---

## In brief

About 35 scored papers, this week with **five 🔴 must-reads**, all abstract-based. Focal points: **lake highstands and Bayesian chronology**, a **drilling initiative along the East African Rift – Dead Sea Transform corridor**, **millennial Holocene variability**, and two papers on **human–environment coupling in the African Middle and Late Pleistocene**.

At the top: **Lake Woods (Jurrkulu)**, northern Australia's largest freshwater lake, with OSL-dated highstands at 195 ± 16 ka and 98 ± 7 ka and a Bayesian-modelled filling event between 95.9 and 53.8 ka (*JQS*); the workshop report on **Lake KINneret Drilling (KIND)**, which aims at a Plio-Quaternary reference record for the Levant and a meridional synthesis along the East African Rift (*Scientific Drilling*); and a **PNAS modelling study** linking a self-sustained ~1350-year Holocene oscillation to tropical–Antarctic ocean–atmosphere interactions. Further 🔴: **non-analog plant environments at early *Homo sapiens* sites in South Africa** (Florisbad, new luminescence ages) and a **continent-wide morphometric analysis of Middle Stone Age points**, whose shape is best explained by palaeoenvironment and region.

In the 🟡 block, among others: a **nonlinear ENSO–seasonal-cycle combination mode** for bimodal Horn of Africa rainfall (*GRL*), **8200 years of wind variability from Svalbard lake sediments**, and a **review of early-warning systems for terrestrial biosphere tipping points** (*Reviews of Geophysics*).

Funding: **ICDP AGU travel grants (8 Nov)**, **ERC Proof of Concept opens (12 Nov)**, **Horizon Europe palaeoclimate call opens (17 Nov)**, **ERC Consolidator (12 Jan 2027)**, **ICDP**

**proposals (15 Jan 2027)**. Policy: **public hearing on the German WissZeitVG reform on Wednesday, 4 Nov 2026**; **final budget committee session on the 2027 federal budget on 12 Nov**; deadline for a Council–Parliament agreement on the **2027 EU budget** around **16 Nov**.

---

## Part A — Papers of the week

Covered are papers with a Crossref `created` date from 19 October 2026 onward that were not listed in a previous issue. For every entry with an accessible abstract (OpenAlex or Crossref), the assessment is based on that abstract; titles without an available abstract are marked ⚠ *title-based*.

### Must-read (75+)

**1. Late Quaternary lake filling and drying in northern Australia's largest freshwater lake** · *Journal of Quaternary Science*, 2026 · **Score 84** Lake Woods (Jurrkulu) fills today during strong Indo-Australian summer monsoons to a maximum depth of ~5 m; two arcuate ridges 20–40 m above the lake floor point to much higher past fillings. The authors combine **20 single-grain OSL ages**, basin hypsometry, catchment-wide denudation rates and rare earth elements for provenance. Findings per abstract: two highstands at **195 ± 16 ka** and **98 ± 7 ka** (20 and 15 m above the lake floor) plus a third filling event (18.5 m) with a **Bayesian-modelled boundary age of 95.9–53.8 ka** (95% credible interval). They find no evidence for the previously reported late MIS 3 highstand; Holocene deposits do not allow robust quantification of filling episodes, and the lake floor shows strong post-depositional mixing (maximum ages < 143 ka). Denudation rates (0.83–2.05 m/Myr) are among the lowest in northern Australia, so the monsoon record is, according to the authors, only discontinuous — with exceptional episodic filling in MIS 5–4. Assessment: a case study for shoreline chronologies in shallow, episodically filled basins; methodologically transferable to reconstructing lake highstands in East African rift basins. *Topic 38 · Method 20 · Venue 8 · Connectivity 8 · Recency 10 (created 28 Oct 2026)* · DOI: 10.1002/jqs.70130

**2. Workshop report: Lake KINneret Drilling (KIND) – reconstructing Pliocene–Quaternary hydroclimate and seismic history of the Levant** · *Scientific Drilling*, 2026 · **Score 83** Workshop report (no new measurements). Lake Kinneret occupies a pull-apart basin on the Dead Sea Transform; seismic profiles and short cores show a thick, authigenic-calcite-dominated, largely stratigraphically ordered sequence since the late Miocene. Project aim per abstract: an **exceptionally long Plio-Quaternary chronostratigraphic reference record** with potentially sub-centennial resolution for hydroclimate, palaeoseismicity, geomagnetism and palaeobotany — as a building block of a **meridional synthesis of continental and marine core records along the East African Rift – Dead Sea Transform corridor**. The region is also a main corridor of out-of-Africa dispersal. Assessment: directly connected to existing rift-lake drilling in East Africa and relevant for planning comparative syntheses along the corridor. Drilling status and funding are not stated in the abstract. *Topic 40 · Method 16 · Venue 7 · Connectivity 10 · Recency 10 (created 29 Oct 2026)* · DOI: 10.5194/sd-35-255-2026

**3. Holocene millennial climate variability influenced by tropical–Antarctic ocean–atmosphere interactions** · *PNAS*, 2026 · **Score 81** A transient Holocene simulation with EC-Earth (8–0 ka) plus a global compilation of **304 temperature reconstructions** identify, per abstract, a robust, **self-sustained ~1350-year oscillatory mode**. Within each cycle, Antarctic

Bottom Water (AABW) strengthens with developing warming and an El Niño-like tropical SST response; positive feedbacks (sea-ice albedo, tropical air–sea interaction, SST–AABW) amplify the warming until reduced brine rejection from sea-ice retreat weakens AABW and initiates the cooling phase. The dominant period matches proxy-based estimates and is confirmed by further simulations. The authors propose this as an alternative to solar or deep-ocean explanations (interpretation; a model result, not direct proxy evidence of the mechanism). Relevant for spectral and phase analyses of Holocene proxy series, where millennial cycles are often attributed to solar forcing. *Topic 30 · Method 20 · Venue 15 · Connectivity 6 · Recency 10 (created 27 Oct 2026)* · DOI: 10.1073/pnas.2538010123

**4. Non-analog environments supported early Homo sapiens in southern Africa 200 000 years ago** · *Communications Earth & Environment*, 2026 · **Score 80** A **plant-wax n-alkane isotope record** for the sites **Florisbad** and **Lombard's Drift** (South Africa), anchored by **new luminescence ages that revise the age of the Florisbad cranium** (the revised value is not given in the abstract). Findings per abstract: contrasting, **non-analog plant environments** during the Middle and Late Pleistocene, supported by multiple lines of evidence. The authors see this as a basis for evaluating the cultural adaptation strategies of early *Homo sapiens* through time. Assessment: thematically close to human–environment questions in East African archives; "non-analog" is a caution against modern transfer functions. *Topic 36 · Method 15 · Venue 11 · Connectivity 8 · Recency 10 (created 30 Oct 2026)* · DOI: 10.1038/s43247-026-04116-4

**5. Morphological variation in Middle-Late Pleistocene stone points: a pan-African perspective** · *Quaternary Science Reviews*, 2026 · **Score 79** The first **continent-wide geometric morphometric analysis** (elliptic Fourier analysis, multivariate statistics) of almost 1000 points from 44 assemblages in eastern, northern and southern Africa. Findings per abstract: shape is dominated by two axes (elongation vs. breadth; basal tanging). **Modelled palaeoenvironment (mainly annual precipitation and temperature seasonality) and region** are the strongest predictors; only the North African Aterian forms a consistently distinct group, while Howiesons Poort and Still Bay points (smaller sample) fall largely within "generic" MSA variability. Raw material acts as an additional constraint. Assessment: a quantitative link between artefact variability and palaeoclimate — its robustness depends on the quality of the modelled palaeoenvironmental fields. *Topic 34 · Method 16 · Venue 11 · Connectivity 8 · Recency 10 (created 29 Oct 2026)* · DOI: 10.1016/j.quascirev.2026.110314

## Relevant (50–74)

**6. Seasonal Modulation of ENSO Teleconnections to Horn of Africa Rainfall** · *Geophysical Research Letters*, 2026 · **Score 74** Nonlinear interactions between ENSO and the semiannual and annual ITCZ migration generate, per abstract, a **combination mode (C-Mode)** with significant spectral peaks near 1 and 2 $yr^{-1}$ that modulates timing and amplitude of both rainy seasons (OND, MAM) over the Horn of Africa; including it substantially improves the representation of the bimodal rainfall response. Instrumental timescale, but methodologically close to combination tones in proxy spectra — and relevant for interpreting rainfall seasonality in the southern Ethiopian Rift. *Topic 25 · Method 20 · Venue 11 · Connectivity 8 · Recency 10 (created 30 Oct 2026)* · DOI: 10.1029/2026gl123196

**7. Geochemical and granulometric fingerprints of 8200 years of westerly wind variability in inner-fjord lake sediments from Svalbard** · *Boreas*, 2026 · **Score 72** Laminated sediments from a closed-basin lake on Spitsbergen; geochemical, visual and grain-size fingerprints within a multiproxy geostatistical framework link aeolian input to specific source areas (Fe-rich coarse silt from upwind dolerites). Findings per abstract: four phases of enhanced

aeolian activity in the Mid and Late Holocene; shifts in sources coincide with the end of the Holocene Thermal Maximum and the onset of the Neoglacial, but the wind signal itself remains comparatively stable over the long term, with no clear correspondence to regional palaeoclimate records. Methodologically transferable to provenance analysis of aeolian fractions in lake sediments. *Topic 30 · Method 18 · Venue 8 · Connectivity 6 · Recency 10 (created 26 Oct 2026)* · DOI: 10.1111/bor.70089

**8. Science Needs for a Terrestrial Biosphere Tipping Point Early Warning System** · *Reviews of Geophysics*, 2026 · **Score 72** Review: how can Earth-observation data be used for early-warning systems of biosphere tipping points? A synthesis of potential tipping elements and mathematical methods for resilience estimation, emphasising spatio-temporal resilience indicators that exploit short but spatially dense EO series, including space-for-time substitution and parallel forcing data. No palaeodata; conceptually connected to early-warning analyses of proxy series, which face the opposite problem (long but spatially sparse) (assessment). *Topic 30 · Method 18 · Venue 11 · Connectivity 3 · Recency 10 (created 27 Oct 2026)* · DOI: 10.1029/2025rg000900

**9. Anchoring Anatolian tree-rings and Mesopotamian chronology with single-year radiocarbon patterns and ice-core evidence** · *PNAS*, 2026 · **Score 71** Single-year $^{14}C$ patterns in Anatolian timbers anchor a combined tree-ring chronology for **2072–744 BCE**; together with archaeological and high-resolution ice-core evidence, the remaining dating uncertainty supports, per abstract, the **middle–low chronology** for Mesopotamia (2000–1500 BCE). It also yields an annually resolved palaeoclimate record. Methodologically relevant for wiggle-matching and coupling independent chronologies. *Topic 25 · Method 18 · Venue 15 · Connectivity 3 · Recency 10 (created 26 Oct 2026)* · DOI: 10.1073/pnas.2615871123

**10. Monsoon zonal seesaw: Development of the South Asian monsoon aridified North Africa in the Cenozoic** · *Science Advances*, 2026 · **Score 69** Palaeoclimate ensembles plus a model of the atmospheric response to imposed diabatic heating: per abstract, **early Cenozoic North African aridification** was primarily driven by the intensification (and poleward shift) of the South Asian monsoon via the monsoon–desert mechanism (large-scale subsidence over North Africa), rather than being an independent response to palaeogeography. The authors identify a persistent dynamical pathway between the two monsoon systems whose expression at other times depended on competing forcings. Pre-Quaternary, but relevant as a dynamical framework for Africa–Asia coupling on orbital timescales as well (assessment). *Topic 25 · Method 13 · Venue 15 · Connectivity 6 · Recency 10 (created 30 Oct 2026)* · DOI: 10.1126/sciadv.aee1112

## Radar

- **Evidence from 2024 Storm Boris indicates anthropogenic climate change amplifies floods more than extreme precipitation** · *Science Advances* · **69** · Storyline approach with two climate and two hydrological models across hundreds of catchments: without anthropogenic climate change, per abstract 6–9% less precipitation but 14–22% lower flood peaks — amplification factor 2.1–2.9. Context for Central European flood research. DOI: 10.1126/sciadv.aef7703 · created 28 Oct 2026
- **Late Holocene wetting facilitated human expansion in the Qaidam Basin** · *Palaeogeography, Palaeoclimatology, Palaeoecology* · **67** · ⚠ *title-based* · Human–environment coupling in an arid basin; findings with caveats. DOI: 10.1016/j.palaeo.2026.114318 · created 31 Oct 2026

- **Late-glacial and Holocene vegetation dynamics in response to climate, land use and fire at Lago di Andromia (Ossola, Italian Alps)** · *Quaternary Science Reviews* · **67** 🟡 · Pollen, stomata, charcoal, macrofossils over ~14 kyr: larch from ~11.5 ka, silver fir by ~10.5 ka at the latest; earliest pastoralism with fire clearance ~5.5 ka, marked opening from the Early Bronze Age. DOI: 10.1016/j.quascirev.2026.110302 · created 28 Oct 2026
- **Quantifying structural and proxy composition uncertainty in Common Era climate field reconstructions using online paleo data assimilation** · *EGUsphere* (preprint) · **65** 🟡 · Particle filter with Linear Inverse Models; factorial ensembles separate internal, structural and proxy-composition uncertainty; structural uncertainty (mainly PSM calibration) dominates the amplitude of secular temperature variability. Not yet peer reviewed. DOI: 10.5194/egusphere-2026-6303 · created 28 Oct 2026
- **Decoupled δ²H and effective-moisture records reveal vapor-source control on Holocene precipitation isotope in West Asia** · *Global and Planetary Change* · **64** 🟡 · ⚠ *title-based* · Relevant for interpreting isotope proxies as moisture indicators. DOI: 10.1016/j.gloplacha.2026.105787 · created 30 Oct 2026
- **A snow-jet bridging mechanism driving the Afro-Asian summer monsoon precipitation dipole** · *npj Climate and Atmospheric Science* · **63** 🟡 · Large snow-perturbation ensembles plus information-flow causal analysis: reduced Tibetan Plateau snow cover favours, via an accelerated tropical easterly jet, a dipole (more rainfall over tropical Africa, less over South Asia). DOI: 10.1038/s41612-026-01582-7 · created 21 Oct 2026
- **Distinct phases of abrupt Indian monsoon variability during 4800–3500 yr BP in the northeastern Himalaya with a prominent 4.2 ka event** · *Palaeogeography, Palaeoclimatology, Palaeoecology* · **62** 🟡 · ⚠ *title-based*. DOI: 10.1016/j.palaeo.2026.114321 · created 31 Oct 2026
- **Sea Surface Conditions and Antarctic Bottom Water Formation Offshore Cape Darnley, East Antarctica, Over the Last 600 ka** · *Paleoceanography and Paleoclimatology* · **62** 🟡 · Diatoms and geochemistry: MIS 13 with indications of ice-shelf melt, MIS 1 with a signal of enhanced frazil-ice formation. DOI: 10.1029/2026pa005435 · created 26 Oct 2026
- **Contrasting organic carbon responses to climate variability over the past millennium in hydrologically diverse lakes on the northeastern Tibetan Plateau** · *Quaternary Science Reviews* · **62** 🟡 · ⚠ *title-based*. DOI: 10.1016/j.quascirev.2026.110359 · created 28 Oct 2026
- **Interaction between the East Asian summer monsoon and Westerlies as shown by desert shrub-ring records from the Alxa Plateau** · *Climate of the Past* · **62** 🟡 · 26 shrub-ring chronologies; dry–wet variability spatially heterogeneous and modulated by the interplay of westerlies and the East Asian summer monsoon. Open access. DOI: 10.5194/cp-22-1927-2026 · created 26 Oct 2026
- **An Arctic ice shelf in the Last Glacial Maximum rivalled Antarctica's modern ice shelves** · *Nature Geoscience* · **61** 🟡 · ⚠ *title-based*. DOI: 10.1038/s41561-026-02127-6 · created 26 Oct 2026
- **Stochastic simulation of Yangtze River fluvial terraces via Bayesian generalized Coupled Markov Chains** · *Stochastic Environmental Research and Risk Assessment* · **60** 🟡 · ⚠ *title-based* · Bayesian stochastic modelling of fluvial terraces. DOI: 10.1007/s00477-026-03394-w · created 21 Oct 2026
- **Holocene water mass structure changes in Lake Nurume, Lützow-Holm Bay, East Antarctica** · *Progress in Earth and Planetary Science* · **58** 🟡 · Multiple cores from different

water depths, diatoms, XRF scanning and GIA palaeotopography; sedimentation resumes ~6.6 cal ka BP under marine conditions. DOI: 10.1186/s40645-026-00849-0 · created 29 Oct 2026

- **Impact of Ocean Physical Conditions on Oceanic Carbon Pumps and Atmospheric $CO_2$ at the Last Glacial Maximum** · *Paleoceanography and Paleoclimatology* · **58** 🟡 · One biogeochemical model with physical fields from 12 PMIP forcings: mean glacial $pCO_2$ reduction 40.3 ppmv, of which −30.1 ppmv from SST cooling. DOI: 10.1029/2026pa005487 · created 23 Oct 2026
- **Temperature-sensitivity of water stable isotopes in global precipitation: a groundwater noble gas-based, paleoclimate perspective** · *Advances in Geochemistry and Cosmochemistry* · **57** 🟡 · 746 fossil groundwater samples (39 locations, ~35 kyr): the most consistent $\delta^{18}O$–temperature relationship in Europe. DOI: 10.33063/agc.v2i2.1097 · created 19 Oct 2026
- **Glacial $CO_2$ drawdown by plankton metabolic balance** · *Nature Communications* · **57** 🟡 · ⚠ *title-based*. DOI: 10.1038/s41467-026-78422-y · created 29 Oct 2026
- **Sedimentation in Lake Pleshcheyevo (East European Plain) in the Late Holocene under conditions of increased anthropogenic impact** · *Quaternary* · **57** 🟡 · 3600 years of lake sediments; two age–depth models compared via anthropogenic pollen indicators; eutrophication from ~1900 cal BP with slash-and-burn agriculture. DOI: 10.3390/quat9060073 · created 23 Oct 2026
- **The last glacial cycle in western Norway — extensive MIS 5d glaciation, multiple glacial advances and LGM glacial advance phases** · *Boreas* · **55** 🟡 · $^{14}C$ ages (25–50 ka) treated as minimum ages; amino acid epimerization points to MIS 5c; five major Weichselian advances. DOI: 10.1111/bor.70092 · created 23 Oct 2026
- **Radiocarbon dating of highly diverse Pleistocene faunal assemblage from Graniczna Cave (Poland)** · *Radiocarbon* · **55** 🟡 · 13 bone dates plus OSL; focus on collagen preservation and pre-screening criteria. DOI: 10.1017/rdc.2026.10267 · created 19 Oct 2026
- **GeoAI-driven satellite diagnostics of eco-hydrological tipping points and early warning systems for desiccating lake basins** · *Advances in Space Research* · **55** 🟡 · ⚠ *title-based*. DOI: 10.1016/j.asr.2026.10.066 · created 22 Oct 2026
- **On the Stability and Sensitivity of Climate-Informed Flood Frequency Models** · *Geophysical Research Letters* · **53** 🟡 · A semi-parametric framework with a bounded logistic link function (548 US basins) yields more stable flood quantiles than log-linear approaches. DOI: 10.1029/2026gl124039 · created 24 Oct 2026
- **Chronology and Funerary Practices Associated with Antennae Tumuli in Western Sahara** · *African Archaeological Review* · **50** 🟡 · Two AMS dates place a burial at 4417–4100 cal BP — the first absolute chronology for this monument type in the Atlantic Sahara. DOI: 10.1007/s10437-026-09666-8 · created 20 Oct 2026
- **How the River Network Shapes Spatiotemporal Drought Behavior Across Europe** · *Geophysical Research Letters* · **49** 🟢 · Comparing unrouted runoff with routed discharge: the river network increases drought extent in most basins. DOI: 10.1029/2026gl122313 · created 1 Nov 2026

*Not listed or only noted:* The *Science* Perspective "Has the next ice age already come and gone?" is not a primary article. Not included: numerous off-topic hits from topic queries ("early warning", "Turkana", "Rift" in medical, economic or social-science contexts) and pre-Quaternary papers without profile relevance.

## Part B — Funding & calls

**ICDP — AGU 2026 travel grants for early-career researchers — deadline 8 Nov 2026.** Several grants of US$300 for presenting ICDP-related work at the AGU Fall Meeting (CV + accepted abstract; as of previous issue). Source: PAGES Newsletter October 2026.

**ERC Proof of Concept 2027 — opens 12 Nov 2026**, first cut-off 10 Mar 2027 (per a summary of Work Programme 2027; check the ERC site before planning). Source: Hezelburcht — ERC Work Programme 2027.

**Horizon Europe — "Palaeoclimate science for a better understanding of Earth system dynamics" (HORIZON-CL5-2027-01-D1-08).** Opens **17 Nov 2026**, deadline **4 Mar 2027**, Research and Innovation Action, about €14 million for roughly two projects (as of previous issue). Source: Topic page.

**ERC Consolidator Grant 2027 — deadline 12 Jan 2027.** Usually 7–12 years after the PhD. Source: ERC — Consolidator Grant.

**ICDP — 2027 proposal cycle.** Recurring **15 January** deadline for pre-, workshop, full and fast-track proposals; fast-track letters of intent, per ICDP, "by early November at the latest" — i.e., now. The 2027 date is not formally confirmed. Source: ICDP — Proposal Preparation.

**DFG — Open-access publication funding.** Institutional proposals first due **4 May 2027** (as of previous issue). Source: DFG — Information for Researchers No. 68.

*No new profile-relevant calls with a deadline were found this week; the entries above are carried over from Week 44.*

---

## Part C — Science policy (DE/EU)

**WissZeitVG — public hearing on Wednesday, 4 Nov 2026, 10:30.** The Bundestag research committee hears experts on the government bill "modernising fixed-term employment law in science" (BT-Drs. 21/8281), the Greens' motion "Good work on campus" (21/6345) and Die Linke's motion to abolish the special fixed-term regime (21/8384). Invited per the agenda (as of 27 Oct) are, among others, Amrei Bahr (Univ. Stuttgart), Sonja Bolenius (DGB), Patrick Cramer (Max Planck Society), Andreas Keller (GEW) and Birgitta Wolff (Univ. Wuppertal). The hearing is streamed live. Two written statements and four amendment motions are attached to the bill. Assessment: the hearing opens the phase in which changes become realistic; for third-party-funded postdoc positions, minimum contract durations and their application to grant-funded contracts remain the key issue. Source: Agenda, 33rd session of the research committee (PDF).

**German federal budget 2027 — final budget committee session on 12 Nov, final vote on 27 Nov.** Changes to individual budgets, including BMFTR project funding and DFG allocations, are still possible in the final session; government draft: BMFTR core budget about €22 billion, nominally about +0.7% (as of previous issue). Sources: Bundestag — 2027 budget timeline · Forschung & Lehre — Only a small increase for the BMFTR core budget.

**EU budget 2027 — Council/Parliament negotiations, agreement deadline around 16 Nov.** The Council had proposed €12.6 billion for Horizon Europe (−€231.5 million versus the Commission's €12.8 billion draft, including −€5.5 million for the ERC); Parliament voted on 21 Oct for an increase (per a report, +€63 million for Horizon Europe over the Commission). Without agreement by the deadline, the Commission must table a new draft. The 2028–2034 MFF (FP10/

ERC from 2028) remains open. Sources: Science|Business — Council proposes €231m cut · Brussels Times (21 Oct 2026).

## Part D — Community & dates

- **WissZeitVG hearing** in the Bundestag — **4 Nov 2026**, 10:30 (livestream).
- **ICDP AGU 2026 travel grants** — deadline **8 Nov 2026**.
- **ERC Proof of Concept** — opens **12 Nov 2026**.
- **German federal budget 2027** — final committee session **12 Nov**, final vote **27 Nov 2026**.
- **EU budget 2027** — Council/Parliament agreement deadline around **16 Nov 2026**.
- **Horizon CL5 palaeoclimate call** — opens **17 Nov 2026**, deadline **4 Mar 2027**.
- **AGU Fall Meeting 2026** — **7–11 Dec 2026**, San Francisco. Source: AGU.
- **ERC Consolidator Grant 2027** — deadline **12 Jan 2027**.
- **ICDP** — proposal deadline expected **15 Jan 2027**.
- **EGU General Assembly 2027** — **4–9 Apr 2027**, Vienna; abstract submission open since 28 Oct 2026 (deadline usually mid-January, not verified). Source: PAGES Newsletter October 2026.
- **PAGES dates 2027** — PALSEA-Next **24–26 Jun 2027** (Dartmouth College); **PMIP conference 30 Aug – 3 Sep 2027** (Rio de Janeiro). Source: PAGES Newsletter October 2026.

## Methods & sources

Paper screening follows the fixed criteria (topical fit, methodological fit, impact/venue, connectivity, recency) and excludes DOIs already listed in a previous issue. Funding, policy and dates are based on the sources linked in Parts B to D.

**How this issue was produced.** Direct network access from the computing environment was blocked. The **Crossref REST API** (ISSN queries for about 30 journals, including QSR, Climate of the Past, Quat. Sci. Adv., GPC, EPSL, Boreas, JQS, Quat. Res., P³, Nature, Science, PNAS, Nature Commun., Nature Geosci., Sci. Adv., Commun. Earth Environ., GRL, G-cubed, Paleoceanogr. Paleoclimatol., Quat. Int., The Holocene, J. African Earth Sci., J. Human Evol., Radiocarbon; Copernicus prefix including EGUsphere; 16 topic queries; filter `from-created-date:2026-10-19`) and **OpenAlex** (abstracts by DOI from the `abstract_inverted_index`) were queried through the integrated browser interface (`fetch()`). For several very recently registered Elsevier titles no abstract was available yet.

**What this means for coverage.** Where neither OpenAlex nor Crossref provides an abstract, the entry is marked ⚠ *title-based*; no findings beyond the title were invented. Dates are Crossref `created` dates (registration), which may differ from the online publication date by a few days. Topic queries return a lot of off-topic noise; pre-selection used keywords followed by manual review, so individual relevant papers outside the core journals may be missing. Part B received only limited new research this week; carried-over details are marked "as of previous issue".

*GeoNews is an automatically generated digest published by Dr. Markus L. Fischer, Palaeoclimate Dynamics group, University of Potsdam. The thematic profile follows his research interests; selection is made by a rule-based workflow, not an editorial team. The digest is an independent*

*literature overview with no affiliation to the publishers or funders mentioned and does not constitute a recommendation or advice. Entries may contain errors; when in doubt, the linked original source prevails. Corrections and comments are welcome.*