PathoPhys Online Tests (2)

× Untitled Notebook... × Untitled Notebook... × VAA11_II

high hypoxia
high hyperventilation during artificial lung ventilation

The causes of respiratory alkalosis are:

Select one or more options:

- [x] a. high hypoxia
- [ ] b narcotic poisoning
- [ ] cdamage to the central nervous system, which leads to attenuation respiratory center
- [x] d. hyperventilation during artificial lung ventilation

01:37 28%
Google Lens
https://drive.google.co...
Czech → English

8:17 a.m

moodle.upol.cz

secondary lactose intolerance

Task 5

disorder of bile and pancreatic secretion

Malabsorption syndrome occurs most often (2) as a consequence

Select one or more options:

- [x] a. disorders of bile secretion
- [x] b. disorders of pancreatic secretion
- [ ] c. reduced secretion of gastric juice
- [ ] d. decreased secretion of saliva
- [ ] and diabetes insipidus

Task 6

Choose the correct statement(s) about SIADH

Select one or more options:

- [ ]

Translated text

16:42 51%
Google Lens
https://dr...!
Czech → English

02.CZ 21:57 14%

Done

Task 4

hypertonic hyperhydration; Conn's Cushing

Select which state(s). may lead to hypertonic hyperhydration:

Select one or more options:

- [ ] a. Addison's disease
- [x] b. SIADH
- [✓] c. Conn's syndrome
- [✓] d. Cushing's syndrome
- [ ] e. Drinking large amounts of tap water

Translated text

Send to Translate Home

Select all

Did you find these results useful?

Yes No

18:16 36%
Google Lens
https://drive.google.co...
Czech → English

Ulaha 6

Cloche 1:

Task 9

Bodů 0,50 / 1,00

Losses of water and solutes during excessive sweating lead to

Select one or more options:

- a. hypernatremii
- b. hypertonic dehydration
- c. chypotonic dehydration
- Ed isotonic dehydration
- e. volume increase intracellular fluid due to osmotic shifts of water between extracellular and intracellular compartment

hypernatremia

The correct answers are:
hypertonic dehydration,
hypernatremii

Translated text

Send to Translate Home

Select all

Did you find these results useful?

Yes No

01:42 27%
Google Lens
https://drive.google.co...
Czech → English

8:17 a.m

moodle.upol.cz

Task 6

Choose the correct statement(s) about SIADH

Select one or more options:

- [ ] a. Severe edema of the lower extremities is one of the most important complications
- [x] lung cancer can lead to hypersekreci ADH
- [x] In case of incorrect treatment, pontinitis can occur demyelination
- [ ] d. It is associated with a decreased level ANP
- [x] May cause hyponatraemia

Task 7

In an acutely admitted patient with repeated vomiting during an attack lasting several days acute pancreatitis, these internal disorders can be found during admission to the hospital

environment

Translated text

15:19 69%
Google Lens
https://lens.google.co...
Czech → English



IMG_1214.JPG
IMG_1218.JPG
Translated text

01:53 43%
Google Lens
https://drive.google.co...
Czech → English

4:05 p.m 83%

Task 8

Constipation

Constipation

Select one or more options:

- [ ] a. can occur with hyponatremia
- [ ] b. arises in hyperthyroidism
- [x] can occur during dehydration
- [x] is a negative factor that may contribute to portosystemic encephalopathy in advanced liver cirrhosis
- [x] risk factors include older age

Translated text

17:05 45%
Google Lens
https://drive.google.co...
Czech → English

Toko S40 1094% 1441

patients' blood osmolarity
is high

With the biggest
chances are the patient
has an elevated level of ketone
bodies in the plasma ✓

With the biggest
chances are the
patient is dehydrated
due to osmotic
diuresis ✓

Most
likely, the patient
has a normal or
decreased anion
gap

The correct answers are: S
most likely has
patient tachycardia as a result
dehydration and activation
sympathetic nervous system
system, 5 max
the patient has probabilities
dehydration as a result
osmotic diuresis. It
most likely is
osmolarity of patients' blood
high, with the largest
the patient has probabilities
elevated level of ketone bodies in
plasma

Dream -U... 3.JPG
Dream U... 6.JPG
Translated text

16:33 53%
Google Lens
https://drive.google.co...
Czech → English

8:51 am

MESSENGER now

David Gronych

VŠEO Group

And from 10:45 I hope they don't teach anymore 😂😂



Select one or more options:

- [ ] a. Crohn's disease
- [ ] b. Appendicitis
- [ ] c. Meckel's diverticulum
- [ ] d. Rectal fistula
- [ ] e. Hirschsprung's disease

Job 9

Task 2

Translated text

01:34 47%

https://drive.google.co...

Google Lens

Czech → English

A combination of hypokalemia and hyponatremia is found in

hypokalemia + hyponatremia

Select one or more options:

- [ ] a. failure of the adrenal cortex
- [ ] b. pure water dehydration
- [ ] c. Diabetes insipidus
- [x] d. overproduction syndrome antidiuretic hormone
- [ ] E. primary hyperaldosteronismu

overproduction ADH

Translated text

Send to Translate Home

Select all

Did you find these results useful?

Yes No

00:26 45%
Google Lens
https://drive.google.co...!
Czech → English

8:53 am LTE

Task 2

A part of the compartment of body fluids is also a transcellular component, which

Select one or more options:

- [x] it is formed mainly by secretions in the lumen of the digestive tract and cerebrospinal fluid
- [ ] b. make up about 1-1.5% intracellular fluids c.
- [ ] are formed only in pathological conditions (e.g. ascites)
- [ ] d. is part of all cells

Task 3

moodle.upol.cz

Translated text

16:03 61%

Google Lens

https://lens.google.co...

Czech → English

IMG_1205.JPG

IMG_1206.JPG

Task 4

Select the correct statement(s) about hypokalemia:

Select one or more options:

A. A tall spiked T wave is prominent on the EKG

It develops in patients with primary hyperaldosteronism

Can even cause paralytic ileus

swelling

E. May be associated with metabolic acidosis

The correct answers are: Can cause up to paralytic ileus, Can v
patients with primary hyperaldosteronism

IMG_1212.JPG

IMG_1213.JPG

Task 2

Task 7

IMG_1214.JPG

IMG_1218.JPG

Translated text

01:38 46%
Google Lens
https://drive.google.co...
Czech → English

Task 6

Task 1

Pathological changes in the esophagus during gastroesophageal reflux are caused

Select one or more options

a. by slowing down and deepening peristaltic waves jicnu

b. a rise in pH in the stomach and a lack of proteinases

the presence of HCl and of aspartate proteinases in the intestine

d by the presence of genitives diverticula

Translated text

Send to Translate Home

Select all

Did you find these results useful?

Yes No

16:57 48%
Google Lens
https://drive.google.co...
Czech → English

Task 10

Point 0.00/1.00

A 52-year-old patient, a chronic alcoholic, is admitted to the hospital with liver cirrhosis, acute signs of alcohol intoxication and vomiting. His albumin level is 10 g/L (N = 35-50 g/L), pH 7.38, Cl-90 mmol/L, PaCO2-5.3 kPa (40 mm Hg). Choose the 1 most appropriate statement about the patient:

Select one or more options:

- With the biggest chances are the patient has a combination of metabolic acidosis and metabolic alkalosis
- Most likely, the patient has a combination of metabolic acidosis and respiratory alkalosis
- ✓ Based on available information, no conclusion can be made about acid-base balance patient X
- d. With the largest

Translated text

23:59 64%
https://dri...!
Google Lens
Czech → English



Translated text
Send to Translate Home
Select all
Did you find these results useful?
Yes
No

14:11 37%
https://drive.google.co...
Google Lens
Czech → English

Task 8

Crohn's disease

Select one or more options:

- [ ] and. It is an acute inflammatory disease of the colon
- [x] D. Extraintestinal complications include joint inflammation, canker sores
- [x] C. Granulomatous inflammation penetrates the entire wall
- [x] d. It most often affects the small intestine (ileitis)
- [ ] E. It usually starts with involvement of the anus

Task 2

Translated text

16:36 52%

Google Lens

Czech → English



IMG_1428.JPG IMG_1429.JPG

IMG_3234.JPG IMG_3235.jpg

Translated text

16:36 52%
Google Lens
Czech → English

IMG_1428.JPG IMG_1429.JPG

In the images we see pathology in the stomach area. What are the risk factors for this condition?



Select one or more options:

- [x] Nesteroidní antirevmatika
- [x] Helicobacter pylori infection
- [x] Mental stress
- [ ] d. Secretion of endogenous prostaglandins thicky E
- [x] Duodenogastricky reflux

IMG_3234.JPG IMG_3235.jpg

Translated text

23:11 61%
Google Lens
https://drive.google.co...
Czech → English

O2 - CZ|O2 - SK 10147% 13:57

Task 4

Hiatal hernia

Select one or more options:

- [ ] a. It often leads to cirrhosis of the liver
- [x] It can manifest heartburn
- [x] It can lead to development strangulation hernias or k obturation
- [ ] d. It is often manifested diarrhea
- [ ] e. It is one of the important causes of celiac disease

Task 5

It does NOT belong to alkalizing agents

Translated text

01:52 24%

https://drive.google.co...

Google Lens

Czech → English

8:17 a.m

moodle.upol.cz

Task 7

In an acutely admitted patient with repeated vomiting during a several-day attack of acute pancreatitis, the following disorders of the internal environment

Select one or more options:

- [ ] and metabolic alkalosis with hypercapnil, polyuria, hyperkalemia
- [x] combination of metabolic alkalosis and acidosis, dehydration, reduced moti chloride excretion
- [ ] c. pure metabolic alkalosis with complete compensation, hypochloremia, hyperchloruria, and hyperkalemia
- [ ] d. pure metabolic acidosis with hypocapnia, dehydration and hypokalemia

Translated text

Send to Translate Home

Select all

Did you find these results useful?

Yes No

01:26 31%

Google Lens

https://lens.google.co...

Czech → English



Translated text

Send to Translate Home

Select all

Did you find these results useful?

Yes No

17:59 40%
Google Lens
https://drive.google.co...
Czech → English

Ulaha 6

Clove 1

Task 4 Point) 1/20/1.02

For the classification of edema, the following characteristics are not used:

Choose one or more options:

- [ ] Pitting and non-pitting
- [x] Acquired and inherited ✓
- [ ] Primaries and secondary
- Tymphetium
- [x] Hyperosmolar hypoosmolar ✓✓
- [ ] Local and generalized

The correct answers are:
Hyperosmolar and hypoosmolar,
Acquired and inherited

Translated text
Send to Translate Home
Select all
Did you find these results useful?
Yes No

15:57 63%
Google Lens
https://drive.google.com...!
Czech → English

58% 9:26

Task 4

Children with kwashiorkor are characterized by:

Select one or more options:

- [x] Zvětšenými játry s tukovými Infiltrates
- [ ] b. Lack of dietary carbohydrates
- [x] A protruding belly
- [x] Low resistance to infections
- [x] Pitting edema with swelling of ankles and feet

The correct answers are: Protruding abdomen, Low resistance to infections, An enlarged liver with fatty infiltrates, Pitting edema with swelling of ankles and feet

Task 5

Translated text

16:31 54%

Google Lens

https://drive.google.co...

Czech → English

.02.CZ 9:25 84%

moodle.upol.cz

615

Task 3

Malabsorption may be present

Select one or more options:

a. reduced resorption of water and salt

b. increased water resorption a salt

c. insufficient breakdown of fats

d. discounted intraluminal osmotic pressure in the alimentary canal

Translated text

Send to Translate Home

Select all

Did you find these results useful?

Yes No

16:36 53%

Google Lens

Cristiano ... Piqu...

Czech → English

15:41 O 63%

moodle.upol.cz/mod/quiz/review.

Task 1

A non-physiologically high value of base excess (BE) can be associated with

Select one or more options:

- [x] metabolic alkalosis without compensation
- [ ] b. no answer is correct
- [ ] Crespiratory alkalosis without metabolic compensation
- [ ] d. respiratory acidosis without metabolic compensation
- [ ] e respiratory alkalosis with renal compensation

Task 2

Translated text

Send to Translate Home

Select all

Did you find these results useful?

Yes No

15:13 71%
Google Lens
https://drive.google.co...
Czech → English

4:22 p.m

malignant lymphoma

Task 3

Lactase deficiency

Select one or more options:

- [ ] causes intolerance to milk and some dairy products
- [ ] b. causes Crohn's disease disease
- [ ] some patients mail diarrhea and pain after milk and dairy products, some have minimal difficulties
- [ ] d. causes intolerance animal fats
- [ ] e. causes intolerance to foods containing gluten

Translated text

23:12 63%

Google Lens

https://drive.google.co...

Czech → English

moodle.upol.cz/mod/qu

Task 5

It does NOT belong to alkalizing agents

Select one or more options:

- [ ] and sodium lactate
- [x] NH4Cl
- [ ] bicarbonate
- [ ] sodium citrate

Task 6

Write down the question number according to the question list and the answer to that question. You do not need to rewrite the wording of the question.

Translated text

01:54 24%
Google Lens
https://drive.google.co...
Czech → English

8:17 a.m

moodle.upol.cz

d. pure metabolic acidosis with hypocapnil, dehydration and hypokalemia

Task 8

Causes of metabolic alkalosis include

Select one or more options:

and the effect of inhibitors karboanhydrázy

b. hyperkalémie

primary hyperaldosteronism

d. Vomiting

Task 9

Translated text

Send to Translate Home

Select all

Did you find these results useful?

Yes No

01:30 30%
Google Lens
https://drive.google.co...
Czech → English

8:17 a.m
moodle.upol.cz

Task 4

Celiac sprue (celiac disease) malignant lymphoma

Select one or more options:

represents an increased risk of malignant lymphoma

b. is often complicated by the formation of adhesions between individual ones parts of the gastrointestinal tract tract is

c. characterized by malabsorption gluten

may be accompanied by secondary lactose intolerance

Task 5

Translated text
Send to Translate Home
Select all
Did you find these results useful?
Yes No

17:06 45%

Google Lens

https://drive.google.co...

Czech → English

Task 6

The anion gap is calculated from plasma concentration values

No Cl HCO_3

Select one or more options:

- [[Na+]+[K+]-[HCO3-]
- b[Na+] + [K+] + [Ca] + [HCO3-]-[Cl]
- [Na+] + [K+] - [Cl-]
- [Na]+[K+]-[HCO3-]
- [Na+] - [Cl] - [HCO3-] ✓

The correct answer is: [Na+ - [Cl-] - [HCO3-]

Dream -U... 3.JPG

Dream Hu... 6.JPG

Translated text

10:53 89%
Google Lens
https: //d...e.com/d ...
Czech → English
02.cz 14:37 39%
moodle.upol.cz

Job 2

Which factor / y increases / and feeling hunger:

Select one or more options:

- [ ] a. leptin
- [ ] b. Insulin
- [x] c. Ghrelin
- [ ] d. interleukin 1
- [x] e. neuropeptide y

Translated text
Offline · Some features are not available
Send to Translate Home
Select all
Sh
Did you find these results useful? Yes No

10:53 89%
Google Lens
https: //d...e.com/d ...
Czech → English
IL O2.cz 14:37 意 39% 1
moodle.upol.cz

Job 1

Constipation can be caused

Select one or more options:

- By adopting food with a nise content of fiber
- 0 B bacterial infection in the colon
- c. By slowing the intestinal passage caused by hyperthyroidism
- d. tumor pressure from the
- e. surrounding defectant reflection

Translated text
Offline · Some features are not available
Send to Translate Home
Select all
Sh
Did you find these results useful? Yes No

10:53 89%
Google Lens
https: //drive.google.com/d ...
Czech → English

10:42 4G

Task 9

Tissue lactic acidosis cannot be caused

Select one or more options:

- [ ] a. by the lack of oxygen in tissues from circulatory concerning
- [ ] b. Excessive muscle work
- [ ] c. bounds of lactate metabolism in liver
- [ ] d. The lack of oxygen at respiratory diseases
- [x] e. Hypoglycemia

To complete the browser

Translated text
Offline · Some features are not available
Send to Translate Home
Select all
Sh
Did you find these results useful? Yes No

10:52 89%
Google Lens
https: //d...g...le.com/d ...
Czech → English

10:42 4G

Job 8

In the picture Vidime pathology in the transition of the esophagus / stomach. What pathology, or. the type is?



Select one or more options:

- [ ] and Jikn diverticl
- [ ] b. Ahalasia
- [ ] c. Savelní ulcer
- [ ] d. Paraezophageal Hernie
- [x] e. Slide

moodle.upol.cz

Translated text
Offline · Some features are not available

10:52 89%

Google Lens

https: //d...g...le.com/d ...

Czech → English

10:42 l 4g

Job 7

When gastroesophageal reflux can

dozeft

Select one or more options:

- [ ] A. To shorten the relaxation time

cardie

- [x] To inflammatory changes

them

- [ ] To develop Barrett's esophagus
- [x] To regurgitate acid to

lung

- [ ] To the development of adenocarcinoma

esopl-agus

Translated text

Offline · Some features are not available

Send to Translate Home | Select all | Sh

Did you find these results useful? Yes No

10:52 89%
Google Lens
https: //d...e.com/d ...
Czech → English

Job 6

Young diabetics unconscious, treated insulin (found by card

nooпe

Plasma potassium

Select one or more options:

- increases with a drop drop
- b. is reduced at hyperaldosteronism
- CJs reduced in the terminal stage of renal insufficiency
- d. decreases with the drop pH
- e. Is usually 10-15 mmol

Translated text
Offline · Some features are not available
Send to Translate Home
Select all
Sh
Did you find these results useful? Yes No

10:52 89%
Google Lens
https: //d...e.com/d ...
Czech → English

8:17:17

moodle.upol.cz

Job 7

Acutely accepted patient with repeated vomiting in a multi-day acute pancreatitis seizure can be found in the hospital to find these disorders

Select one or more options:

- a. Metabolic alkalosis with hypercapnia, polyurius and hyperkalumi
- b. combination of metabolic alkalosis and acidosis, dehydration, reduced chloride secretion
- c. pure metabolic alkalosis with complete compensation, hypochloris, hyperchloruria and hyperkalumi
- d. Pure metabolic acidosis with hypokapenia, dehydration and hypokalemia

Úloha 8

Translated text
Offline · Some features are not available

10:52 89%
Google Lens
https: //d...e.com/d ...
Czech → English

8:17:17

moodle.upol.cz

Job 7

Acutely accepted patient with repeated vomiting in a multi-day acute pancreatitis seizure can be found in the hospital to find these disorders

Select one or more options:

- a. Metabolic alkalosis with hypercapnia, polyurius and hyperkalumi
- b. combination of metabolic alkalosis and acidosis, dehydration, reduced chloride secretion
- c. pure metabolic alkalosis with complete compensation, hypochloris, hyperchloruria and hyperkalumi
- d. Pure metabolic acidosis with hypokapenia, dehydration and hypokalemia

Úloha 8

Translated text
Offline · Some features are not available

10:52 89%
Google Lens
https: //drive.google.com/d ...
Czech → English

8:17:17

moodle.upol.cz

d. Pure metabolic acidosis with hypokapnia, dehydration and hypokalemia

Role 8

Among the states of metabolic alkalosis

Select one or more options:

a. the effect of inhibitors carboxyydrase

b. Hyperkalaemia

c. Primary hyperaldosteronism

d. vomited

Task 9

Primary metabolic acidosis may occur

Select one or more options:

a. [redacted]

Translated text
Offline · Some features are not available

10:52 89%
Google Lens
https: //drive.google.com/d ...
Czech → English
8:17:17
moodle.upol.cz

d. Pure metabolic acidosis with hypokapnia, dehydration and hypokalemia

Role 8

Among the states of metabolic alkalosis

Select one or more options:

- a. the effect of inhibitors carboxyydrase
- b. Hyperkalaemia
- c. Primary hyperaldosteronism
- d. vomited

Task 9

Primary metabolic acidosis may occur

Select one or more options:

- a. [redacted]

Translated text
Offline · Some features are not available

10:52 89%
Google Lens
https: //drive.google.com/d ...
Czech → English

8:17:17

moodle.upol.cz

secondary lactose intolerance

Job 5

Malabsorbent syndrome is most commonly formed (2) as a result

Select one or more options:

a. bile secretion disorders

b. Pancreatic secretion disorders

c Reduced secretion of the gastric wage

d. Reduced secretion of saliva

Diabetes Insipidus

Job 6

Translated text
Offline · Some features are not available
Send to Translate Home
Select all
Did you find these results useful? Yes No

15:59 62%
Google Lens
https://drive.google.co...
Czech → English

10:40 a.m.

metabolic acidosis

Task 5

During a visit to the pediatrician, a fourteen-year-old girl is diagnosed with anorexia nervosa. Select the symptoms that helped the doctor make this diagnosis:

Select one or more options:

- [ ] a. Body height above 95th percentile
- [x] b. Absence of periods for more than 3 months after having had regular periods since age 12 to 13
- [ ] GBMI 21.5 kg/m2
- [x] Body weight below 5th percentile
- [ ] Ascites caused hypoproteinemii

Task 6

Translated text

00:16 48%
Google Lens
https://drive.google.co...:
Czech → English
58%8:20
moodle.upol.cz/mod/qu



Select one or more options:

- Ulcerative colitis
- b. Hirschsprungova disease
- C. Appendicitis
- d. Meckel's diverticulum
- Crohn's disease

Translated text

01:59 22%

Google Lens

Czech → English



Translated text

Send to Translate Home

Select all

Did you find these results useful?

Yes No

15:21 68%
Google Lens
https://lens.google.co...
Czech → English

T-Mobile CZ 8:24 @ 100%
moodle.upol.cz

Task 7

Úbytek

Sodium depletion is regularly accompanied

Select one or more options:

- [ ] and colloidal reduction osmotic pressure plasma
- [x] b. by reducing the volume extracellular fluid
- [ ] and low values hematocrit
- [ ] d. by increasing skin turgor

Task 8

IMG_1214.JPG IMG_1218.JPG
Translated text

23:13 63%
Google Lens
https://drive.google.co...
Czech → English

O2 - CZ|O2 - SK 10147% 13:58

Task 8

Select the important cause(s) of hypernatremia:

Select one or more options:

- [x] A. Profuse sweating in patients with fever
- [ ] B. Sterile wound on forearm without marks SIRS
- [ ] C. SIADH
- [ ] D. Type 2 diabetes mellitus, fully compensated state

Task 9

Select the processes that are common in the pathogenesis of gout:

Translated text

23:44 49%
Google Lens
https://drive.google.co...:
Czech → English

58% 8:19 am

Task 5

long term starvation = metabolic acidosis

Which acid-base disorder balance develops primarily in patients with long-term simple starvation

Select one or more options:

- [ ] A. Metabolic alkalosis
- [ ] B. Respiratory acidosis
- [ ] C. Respiratory alkalosis
- [ ] D. Metabolic acidosis
- [ ] E. Combined metabolic and respiratory acidosis

Task 6

Translated text

00:26 45%
Google Lens
https://drive.google.co...!
Czech → English

8:53 am LTE

Task 1

Choose the correct statement(s) about stress starvation

Select one or more options:

- [ ] a. Ketone bodies are often elevated
- [ ] b. The insulin level is usually reduced
- [x] c. Glucose levels often rise
- [x] d. It is characterized by intense protein catabolism
- [ ] e. Severe emotional stress (e.g. death of a partner) is one of the most important causes

AA moodle.upol.cz
Translated text

16:55 48%
Google Lens
https://drive.google.co...
Czech → English

T-Mobile CZ 8:24 am 100%

moodle.upol.cz

B12 → hyperhomocysteinemia

Choose the correct statement(s) about the vitamin.

B12:

Choose jeuna new vice

- [ ] a. The recommended daily intake of B12 is 70-150 mg for adults
- [ ] b. B12 is mostly reabsorbed in the duodenum
- [x] c Vitamin B12 deficiency can lead to macrocytic anemia and demyelination of nerves
- [x] d Vitamin B12 deficiency leads to hyperhomocysteinemia with an increased risk of developing endothelial damage
- [x] e Gastric resection leads to B12 reabsorption disorders

Translated text

23:00 52%
Google Lens
https://drive.google.co...
Czech → English

58% 8:19 am

Task 4

Pure water hyperhydration leads in the acute stage to

Select one or more options:

- [ ] a. increase in the volume of erythrocytes
- [ ] b. leads to a reduction plasma protein concentration
- [ ] c. hypernatrémii
- [ ] d. hyperchlorémii
- [ ] e. reduction in the volume of erythrocytes

Translated text
Send to Translate Home
Select all
Did you find these results useful?
Yes No

01:54 43%

https://drive.google.co...

Google Lens

Czech → English

Task 6

Task 1

IMG 3198.JPG

IMG 3199.JPG

In case of salicylate poisoning, it is developing

apply acid-base disorders

Select one or more options:

a. formation of ketone bodies- metabolic acidosis

b irritation of the respiratory center alkalosis

c diarrhea→metabolic alkalosis

d. metabolic acidosis

e depression of the respiratory center, respiratory acidosis

Translated text

Send to Translate Home

Select all

Did you find these results useful?

Yes No

13:54 23%
Google Lens
https://lens.google.co...
Czech → English

Task 4

Secondary hyperaldosteronism

Select one or more options:

a. decreases the ratio of Na/K concentration in urine

b. leads to metabolic acidosis

X c. increases the ratio of Na/K concentration in urine

d. leads to metabolic alkalosis

e. can lead to an increase arterial blood pressure

IMG_1214.JPG
IMG_1218.JPG
Translated text

16:31 54%
Google Lens
https://drive.google.co...
Czech → English

They are like mature

02.CZ 9:25 84%
moodle.upol.cz

736

Task 4

Renal compensatory response to metabolic acidosis

Select one or more options:

- leads to a reduction of the so-called titratable acidity of urine
- b. reaches its maximum within 6-12 hours
- reaches its maximum up to 3-5 days
- d. leads to an increase in the so-called titratable acidity of urine
- it reaches its maximum within 12-24 hours

IMG_1214.JPG IMG_1218.JPG
Translated text

17:49 43%

Google Lens

https://drive.google.co...

Czech → English

Ulaha 6

Clove 1

Task 4

Bodů 0,50 / 1,00

Select conditions associated with hypercholesterolemia

Select one or more options:

- Hepatocellular carcinoma with cachexia
- Anorexia nervosa✓
- Type diabetes mellitus
- Nephrotic syndrome✓
- Hypothyroidism

The correct answers are: Type II diabetes mellitus, Anorexia nervosa, Nephrotic syndrome, Hypothyroidism

Dream -U... 3.JPG

Dream Hu... 6.JPG

Translated text

d. is influenced by the current state of renal function

Task 8

Mortality risk in patients with mechanical ileum represents

Select one or more options:

- a. weight loss
- b. resorption of bacterial toxins
- c. bile secretion disorder
- d. hyperhydration
- e. dehydration

16:59 47%
Google Lens
https://drive.google.co...
Czech → English

@1094% 1441

Task 8

0.50 / 1.00

In a 12-year-old boy, developed decompensated type 1 diabetes mellitus due to insufficient insulin dose His glucose level is 20.9 mmol/L, blood pH is 7.1, the patient is severely dehydrated

Select all correct statements about the patient:

Select one or more options:

With the biggest the patient is likely to have tachycardia due to dehydration and activation of the sympathetic nervous system

With the biggest probability is the osmolarity of the patients' blood high

With the biggest chances are the patient has an increased level of ketone bodies in the plasma ✓✓

With the biggest chances are the patient has dehydration due to osmosis

Translated text

16:09 59%

Google Lens

Czech → English

IMG_1205.JPG

IMG_1206.JPG

Task 4

Vybe typical p

Values of pH = 7.25, pCO2 = 4.0 kPa (30 mm Hg), HCO3-15 mmol/l belong to a patient with

Select one or more options:

Chronic metabolic acidosis with activated compensatory mechanisms

B. Chronic metabolic alkalosis with activated compensatory mechanisms

C. Acute respiratory acidosis

D. Combined respiratory and metabolic acidosis

E Alutoi respiratory alkalo

The correct answer is: Chronic metabolic acidosis with activated compensatory n

IMG_1212.JPG

IMG_1213.JPG

Task 2

Task 7

IMG_1214.JPG

IMG_1218.JPG

Translated text

16:39 52%
Google Lens
https://drive.google.co...
Czech → English

3:41 p.m 1963%

d. is an intestinal manifestation of
congenital progressive myodystrophy

Task 4

At the beginning of hypotonic dehydration

Select one or more options:

- a. there is an osmotic movement of water from the interstitium into the cells
- with the volume of intracellular fluid increases
- c. the level decreases antidiuretic hormone
- d. the volume of intracellular fluid decreases

Translated text

15:58 62%

Google Lens

Czech → English

k 4

kwashiorkor are characterized by:

e or more options:

Zvětšenými játry s tukovými

Infiltrates

Lack of dietary carbohydrates

A protruding belly

Low resistance to

Translated text

Send to Translate Home

Select all

Did you find these results useful?

Yes

No

16:06 60%
Google Lens
Czech → English

IMG_1205.JPG

IMG_1206.JPG

Task 4

Task 13

Hyperkapnie

Select one or more options:

- Such Hyperkaphia leads to vasodilatation of the CNS tube as the brain flows
- It can develop in patients with COPD
- With a high increase in PaCO2, the brain is inhibited, the patient may develop a hypercapnic coma It
- is a consequence of Kussmaul breathing
- by increasing PaCO2, the respiratory center is stimulated

The correct answers are: With a slight increase in PaCO2, the respiratory center is stimulated, Telka hypercapnia leads to and increases the risk of brain swelling. With a high increase in PaCO2, the brain is inhibited and the patient may develop h
It can develop in patients with severe COPD

IMG_1212.JPG

IMG_1213.JPG

Task 2

Task 7

IMG_1214.JPG

IMG_1218.JPG

Translated text

10:52 89%
Google Lens
https: //drive.google.com/d ...
Czech → English
8:17:17
moodle.upol.cz

Job 2

Choose an important / important cause / states of hypernatremia:

Select one or more options:

- [ ] and lack of glucocorticoids
- [ ] Increased sweating in patients with fever
- [x] Diabetes Insipidus
- [ ] Increased Aldosterone production
- [ ] e. Siadh

Task 3

Translated text
Offline · Some features are not available
Send to Translate Home
Select all
Did you find these results useful? Yes No

10:52 89%
Google Lens
https: //d...e.com/d ...
Czech → English
39.jpg
0 國 40.jpg

The organism better tolerate the pH in the area

Choose one or more options

- and acidemia because dissociation IVKA axyhemoglobin is moved to the left
- b. ATVALEEMY, PROTOTE dissociation ivka axyhemoglobin is moved to the right
- c. acidemia because dissociation IVKA oxyhemoglobin is moved to the right
- d Alkatémie because dissociation oxyhemoglobin cluster moves the left

Translated text
Offline · Some features are not available
Send to Translate Home
Select all
Did you find these results useful? Yes No

10:51 89%
Google Lens
https: //d...google.com/d ...
Czech → English
39.jpg
0 國 40.jpg
Hyperchloremic acidosis
Job 8

In hyperosmolar dehydration

Select one or more options:

- Aje a sense of mensi masculine than u isoosmoles dehydration
- It is a lot of lost water than the Low of Lost Solut
- c. Femble of lost water less than the amount of lost Solutd
- D is the amount of lost water the same as the amount of lost Soluto

Ulcus bulbi duodeni is a multifactorial the disease. But it turned out that it was
In hyperosmolar dehydration
Choose one or more options

Translated text
Offline · Some features are not available
Send to Translate Home
Select all
Sh
Did you find these results useful? Yes No

10:51 89%
Google Lens
https: //d...ge.com/d ...
Czech → English
39.jpg
0 國 40.jpg

Celiacal Sprue (celiacia)

Select one or more options:

- [x] is induced by gliadinin intolerance
- [x] Prevalence in CR is approximately 1: 250-300 throughout the age spectrum
- [ ] The basis of the air is a gluten-free diet only in terms of interest
- [ ] DJe caused by lactose intolerance
- [x] Immunic factors are employed in Etiopathogenesis of disorder

It is an immunic unplaceabody so neviem

Choose one or more options

Translated text
Offline · Some features are not available
Send to Translate Home
Select all
Did you find these results useful? Yes No

10:51 89%
Google Lens
https: //d...e.com/d ...
Czech → English

Sodium retention j
1st - 2nd hyperaldosteronism

19.JPEG 20.jpg

Between main puffs of sodic retention (2)
Select one or more options:

a. Increased blood flow by kidney

b. Undetty cortex function

c. Primary hyperaldosteronism

d. Secondary hyperaldosteronism

e. the rise of oncotic plasma pressure

23.jpg 24.jpg

Translated text
Offline · Some features are not available
Send to Translate Home
Select all
Did you find these results useful? Yes No

10:51 89%
Google Lens
https: //d...e.com/d ...
Czech → English
19.JPEG
0 0jpg
0

For Crohn's disease paid

Choose one or more options

- and most often affected
  Sigmoidum and rectum
- b. The most frequently affected
  terminin ileum and reside
  the deal
- c. H. pylori is the most important
  etiological factor
- d inflammation with breadth continuously
  Stisna is disabled diffusive

23.jpg
0 四 24.jpg
Translated text
Offline · Some features are not available
Send to Translate Home
Select all
Sh
Did you find these results useful? Yes No

10:51 90%
Google Lens
https: //d...g...e.com/d ...
Czech → English
19.JPEG
0 國 20.jpg

In the esophagus Achalasia occurs, even except to select one or more options

55

a. aspiration pneumonia

b. loss of myenteric ganglium cells in the esophagus wall

c. Development of Barrett's Jikn

d. Disorders relaxing the Dolni Jíno Sphincer

e. esophageal peristaltic disorders

23.jpg
24.jpg
Translated text
Offline · Some features are not available
Send to Translate Home
Select all
Did you find these results useful? Yes No

10:51 90%
https: //d...g.google.com/d ...
Google Lens
Czech → English
Vyerte one or more mo
19.JPEG
0 國 20.jpg

In the esophagus Achalasia occurs, even except to select one or more options

55

a. aspiration pneumonia

b. loss of myenteric ganglium cells in the esophagus wall

c. Development of Barrett's Jikn

d. Disorders relaxing the Dolni Jíno Sphincer

e. esophageal peristaltic disorders

23.jpg
24.jpg
Translated text
Offline · Some features are not available
Send to Translate Home
Select all
Sh
Did you find these results useful?
Yes
No

10:51 90%
https: //d...g...gle.com/d ...
Google Lens
Czech → English

I can't imbue hyperkalaemia

Select one or more options:

and primary hyperaldosteroneism

b. Extensive hemolysis after administration of incompatible blood

Cakutni Failed kidney

d. Acidosis

Translated text
Offline · Some features are not available
Send to Translate Home
Select all
Sh
Did you find these results useful? Yes No

10:51 90%
https: //d...g...gle.com/d ...
Google Lens
Czech → English

I can't imbue hyperkalaemia

Select one or more options:

- and primary hyperaldosteroneism
- b. Extensive hemolysis after administration of incompatible blood
- Cakutni Failed kidney
- d. Acidosis

Translated text
Offline · Some features are not available
Send to Translate Home
Select all
Did you find these results useful? Yes No

17:56 41%
Google Lens
https://drive.google.co...:
Czech → English

Task 8 Points 0.00/1.00

Intermittent fasting can mean that a person:

Choose one or more options: a.

- [ ] Consumes only 1 type of food each week (e.g. apples 1 week, bread 2 weeks, cheese 3 weeks...)
- [ ] Consumes a low-protein diet for 1 day, then a low-fat diet for 1 day, then a low-carb diet for 1 day X
- [x] Consumes food for 6 hours daily and does not consume for 18 hours food
- [x] Consumes food for 8 hours a day and does not consume food for 16 hours
- [ ] Consumes only 1 type of food every day (e.g. 1 day apple, 2 day bread, 3 day cheese...) X

The correct answers are: Consumes food for 8 hours a day and does not consume food for 16 hours, Consumes food for 6 hours a day and does not consume food for 18 hours

Dream -U... 3.JPG
Dream Hu... 6.JPG
Translated text

36 Task 4

Renal compensatory response to metabolic acidosis

Select one or more options:

- a. leads to a reduction of the so-called titratable acidity of urine
- b. reaches its maximum within 6-12 hours
- c. reaches its maximum up to 3-5 days
- d. leads to an increase in the so-called titratable acidity of urine
- e. it reaches its maximum within 12-24 hours

02:11 39%

Google Lens

https://lens.google.co...

Czech → English

Task 6

Task 1

IMG 3198.JPG

IMG 3199.JPG

Protective factors in the pathogenesis of peptic ulcer include:

Select one or more options:

a. žluč

b intact microcirculation in the gastric mucosa

caspartate proteinases

abnormal secretion of prostaglandins

WIMG 3234.JPG

IMG 3235.jpg

Translated text

Send to Translate Home

Select all

Did you find these results useful?

Yes No

14:12 38%
Google Lens
https://drive.google.co...!
Czech → English

4:21 p.m

Test #2 2021: Suddenly...
moodle.upol.cz

Task 1

Select the processes that are common in the pathogenesis of gout:

Select one or more options:

- [ ] a. High level of urea in the blood
- [x] As the gout progresses, tophi appear on the hands, fingers, elbows and ears
- [x] It precipitates in the joint cavity uric acid crystals
- [ ] d. Myocardial infarction is often the initial sign of gout
- [x] Hyperemia and severe pain in the affected joint during an exacerbation

Task 2

Translated text

22:55 51%
Google Lens
https://drive.google.com...
Czech → English

d. is influenced by the current state of renal function

Task 8

Mortality risk in patients with mechanical ileum represents

Select one or more options:

- [ ] a. weight loss
- [x] b. resorption of bacterial toxins
- [ ] c. bile secretion disorder
- [ ] d. hyperhydratace
- [x] e. dehydration

Translated text

16:04 60%

Google Lens

Czech → English

IMG_1205.JPG

IMG_1206.JPG

Metabolic (Reaven) syndrome includes

Select one or more options:

- [ ] A. Proteinuria with hematuria
- [x] B. Impaired glucose tolerance
- [x] C. Lowering HDL cholesterol
- [x] D. Insulin resistance
- [ ] E. Hemolytic anemia

The correct answers are: Insulin resistance

Task 2

Task 7

Translated text

Send to Translate Home

Select all

Did you find these results useful?

Yes No

14:10 37%
Google Lens
https://drive.google.co...
Czech → English

Task 7

Ulcerative changes in the oral cavity

Select one or more options:

- [ ] and do not occur in infectious diseases
- [ ] b. can be manifested in allergic diseases
- [ ] c. occur in states of deficiency
- [ ] d. may develop in patients with Crohn's disease

IMG_1214.JPG
IMG_1218.JPG
Translated text

00:50 39%
Google Lens
https://drive.google.co...!
Czech → English

8:53 am LTE

Task 3

The following can NOT occur
with acid-base imbalances
combination of precipitating factors

Select one or more options:

- [ ] a. respiratory acidosis + respiratory alkalosis
- [ ] b. metabolic acidosis + respiratory alkalosis
- [ ] c metabolic acidosis + metabolic alkalosis
- [ ] d. respiratory acidosis + metabolic acidosis

Task 4

moodle.upol.cz

Translated text

00:20 47%
Google Lens
https://drive.google.co...
Czech → English

58%8:20

moodle.upol.cz/mod/qu

16

Task 9

Severely malnourished children with kwashiorkor or marasmus are at high risk

ALL

Select one or more options:

- a. dehydration
- b. immune deficiency and infection
- c. hypothermia
- d. refeeding syndrome during treatment
- e. hypoglycemia, electrolyte balance disorders

Translated text

22:50 54%
Google Lens
https://drive.google.co...
Czech → English

58% 8:19 am

Task 3

Barrett's esophagus

Select one or more options:

- occurs in patients with gastroesophageal reflux
- b. it is created by replacing the normal one squamous epithelium with jicnu cells carcinoid
- c. is created by substitution normal squamous epithelium esophagus with metaplastic columnar epithelium
- d. represents a risk factor for the development of esophageal adenocarcinoma
- E. occurs in patients with atrophic gastritis

Translated text

01:18 34%
Google Lens
https://drive.google.com...!
Czech → English

8:53 am LTE

to be superficial, they are decreased and paCO2 increased

he breathes deeply, value paCO2 is lowered

Task 7

Malabsorption syndrome

Select one or more options:

and can manifest as iron deficiency

b. is usually accompanied by obstinate constipation

it is usually accompanied by diarrhea

d. diarrhea is often accompanied by a tendency to develop metabolic alkalosis

can manifest itself vitamin B12 deficiency

Translated text
Send to Translate Home
Select all
Did you find these results useful?
Yes No

02:00 41%

https://drive.google.co...

Google Lens

Czech → English

Task 6

Task 1

IMG 3198.JPG

IMG_3199.JPG

The cause of late postprandial syndrome is

Select one or more options

rapid movement of carbohydrates into the intestine and their rapid resorption

bdiabetes mellitus 2 type

posthyperglycemic hypoglycemia

d bile secretion disorder

IMG 3234.JPG

IMG_3235.jpg

Translated text

Send to Translate Home

Select all

Did you find these results useful?

Yes No

02:04 21%

Google Lens

Czech → English



Translated text

Send to Translate Home

Select all

Did you find these results useful?

Yes No

02:11 19%
https://drive.google.co...
Google Lens
Czech → English

10:42 a.m 4G

Task 4

We can expect isotonic hyperhydration

Select one or more options:

- [ ] and decreased central venous pressure
- [x] a decrease in the concentration of total protein in the plasma ✓
- [x] formation of edemas ✓
- [ ] d. increased serum sodium level (if there is no concomitant kidney disease)
- [x] e reduced number of erythrocytes per mm3 in the blood ✓

Task 5

Translated text
Send to Translate Home
Select all
Did you find these results useful?
Yes
No

23:26 70%
Google Lens
https://drive.google.co...
Czech → English

16:04 83%

Task 1

Hirschsprung's disease
(megacolon congenitum)

Select one or more options:

a. is a congenital disease, which lacks ganglion cells in intestinal plexus

b. is manifested by diarrhea due insufficient digestion of fats

c. manifests itself organic constipation in due to congestion the intestinal contents

d. is an intestinal manifestation of congenital progressive myodystrophy

Translated text

01:53 • 43%

Google Lens

https://lens.google.co...

Czech → English

Task 6

Task 1

IMG 3198.JPG

IMG 3199.JPG

Respiratory alkalosis is characterized

Select one or more options:

and a rise in serum bicarbonate

b. a decrease in pCO2

c increased excretion of H+ in the urine

d. reduced release of oxygen from hemoglobin de weaving

IMG 3234.JPG

IMG 3235.jpg

Translated text

Send to Translate Home

Select all

Did you find these results useful?

Yes

No

18:26 33%
https://drive.google.com...
Google Lens
Czech → English

Role 50/100

A 2.5-year-old bitch developed diarrhea, most likely due to a rotavirus infection. Her temperature is 37.8 degrees Celsius. during examination with sweat where it is pink. She vomited several times and had diarrhea 5-6 times every day for 2 days, her heart rate is 130 bpm (N 80-130 bpm). arterial blood pressure 100/65 mm Hg still normal), answers simple questions and communicates with the doctor and her mother. Choose the 1 most appropriate statement about the best way to treat the patient:

Select one or more options:

- It requires immediate hospitalization, but infusions cannot be given until the doctor receives the results laboratomiho vyleteni It can
- be treated at home with the recommendation to drink orally rehydration drinks✓
- He needs immediate hospitalization with infusion therapy of 0.9% NaCl
- It can be treated at home with the recommendation of an autopsy of sweet tea and fruit juice
- He needs immediate hospitalization with infusion therapy with 5% glukday solution

The correct answer is: It can be treated at home with the recommendation to drink oral rehydration drinks.

Dream –U... 3.JPG
Huh...
Translated text

01:20 33%

Google Lens

https://drive.google.co...

Czech → English



Translated text

Send to Translate Home

Select all

Did you find these results useful?

Yes No

17:57 41%
Google Lens
https://drive.google.co...
Czech → English

Task 9 Point 0.00/1.00

A 65-year-old patient was hospitalized with symptoms of acute respiratory disease. He has a history of smoking for 25 years and obstructive pulmonary disease for 10 years. Within hours, a PCR analysis showed that he was positive for COVID-19 and, according to a lung X-ray, he had symptoms of pneumonia. His respiratory rate is 23/min, heart rate 105 tepl/min, arterial blood pressure 130/80 mm Hg.t-38.3 st.C, leukocyte count 3.9109/L Choose the correct statement about the patient with regard to the largest possible complications in the next 2 days:

Choose one or more options: The

- patient is at high risk of developing respiratory acidosis due to hypoventilation
- The patient is at high risk of developing metabolic and callosis due to diarrhea
- The patient is at high risk of developing metabolic acidosis due to vomiting
- The patient has a high risk of developing respiratory alkalosis due to hypoventilation

(e) The patient is high risk development of dehydration in due to kidney damage

Dream -U... 3.JPG
Dream -u... 6.JPG
Translated text

21:18 565%
https://drive.google.co...
Google Lens
Czech → English

8:53 am LTE

Task 5

On the right side of the picture we see the pathology in the area of the esophagus/stomach junction. About what pathology, or what type is it?



Select one or more options:

- [ ] a. Stomach ulcer
- [ ] b. Achalasia
- [ ] Colic diverticulum
- [ ] Sliding hernia
- [ ] Paraesophageal hernia

Translated text
Send to Translate Home
Select all
Did you find these results useful?
Yes No

16:32 54%
Google Lens
https://lens.google.co...
Czech → English

02.CZ 9:25 @ 83%
moodle.upol.cz

605

When proton pump inhibitors (K+/H+ ATPase) are administered, k

Select one or more options:

- [x] and decrease in HCL production in stomach
- [ ] b. increase in HCL production v stomach
- [ ] c. reduction of gastrin production
- [x] d. increase in gastrin production

Complete the tour

Test navigation

Translated text
Send to Translate Home
Select all
Did you find these results useful?
Yes No

02:13 38%

Google Lens

https://lens.google.co...

Czech → English

...



The value of the anion deficit (anion gap) rises at

770

Select one or more options:

- arenaini tubules of acidosis
- b hypoxic lactic acidosis
- c diabetic ketoacidosis
- dacidosis with diarrhea

Translated text

Send to Translate Home

Select all

Did you find these results useful?

Yes No

23:15 64%
Google Lens
https://drive.google.co...!
Czech → English

4:04 p.m

Task 4

Select the typical symptoms of vitamin deficiency:

Select one or more options:

- [ ] a. Vitamin C deficiency significantly increases the risk of thrombosis
- [ ] Long-term use of broad-spectrum antibiotics can lead to vitamin K deficiency and increase the risk of bleeding
- [ ] Vitamin A deficiency leads to cross-blindness, xerophthalmil keratinization of the skin and mucous membranes
- [ ] Vitamin D deficiency increases the risk of muscle weakness, muscle cramps
- [ ] e. Lack of vitamin D increases the risk of night blindness, xerophthalmia

Translated text

23:10 61%
Google Lens
https://drive.google.co...
Czech → English

12:36 p.m 71%

Task 9

The lack of bicarbonates arises as a result

Select one or more options:

- [ ] a. hypernatremie
- [ ] b. often because of insufficient bicarbonate intake
- [ ] c. hyperaldosteronismu
- [x] d. renal tubular acidosis
- [x] and diarrhea

Something real + diarrhea

Complete the tour

Translated text

18:27 33%
Google Lens
https://lens.google.co...
Czech → English

Ulaha 6

//moodle.upol.cz/pluginfile.php/1127661/mod_resource/content/5/Cv.%204%20Nyživa.pdf

Credit

Task 9

Symptoms and complications of atrophic gastritis include:

Select one or more options:

- [ ] and prolonged fever
- [x] b. megaloblastic anemia
- [x] stomach cancer
- [x] d. decreased iron absorption
- [ ] e. Severe heartburn is the most common symptom

The correct answers are: reduced iron absorption, cancer

Task 10

Dream -U... 3.JPG
Dream U... 6.JPG
Translated text

15:15 70%
Google Lens
https://drive.google.co...
Czech → English

16:22

Task 4

Causes of normochloremic metabolic acidosis do NOT include

Select one or more options:

- [ ] a. starvation
- [ ] b. alcohol poisoning
- [ ] c. salicylate poisoning
- [ ] diabetes insipidus

Task 5

Plasma chloride concentration is increased at

Select one or more options:

- [ ] A. ----------
- [ ] B. prolonged vomiting
- [x] with severe diarrhea
- [ ] D. respiratory alkalosis
- [ ] E. renal tubular acidosis

Translated text

17%

PathoPhys Online Tests (2)

× Untitled Notebook... × Untitled Notebook... × VAA11_II ×

patients' blood osmolarity

is high

With the biggest

chances are the patient

has an elevated level of ketone

bodies in the plasma ✓

With the biggest

chances are the

patient is dehydrated

due to osmotic

diuresis ✓

Most

likely, the patient

has a normal or

decreased anion

gap

21:14 66%
Google Lens
https://drive.google.co...
Czech → English

58% 8:19 am

moodle.upol.cz/mod/qu

16

Task 2

Causes of increased interstitial fluid volume and edema include

Select one or more options:

- a. increased venous pressure due to heart failure
- b. decreased plasma protein levels due to liver failure
- c. venous obstruction v due to thrombus
- effect of histamine
- E. arteriolar narrowing and venular dilation due to angiotensin II

Translated text

13:53 22%
Google Lens
https://drive.google.co...
Czech → English

Task 2

Which acid-base balance disorder develops primarily in patients with long-term simple starvation

Select one or more options:

- [ ] A. Combined metabolic and respiratory acidosis
- [ ] B. Metabolic acidosis
- [ ] C. Respiratory acidosis
- [ ] D. Metabolic alkalosis
- [ ] E. Respiratory alkalosis

IMG_1214.JPG IMG_1218.JPG
Translated text

18:08 38%
Google Lens
https://lens.google.co...
Czech → English

Ulaha 6

Cove 1

Role 40.50/1.00

Major vomiting will lead to

Select one or more options:

and metabolic acidosis in

due to large chloride losses

metabolic alkalosis due

to a large loss of

hydrogen ions in the acidic

gastric juice ✓

c. Hypochloremia

From metabolic acidosis

due to compensatory formation

HCl

Be e metabolic alkalosis

due to compensatory

hyperkalemia

The correct answers are:

metabolic alkalosis as a result

large losses of hydrogen ions in

acid gastric juice,

Hypochlorémil

Dream Hu... 3.JPG
Dream D... 6.JPG
Translated text

10:51 90%
Google Lens
https: //d...e.com/d ...
Czech → English



19.JPEG
0 四 20.jpg

O Helicobacter pylori pay that

Select one or more options:

A. Increases the risk of mortar (mucosa-associated lymphoid tissue) lymphoma

b. In particular, it is applied in pathogenesis Stress ulcers

c. Transmission path is probably fecally oral and oral

d. His toxic-metabolic municipation can often induce acute pancreatitis

23.jpg
24.jpg
O
Translated text
Offline · Some features are not available
Send to Translate Home
Select all
Sh
Did you find these results useful? Yes No

10:50 https://drive.google.com/d... 90%

Google Lens

33.jpg

Czech → English

19.JPEG

0 四 20.jpg

O Helicobacter pylori pay that

Select one or more options:

A. Increases the risk of mortar (mucosa-associated lymphoid tissue) lymphoma

b. In particular, it is applied in pathogenesis Stress ulcers

c. Transmission path is probably fecally oral and oral

d. His toxic-metabolic municipation can often induce acute pancreatitis

23.jpg

24.jpg

Select one or more options:

Translated text

Offline · Some features are not available

Send to Translate Home

Select all

Did you find these results useful? Yes No

10:50 90%
Google Lens
https: //d...e.com/d ...
Czech → English
19.JPEG
0 20.jpg

Anion deficit value (anion GAP) rises at

770

Select one or more options:

- [ ] and Renal tubular acidosis
- [x] hypoxic lactate acidosis
- [x] c diabetic ketoacidosis
- [ ] D acidosis in diarrhea

23.jpg
0 國 24.jpg
Translated text
Offline · Some features are not available
Send to Translate Home
Select all
Sh
Did you find these results useful? Yes No

10:50 90%
Google Lens
https: //d...ogle.com/d ...
Czech → English
19 JPEG
20 jpg
Between the anti-factors in the
pathogenesis of the peptic ulcer pati
Select one or more options:
Bintakni microcirculation
highlights of mucosa
Caspartate Proteinasis
d normer secretion
prostaglandins
23.jpg
0 國 24.jpg
Translated text
Offline · Some features are not available
Send to Translate Home
Select all
Did you find these results useful?
Yes
No

10:50 90%
Google Lens
https: //d...g.google.com/d ...
Czech → English



Translated text
Offline · Some features are not available
Send to Translate Home
Select all
Did you find these results useful? Yes No

10:50 90%
Google Lens
https://drive.google.com/d ..
Czech → English

19.JPEG

620.jpg

In isotonic dehydration

Select one or more options:

There is a vasoconstriction in splanchnice areas

B There is a osmotic shift of water from Intersticia to cells

CSE volume intracellular fluid increases

d. The skin turgor decreases

23.jpg

0 國 24.jpg

When passing succulents in the evaluation of an acid-base disorder applied

Select one or more options:

Among the states of secondary obstipation

Select one or more options:

Translated text
Offline · Some features are not available

10:50 90%
Google Lens
https: //d...g...gle.com/d ...
Czech → English

When poisoning salicylates, in the development of an acid-base disorder applied

Select one or more options:

- a. The formation of ketolenas → metabolic acidosis
- b tiny breathing → centraryespiratory alkalose
- c. pride
- d metabolic acidosis
- Eutlum windscreen center → respiratory acidosis

Translated text
Offline · Some features are not available

10:50 90%
https: //d...g...gle.com/d ...
Google Lens
Czech → English

In investigating acid-base balance pay for Base Excess (BE) to follow

Select one or more options:

- [ ] and the value depends on hemoglobin concentration
- [ ] The sum of all bushing bases
- [ ] It's Distributed the Numental BB (Buffer Base) values from it (normal) values, BE-8B-NBB
- [ ] d normally moves around 42 meq / l
- [x] PLI metabolic alkalia

Translated text
Offline · Some features are not available

10:50 90%
Google Lens
https: //d...e.com/d ...
Czech → English
Choose one or more options

Ethanol poisoning without vomiting leads to

Select one or more options:

- and metabolic acidosis in the Due ethanol
- In the terminal state to respiration of acidosis due to the attenuation of the wind center
- c. Metabolic alkalosis in the result of HCL secretion stimulation
- d. metabolic alkalosis due to ethanol conversion
- e. Respiratory alkalosis due to hyperventilation

Translated text
Offline · Some features are not available

10:49 90%
https: //d...g...gle.com/d ...
Google Lens
Czech → English

Filles

What is typical of celiac?

Select one or more options:

- [x] and gluten induced autoimmune response
- [ ] b. autoimmune odler induced by lactose
- [ ] c. It is often associated with other autoimmune diseases
- [x] d Compliance with a lifetime Acknowledged Gluten Free Diet is the best treatment
- [ ] e Unnipped number of lymphocytes in bowel mucosal

neviem

Translated text
Offline · Some features are not available
Send to Translate Home
Select all
Sh
Did you find these results useful? Yes No

01:37 46%
Google Lens
https://lens.google.co...
Czech → English
...

Task 6

Task 1

Ethanol poisoning without vomiting leads to

Select one or more opacities

- [ ] and metabolic acidosis as a result of the ethanol strain
- [ ] in terminal state k respiratory acidosis due to depression of the respiratory center
- [ ] c. metabolic alkalosis in the dose of stimulation of HCL secretion
- [ ] d. metabolic alkalosis due to ethanol conversion
- [ ] e. respiratory alkalosis due to hyperventilation

Dream Fu.. 3.JPG
Dream FÜ ... 6.JPG
0
Translated text

23:33 69%
Google Lens
https://drive.google.com...!
Czech → English



Translated text

18:28 33%

Google Lens

Czech → English

Ulaha 6

Clove 1

IMG_3198.JPG

IMG_3199.JPG

Task 4

Choose the correct statement microcirculation

Select one or more matres

A. It usually increases vallitation and increases the risk of iron formation

B. In inflammation, the permeability of the capillaries and post-capillary venules is usually increased

C. Normální (nepoškozené) endoteliální buňky neprodukují NO

D. Nepoškozený endotel má antitrombotický potenciál

E. Pándátni bulk produce relatively high concentrations of NO as a result of inducible activation NO-syntázy

The correct answer is to use a botic potential to produce a relatively high level of NO-induced innate NO-synczy minit, which usually results in permeability

Kaptar and pokapamiche

IMG_3234.JPG

IMG_3235.jpg

Dream Hu... 3.JPG

Snl D... 6.JPG

Translated text

16:32 54%
Google Lens
https://drive.google.com...!
Czech → English

02.CZ 9:25 @ 84%
moodle.upol.cz

Task 7

673

It is an important finding in pure water dehydration

Select one or more options:

- [ ] a. reduced concentration plasma proteins
- [ ] b. unchanged sodium level
- [ ] c. increase in MCV values (mean red cell volume), i.e. mean volume of erythrocytes
- [ ] increased sodium level.

Task 8

Translated text
Send to Translate Home
Select all
Did you find these results useful?
Yes No

23:14 64%
Google Lens
https://drive.google.co...
Czech → English

O2 - CZ|O2 - SK 10147% 13:57

Task 7

Lactose intolerance in adults is caused by

Select one or more options:

- [ ] and excessive fission of lactose to glucose and galactose in the intestine
- [ ] hydrolysis disorder lactose in the intestine
- [ ] C. disruption of the mechanisms ensuring the transport of lactose and galactose through the intestinal wall
- [ ] d. a disorder of lactose phosphorylation in the intestinal wall

Task 8

Select an important cause/important

Translated text

00:16 47%
Google Lens
https://drive.google.co...
Czech → English

58%8:20

Task 8

The following can be said about a patient in lactic acidosis

Select one or more options:

- [ ] A. has increased value standard bicarbonates
- [ ] B. the patient is most likely suffering from severe hypoxia
- [ ] C. has a normal value of standard bicarbonates
- [ ] D. he cannot have lung disease because it only leads to respiratory acidosis
- [x] E. has an increased anion gap

Translated text

23:17 65%
Google Lens
https://drive.google.co...
Czech → English

Translated text

15:20 69%

https://drive.google.co...

Google Lens

Czech → English

T-Mobile CZ 8:24 @ 100%

moodle.upol.cz

Task 5

The main 2 etiological factors of Mellory Weiss syndrome include

Select one or more options:

- [ ] a. Bulimia
- [x] b. GERD (gastroesophageal reflux disease)
- [ ] c. Anorexia nervosa
- [x] d. Alcoholism
- [ ] e. Cirrhosis of the liver

Task 6

IMG_1214.JPG IMG_1218.JPG

Task 2

Translated text

16:34 53%
Google Lens
https://drive.google.co... :
Czech → English



Translated text

10:53 89%
Google Lens
https: //d...e.com/d ...
Czech → English
ll o2.cz 14:37 38%
moodle.upol.cz

Task 3

Secondary constipation may occur at

Select one or more options:

- and hyperthyroosity
- immobilization
- hypokalemia
- d. Hyperhydration

Task 4

Translated text
Offline · Some features are not available
Send to Translate Home
Select all
Sh
Did you find these results useful? Yes No

10:49 90%
https: //drive.google.com/d ...
Google Lens
Czech → English

drive.goog

UPOL.CZ/pluginfile.php/1127661/MOD_RESOURCE/CONTENT/5/CV.%204%20V. \ tPdf \ t

Job 11

Choose the correct / ä assembly and chronic gastritis type A

Choose one or more floss

- [x] Patients have an increased risk of emergence of stomach cancer
- [ ] b. Usually develop in 2-3 weeks after acute gastrite
- [x] Patients often lack of vitamin 812
- [ ] d. Patients have an abusle increased production of HCl and duodenal ulcers
- [x] patients often reduced HCl production

Correct answers are: patients have often reduced HCl production, patients n increased risk of stomach cancer

Translated text
Offline · Some features are not available
Send to Translate Home
Select all
Sh
Did you find these results useful? Yes No

10:49 90%
Google Lens
https: //drive.google.com/d ...
Czech → English

Filles

Job 1

Drive.Google.com

11 Resource Ntent / 5 / CV. % 204% 20V.

Task 4

Select the correct / correct microculation statement:

Choose one or more lives:

and will usually leads to vasodilation and increases the risk of thrombus formation

b. Beyem of zánětu is usually increased by the permeability of capillary and postkapillary Venul

c Nomali (undamaged) endothelial cells I do not produce no

Undototel Má antithrombotic potential

In inflammation, endothelial cells can produce relatively high quantities due to activation inducible

No-synthasis

Correct answers are: Undambled endothelí má antithrombotic potential, when zánětu can endothelial cells produce relatively high MNABasis due to activation of inducible no-synthase, during inflammation usually increases the permeability of capillarics and postkeplámích Venul

Translated text
Offline · Some features are not available
Send to Translate Home
Select all
Sh
Did you find these results useful? Yes No

10:49 90%
Google Lens
https://drive.google.com/d ..
Czech → English

Filles

Street 5 Boda 1.00 / 1.00

In the case of 2.5 years of view, there was a use of the greatest likelihood in the result of rotavirus infection. Several times vomited and diarrhea 5-6 times every day for 2 days, its heart rate is 130 beats per minute (N 80-130 pulse per minute) arterior blood pressure 100/65 mm HG (still in normal), corresponding to simple questions and communicates with Lékalem and her mother. Choose 1 most appropriate claims about the best of treatment patients

Choose one or more options

- Needs immediate hospitalization, but infusions cannot be done than to succeed the results of the laboratory examinations
- Can be treated at home with a pit recommendation oral rehydration drinks ✓
- Needs immediate hospitalization with infusion therapil 0.9% NAC
- Muže be treated at home with Recommended Pit Vice Sweet tea and fruit juice
- E need immediate hospitalization with infusion therapy 5% glucose solution

The correct odler is: Can be treated at home with Recommended pit ororate rehydration drinks

Translated text
Offline · Some features are not available

10:49 90%
https: //d...g...gle.com/d ...
Google Lens
Czech → English
File

osmolarity of the blood of the patient

high

With the most

truthenedobnesti ma

patient increased levels

of ketolena in plasma ✓

The patient is likely

to dehydration in the

result of the Osmatric

Diune ✓

truthenedobnesti

ma patient normally

or reduced anionic

gap

Manage Odoids are: S

The shortcipimacy of the patient

Tachycardia VDüstedku dehydration

and activation of

sympathetic nervous

the system, with

the most likely to defend the

patient dehydration due to

Asmotic diagnostics, with the

world's most likely

Osmolarity of blood patient

high, with the largest

The likelihood of a

pececient perceived ketolus

plasma

Translated text
Offline · Some features are not available

10:49 90%
Google Lens
https://drive.google.com/d... :
Czech → English

Files

Task 9 0.50/1.00 points

Losses of water and solutes during excessive sweating lead to

Select one or more options: a.

- hypernatremia
- hypertonic dehydration
- hypotonic dehydration
- d. isotonic dehydration
- e increase in volume intracellular fluid due to osmotic shifts of water between the extracellular and intracellular compartment

The correct answers are:
hypertonic dehydration,
hypernatremii

Translated text
Send to Translate Home
Select all
Lis
Did you find these results useful? Yes No

10:49 90%
https://drive.google.com/d... :
Google Lens
Czech → English
Files

Task 4 Point) 1/20/1.00

The following characteristics are not used for the classification of edema:

Select one or more options:

- Pitting a non-pitting
- Acquired and Inherited✔
- Primamia sekundäm lymphedema
- Hyperosmolární a hypoosmolar✔
- Local and generalized

The correct answers are:
Hyperosmolar and hypoosmolar
Acquired and inherited

Translated text
Send to Translate Home
Select all
Lis
Did you find these results useful? Yes No

10:48 90%
Google Lens
https://drive.google.com/d...
Czech → English

Task 8 0.00/1.00 points

Intermittent fasting can mean that a person:

Select one or more options:

- He consumes every week only 1 type of food (e.g. 1 week apples, 2. week bread, 3. week cheese...)
- Consumes a low-protein diet for 1 day, then a low-fat diet for 1 day, then a low-carb diet for 1 day X
- Consumes food for 6 hours a day and does not consume food for 18 hours
- Consumes food for 8 hours a day and does not consume food for 16 hours
- Consumes only 1 type of food every day (e.g. 1 day apple, 2 day bread, 3 day cheese...)

The correct answers are: Consumes food for 8 hours a day and does not consume food for 16 hours Consumes food for 6 hours a day and does not consume food for 18 hours

Translated text

14:40 Wed, 7 Dec

Google Lens

Czech → English

Task 3

Leiden mutation

Select one or more options:

- a. carriers have an extended Quick's test
- b. leads to thrombophlebitis in carriers
- c. vede k prodloužení krvácivosti
- d. carriers have a shortened aPTT test

Abhinav Jethi

Comment or add others with @

Viewers of this file can see comments and suggestions.

Comment Cancel

Tina 2

Task 4

Usha 9

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

14:40 Wed, 7 Dec

Google Lens

Czech → English

Type IV = T cells monocyte macrophages

Which cells are most important for immunopathological reactions IV. type? (choose 1 best answer)

Select one or more options:

- [ ] b. Basophilic granulocytes, Eimous cells, B-lymphocytes
- [ ] b. Neutrophil granulocytes eosinophilic granulocytes
- [ ] c. Basophilic granulocytes plasmids
- [ ] d. Basophilic granulocytes eosinophilic granulocytes
- [x] T-cells, monocytes and macrophages

Abhinav Jethi

Comment or add others with @

Viewers of this file can see commer suggestions.

Comment Cancel

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

DIC - hemolytic anemia

Task 7

Anemia accompanying the ischemic phase of disseminated intravascular coagulopathy (DIC) is

Select one or more options:

- a. hemolytic
- b. hemorhagická
- c. diminutive
- d. macrocytes

Abhinav Jethi

Comment or add others wit

Viewers of this file can see com suggestions.

Comment Cancel

Task 8

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Task 6

Primary biliary cirrhosis is not:

Select one or more options:

- [ ] consequence of portal hypertension
- [ ] liver damage due to the action of chemotoxic substances
- [ ] primary liver damage due to the effects of alcohol
- [ ] d autoimmune disease

Abhinav

Comment or a

Viewers of this fil
suggestions.

Comment

Task 9

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

Google Lens

Czech → English

aspicin inhibits COX

02.02

Which of the following statements about aspirin is false?

Select one or more options:

a. Aspirin leads to a decrease in the synthesis of thromboxane A2

b. Aspirin reduces the size and stability of platelet aggregates

c. Vitamin K is an essential cofactor in the reaction that is inhibited by aspirin

d. Aspirin inhibits the destick enzyme cyklooxygenázy-1 (COX-1)

Účinky aspirinu na koagulaci [illegible] mom

Abhinav Jet

Comment or ado

Viewers of this file
suggestions.

Comment

Ulaha 9

113

ranslated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Among the most common causes of acute pancreatitis are:

Select one or more options:

- [ ] a. hyperglycemia
- [x] b. cholelithiasis
- [ ] c. gastrin hypersecretion
- [x] d. alcoholism

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

13:30 Wed, 7 Dec

Google Lens

Czech → English

formed in liver

Fe absorption supression

Hepcidin

Select one or more options:

- a. it is mainly formed in the liver ✓
- b. suppresses the absorption of iron in the duodenum ✓
- c. is excessive produced at hereditary hemochromatóze
- d is a peptide controlling the metabolism of the medium

Abhinav Jethi

Comment or add other

Viewers of this file can see suggestions.

Comment Cancel

Complete the tour

Tact navigation

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

15:46 Tue, 6 Dec

Google Lens

Czech → English

Ascites > dyspnea

Ascites:

Select one or more options:

- [ ] A It is accompanied by reduced secretion of aldosterone
- [x] B May cause shortness of breath in the patient
- [ ] C Can be caused by decreasing Na resorption in the proximal tubules
- [x] D. It is accompanied by water and sodium retention
- [x] E May be accompanied by a diluent hyponatremia

dyspnea

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

A final common coagulation pathway involves:

Select one or more options:

- [ ] a. Prothrombin formation
- [ ] b. N-acetylation by Vitamin A.
- [x] c. Formation of the prothrombinase complex
- [ ] d. Formation of fibrinogen
- [ ] e. Degradation of factor XIII

prothrombinase complex

Abhinav J

Comment or a

Viewers of this fil suggestions.

Comment

Task 6

ranslated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

Google Lens

Czech → English

Task 4

thrombocytosis

Thrombocytosis of a reactive nature is most often associated with (3):

inflammation
bleeding
splenectomy

Select one or more options:

a. Acute or chronic inflammation ✓

b. I'm bleeding ✓

c. Splenomegalii

d. Klonain expansion of megakaryocytes due to myeloproliferative diseases

e. Splenectomy ✓

Abhinav J

Comment or a

Viewers of this fil
suggestions.

Comment

Ulona 6

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

Google Lens

Czech → English

Choose the correct statement about hepatitis

Select one or more options:

- a. Hepatitis A is often (in 50% of patients) manifested by acute liver failure
- b. Viral hepatitis C is a risk factor for liver cancer
- c. Complications of chronic viral hepatitis C include anaemia
- d. Viral hepatitis B is a risk for liver cancer
- e. Complications of chronic viral hepatitis C are extensive hypoalbuminémií

Abhinav Jethi

Comment or add ot

Viewers of this file can
suggestions.

Comment

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

Google Lens

Czech → English

D-dimers are

Select one or more options:

- [ ] a. contain in their amino acid structure of serine and are therefore further cleaved by plasmin
- [ ] are important markers of ongoing thrombolysis
- [ ] they are no longer cleaved by plasmin
- [ ] proteolytic derivatives of fibrin
- [ ] proteolytic derivatives of fibrin and procalcitonin

Abhina

Comment or

Viewers of this
suggestions.

Comment

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

Google Lens

Czech → English

Reticulocytosis in a patient with pernicious anemia that occurs after initiation of treatment is suggestive of (2)

Select one or more options:

- [x] and activation of hematopoiesis
- [ ] b. shortened erythrocyte survival in this patient
- [ ] chemolytic crisis
- [x] positive effect of treatment with vitamin B12

Abhinav Is

Comment or ad

Viewers of this file suggestions.

Comment

Ulaha é

Task 5

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

14:39 Wed, 7 Dec

Google Lens

Czech → English

Liver cells create

Select one or more options:

- [ ] A. Immunoglobulin G
- [ ] B. Imunoglobulin M
- [x] Albumin
- [x] D Plasma coagulation factors II, V, VII, IX, X
- [x] EC-reactive protein (CRP)

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Task 9

Coombs test is Autoimmune hemolytic anemia

The direct Coombs test is usually positive in patients with

Select one or more options:

- pernicious anemia
- autoimmune hemolytic disease
- ahemia
- rheumatoid arthritis
- and sideropenic anemia

Abh

Comment

Viewers of th
suggestions.

Commen

Complete the tour

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

Wed, 7 Dec

Google Lens

Czech → English

It is true about the pathogenesis of thrombosis

Select one or more options:

- a. is the formation of blood clots within the circulatory system, excluding the heart
- b. defects in the fibrinolytic system cannot lead to the formation of a thrombus
- when the level of antithrombin II is reduced to 40%, the risk of thrombus formation is increased
- the resistance of the Mik factor to the proteolytic step is the most common cause of hypercoagulable rules

Abhinav

Comment or a

Viewers of this fil suggestions.

Comment

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

Google Lens

Czech → English

Task 5

When examining blood laboratories, in patients with a moderate gram-negative bacterial infection (Escherichia coli, Pseudomonas species), it is usually assumed:

Select one or more options:

- [ ] Polycytémie
- [ ] b. Agranulocytosis
- [ ] Neutrophia
- [ ] d. Lymphocytosis
- [ ] Eosinofilie

Abhina

Comment or

Viewers of this suggestions.

Comment

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

14:37 Wed, 7 Dec

Google Lens

Czech → English

nonspecific immunity j
macrophages, NK cells
complement

Task 4

Among the components of non-specific immunity we include

nonspecific immunity j

Select one or more options:

- a antibodies
- b macrophages, NK cells
- c complement
- d Those 8 lymphocytes

Abhinav

Comment or a

Viewers of this fil
suggestions.

Comment

Task 5

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Task 9

Blood viscosity is

↑ hematocrit ↑

Select one or more options:

- [ ] a. on hematocrit independent
- [ ] b decreases with increasing hematocrit
- [x] rises with increasing hematocrit
- [ ] d. tide proportional to hematocrit

Com

Viewers c

suggestion

Comm

oha 6

Complete the tour

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

14:36 Wed, 7 Dec

Google Lens

Czech → English

Task 1

The primary cause of anemia may be

Select one or more options:

- [ ] and increased production of erythrocytes
- [x] reduced production of erythrocytes
- [x] increased loss of erythrocytes
- [ ] d shortened prothrombin time

Ab

Commen

Viewers of suggestion

Com

Ulona &

Task 2

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Task 1

Anemia of chronic diseases arises

Select one or more options:

- during inflammation by inflammatory cytokines
- th as a result of bleeding
- drink, in case of chronic infections, rheumatic inflammation
- d. with a lack of folic acid

Abhinav Jef

Comment or add

Viewers of this file c
suggestions.

Comment

Task 2

ranslated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

Google Lens

Czech → English

The most common causes of acute liver failure in Europe are (choose 1 answer):

Select one or more options:

- a. Acute autoimmune hepatitis
- b. Acute hepatitis caused by intoxication of hepatitis B virus or B+ Da acetaminophen paracetamol)
- c. Acute intoxication caused by consumption of Amanita phalloides mushrooms
- d. Acute hepatitis caused by viternous hepatitis Aa virus
- e. Acute hepatitis

Abh

Comment

Viewers of t suggestions

Commen

OLA

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

14:34 Wed, 7 Dec

Google Lens

Czech → English

Task 4

Thrombocytosis can occur in the following diseases or conditions:

Select one or more options:

- [ ] and Chronic inflammatory diseases (for example, Crohn's disease)
- [ ] b. Hypersplenismus
- [ ] c. DIC
- [x] d. Polycythemia vera
- [ ] Thrombotic purpura

thrombocytosis = high number of platelets

Polycythemia vera = erythrocytosis, however rare

Abh

Comment

Viewers of th

suggestions.

Commen

Task 5

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Task 7

Elevated conjugated bilirubin, unconjugated bilirubin, AST, ALT, GGT in a 58-year-old adult are most likely to be signs of (choose 2 correct answers)

Select one or more options:

- [ ] a. Acute pancreatitis
- [ ] b. Acute obstruction of the bile duct
- [x] c. Alcoholic cirrhosis of the liver
- [ ] d. Cystic fibrosis
- [x] e. Hepatitis C virus

Abhin

Comment about

Viewers of thi suggestions.

Comment

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Task 3

Cholesterol cholelithiasis is NOT among the risk factors

Select one or more options:

- cholesterol hypersecretion
- male gender
- pregnancy
- hyposecretion of Hut's acids

Task 4

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

Google Lens

Czech → English

Vodafone CZ LTE 11:20 a.m 93%

Task 2

Vitamin B12 deficiency anemia a

tetrahydrofolic acid is

morphologically characterized

Select one or more options:

a makrocytózo

b microcytosis

c the presence of megaloblas s

d costar dient

e normocytózo

Com

Viewer

sugges

Con

Task 3

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Quick test (prothrombin time)

Select one or more options:

- [ ] a. is prolonged in hemophilia A
- [x] b. tests the coagulation activity of the external plasma coagulation system
- [ ] c. it is prolonged in patients with vori Willebrand disease
- [ ] d. tests the coagulation activity of the internal plasma coagulation system it is prolonged in advanced liver parenchyma damage

Commen

Viewers of suggestion

Com

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Task 1

Relaxation is largely responsible

for the discomfort accompanying

hay fever

Select one or more options:

- [x] histamine
- [ ] b. interferons
- [ ] c leukotrienes
- [ ] d. prostaglandins

Com

Viewer

sugges

Con

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

13:58 Wed, 7 Dec

Google Lens

Czech → English

Task 5

Which of the following manifestations decreases the activity of factor VIII due to an increased rate of degradation?

Select one or more options:

- a. Liver disease
- b. Von Willebrand disease
- c. Vitamin K deficiency
- d. Hemofilie B
- e. Hemophilia d

factor X = V

Abh

Comment

Viewers of th
suggestions.

Commen

Task 4

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

It is used in the pathogenesis of acute pancreatitis

Select one or more options:

- [ ] and. decreased activity of phospholipáA2
- [ ] b. reduced cathepsin B activity
- [ ] increased intracellular trypsin activation
- [ ] d. increased activity elastase

Complete the tour

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Macrocytic and normochromic anemias may include:

Select one or more options:

- [ ] and. sickle cell disease
- [x] deficiency anemia folic acid
- [ ] c. hereditary spherocytosis
- [x] megaloblastic anemia
- [ ] e. sideropenic anemia

megaloblastic anemia (B12 – folic acid deficiency)

Com

Viewers

suggestions

Comm

ranslated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

Google Lens

Czech → English

What is the typical set of events in the pathogenesis of alcoholic liver cirrhosis

Select one or more options:

- [ ] and Alcohol consumption → pancreatitis → cholecystitida → steatosis - cirrhosis
- [ ] b. Alcohol consumption → cholecystitida → steatosis → fibrosis → cirrhosis
- [ ] Alcohol consumption → fibrosis → steatohepatitis → black steatosis
- [ ] Alcohol consumption → shentosis → steatohepatitis → fibrosis → cirhóza
- [ ] e. Konzumace alkoholu → steatohepatitida → fibróza → steatóza → cirhóza

Abhinav

Comment or a

Viewers of this fil suggestions.

Comment

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

A mean erythrocyte volume (MCV) of 75 fl is compatible with the following diagnoses

Select one or more options:

- [ ] a. deficiency anemia Vitamin 812
- [ ] b autoimmune hemolyticka anemia
- [x] sideropenic anemia
- [ ] d. deficiency anemia erythropoán
- [x] polycythemia vera

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Task 6

Combination of anemia, normal serum
Fe concentration, reduced plasma
haptoglobin, high LDH levels
meaning most likely (1)

Select one or more options;

- anemia with chronic blood loss
- [illegible]
- riemolytic anemia
- lack of iron in the body

Abhinav Jethi

Comment or add others with

Viewers of this file can see com
suggestions.

Comment Cancel

Task 7

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

Google Lens

Czech → English

Task 7

An Rh- individual will develop anti-D antibodies

Select one or more options:

- a. already during intrauterine life
- b. shortly after birth
- after Rh+ blood transfusion
- as a result of the delivery of an Rh+ fetus

Abhinav Jethi

Comment or add others

Viewers of this file can see suggestions.

Comment Canc

Task 8

Genetic disorders related to later damage

(Wilson's disease, hemochromatosis

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

14:31 Wed, 7 Dec

Google Lens

Czech → English

Task 7

Sideropenia

Select one or more options:

- [ ] can be manifested in the body by an increase in serum ferritin
- [ ] macrocytic anemia can manifest in the body
- [ ] can be manifested in the body by an increase in transferrin in the serum (increased total binding capacity)
- [ ] anemia can manifest in the body

microcyta

Abhinav

Comment or a

Viewers of this file suggestions.

Comment

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

14:31 Wed, 7 Dec

Google Lens

Czech → English

Task 6

The risk of serious life-threatening infections is increased in patients with an absolute neutrophil count lower than:

Select one or more options:

a. 50 buněk / mm3 (0,05*10^9/L)

b. 5 buněk / mm3 (0,005*10^9/L)

c. 2500 cells n 3 (2,5*10^9/L)

d. 1500 buněk / mm3 (1,5*10^9/L)

e. 500 buněk / mm3 (0,5*10^9/L)

Abhinav Is

Comment or add

Viewers of this file ca suggestions.

Comment

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Task 4

The following values of the blood count - the number of erythrocytes 2.8 10 12/L, hematocrit 0.30, hemoglobin 85 g/L, MCHC 28% (N 32-36%), MCV = 107 FL (N 80-100 fL), MCH 35pg (N 26-36 pg) - correspond best diagnosis of anemia

Select one or more options:

- [ ] a. related to anemia due to acute hematoma
- [x] iron deficiency
- [ ] c. it is not about anomie Because the count erythrocytes le normali
- [ ] zerostark vitamin

Abhina

Comment o

Viewers of this suggestions.

Comment

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Task 3

Resistance to activated protein C

je

Select one or more options:

- [x] a. most often caused by a mutation in the coagulation factor V gene
- [ ] b.. always caused by a mutation in the coagulation factor V gene
- [ ] c. always caused by a mutation in the protein C gene
- [ ] d. associated with bleeding condition

Abhina

Comment or

Viewers of this f
suggestions.

Comment

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

Google Lens

Czech → English

Task 8

type 1 reaction → mast cells

Cells directly involved in

type 1 allergic reaction (anaphylaxis)

are

Select one or more options:

- [ ] a. CD4 lymphocytes
- [ ] b. CD8+ T-lymphocytes
- [ ] c. plasma cells
- [x] d. live mast cells

Abhinaya

Comment or

Viewers of this
suggestions.

Comment

Task 9

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

14:28 Wed, 7 Dec

Google Lens

Czech → English

Task 7

The anticoagulant effect of heparin is mediated

Select one or more options:

- a. protein C inhibition
- b. antithrombin
- c. factor Xa inhibition
- d. by thrombomodulin

Abhinav J

Comment or ado

Viewers of this file c
suggestions.

Comment

Task 8

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Task 9

lazy leukocyte syndrome

The syndrome of lazy leukocytes is characterized

Select one or more options:

- [ ] a. phagocytosis disorder with the impossibility of cells to form H202
- [ ] b. phagocytosis disorder
- [x] c. chemotaxis disorder
- [ ] d. inability to destroy phagocytosed bacteria

Abhinav Jethi

Comment or add othe

Viewers of this file can se suggestions.

Comment

Ph.D

Complete the tour

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

14:28 Wed, 7 Dec

Google Lens

Czech → English

Task 8

Deficiency anemia in chronic renal failure of erythropoietin is

Select one or more options:

- [x] normocytamf
- [ ] b. makrocytární
- [ ] c. microcytes
- [ ] 4 megaloblastic

Abh

Comment

Viewers of th suggestions

Commer

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

Google Lens

Czech → English

Genetic disorders ass

liver damage (Wilso

hemochromatosis, alpha 1 antitrypsin

deficiency) usually

Select one or more options:

- [ ] 3. They are inherited in an X-linked fashion, so they are much more common in boys
- [ ] b. They lead to chronic liver damage and usually appear in adulthood
- [ ] c They lead to acute liver failure in early childhood
- [ ] d. They do not have serious clinical manifestations related to other organs
- [ ] They can be partially influenced by special preventive measures

Abhinav Jethi

Comment or add c

Viewers of this file ca
suggestions.

Comment

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

Google Lens

Czech → English

Task 6

Predisposing factor chronic pancreatitis represents

Select one or more options:

- diabetes mellitus
- b. gastroesofageal reflux
- c. diabetes insipidus
- regular consumption alcohol

Abhir

Comment d

Viewers of thi suggestions.

Comment

Task 7

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Task 7

HIV infection primarily reduces the number in the blood

Select one or more options:

- a. B-lymphocytes
- CD4+ T-lymphocytes
- c. monocytes
- d. eosinophilic granulocytes

Abhinav

Comment or a

Viewers of this fil
suggestions.

Comment

Task 8

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

Google Lens

Czech → English

Patients with chronic granulomatous disease

Select one or more options:

a. they have a malfunction intracellular killing of the microbe

b. have a migration disorder leukocytes

c. they have a reduced number of leukocytes

d. they may have a disorder in the formation of reactive forms of oxygen

Abhinav Is

Comment or ad

Viewers of this file suggestions.

Comment

Utena &

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

14:43 Tue, 6 Dec

Google Lens

Czech → English

Task 1

It applies to pernicious anemia

Select one or more options:

- [ ] And they were among the hemolytic anemias
- [x] is the result of an autoimmune disease
- [ ] in the peripheral blood there are plutomic microcytes
- [x] there is a deficiency of vitamin 812

Task 2

Atopy means

Select one or more options:

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

APC (activated protein C) resistance

Yippee

Select one or more options:

a. hereditary diseases increasing the risk of thromboembolic disease

b. hereditary diseases increasing the physics of bleeding conditions

c. acquired disease increasing the risk of thromboembolic disease

d. acquired disease increases the risk of bleeding status

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

It does not belong to the corpuscular hemolytic anemias

Select one or more options:

- [ ] and. hereditary spherocytosis
- [ ] b. paroxysmal nocturnal hemoglobinuria
- [ ] Cautoimmune hemolytic anemia
- [ ] sickle cell disease

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

In vitamin B12 deficiency anemia

Select one or more options:

- [ ] and there is typically normochromic normocytic anemia in the blood count
- [ ] b. cytoplasm maturation is impaired, but nuclear maturation is normal
- [ ] there may be pancytopenia in the blood count
- [ ] d. damage to the nervous system may occur

Abhin

Comment c

Viewers of th suggestions.

Commen

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Pancreatic cancer risk factors

do NOT include

Select one or more options:

- [ ] a. diabetes mellitus
- [ ] b. high energy intake food
- [ ] c. smoked
- [ ] d. Hypersecretion of insulin

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

15:51 Tue, 6 Dec

Google Lens

Czech → English

We can see the development on the diagram the concentration of antibodies in the blood, viral RNA and ALT over time for a certain infectious agent. What agent is it?



Choose one or more options

- [ ] Virus hepatitis D
- [ ] Virus hepatitis A
- [ ] Virus hepatitis B
- [ ] Virus hepatitis E
- [x]

Hepatitis C

-> viral hep C

Abh

Comment

Viewers of th
suggestions.

Commen

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Task 1

aPTT

Select one or more options:

- [ ] a. tests the coagulation activity of the external plasma coagulation system
- [ ] is prolonged in hemophilia A
- [ ] is shortened with heparin therapy
- [ ] tests coagulation activity of the whitini system of plasma coagulation

Abhinav Jet

Comment or add

Viewers of this file c
suggestions.

Comment

anslated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

Google Lens

Czech → English

Task 8

What are the most common causes of jaundice in intravenous drug users?

Select one or more options:

- [ ] a. Hepatitis A
- [ ] 6. Hepatitis E
- [x] c. Hepatitis B
- [x] d. Hepatitis C
- [x] e. Hepatitis B+D

jaundice
Hepatitis
B+C+D

Abhinav J

Comment or ad

Viewers of this file
suggestions.

Comment

Ghond 2

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

15:46 Tue, 6 Dec

Google Lens

Czech → English

Ascites:

Select one or more options:

A It is accompanied by reduced secretion of aldosterone

May cause shortness of breath in the patient

C Can be caused by decreasing Na resorption in the proximal tubules

D. It is accompanied by water and sodium retention

E May be accompanied by a diluent hyponatremia

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

15:46 Tue, 6 Dec

Google Lens

Czech → English

C. He is deficient in vitamin K due to antibiotic treatment and suffers from insufficiency protein C

d. Must be hospitalized in ICU for high risk of life-threatening bleeding

With the biggest probability has no acute clinical manifestations of hemostasis disorders

Comm

Viewers suggesti

Com

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

After an acute viral respiratory tract infection in a 65-year-old man, laboratory tests showed: 160*10^9 platelets, INR 2.1, aPPT 28 sec, Duke bleeding time 5 min 30 sec, normal d-dimer level. He has a history of arterial hypertension, ischemic heart disease, arrhythmias, without other subjective health problems. The patient is most likely to:

Select one or more options:

- [x] It is necessary to ask sp about drinking warlarina
- [ ] Got pfauchu hemot type Av dusted Worby protilate brot Will
- [ ] Mi nedostatek vitaminu K v důsledku léčby antibiotiky z tří

Com

Viewers suggesti

Comm

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Repeated blood transfusions can lead to

Select one or more options:

- [ ] copper deposition in bones
- [ ] increase in transferrin saturation by Feiez
- [ ] c lowering of ferritin levels in the blood plasma
- [ ] can lead to increased iron deposition in the body
- [ ] concentration of transferrum in plasma

Translated text

Google Lens

Czech → English

An increased concentration of PAI-1 (plasminogen activator inhibitor-1) is a risk factor mainly for (1)

Select one or more options:

- [ ] thromboembolic diseases
- [ ] b. for GIT bleeding
- [ ] increased bleeding
- [ ] paroxysmal nocni hemoglobinurie
- [ ] M. DIC disen

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Serum sickness is a generalized form of allergy

type 3

Select one or more options:

- [ ] a. 3rd type
- [ ] b. 1. type
- [ ] c. 2nd type
- [ ] d. 4th type

Comm

Viewers

suggestions

Com

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Choose from the 3 most common causes liver cirrhosis in European countries:

Select one or more options:

- [ ] a Hepatitis A and E
- [ ] b Wilson's disease and alpha 1 antitrypsin deficiency
- [ ] c Hepatitis BaC
- [ ] d Alcoholic liver disease
- [ ] e Nonalcoholic steatosis of the liver

Com

Viewers

suggestion

Comm

Complete the tour

ranslated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

15:01 Tue, 6 Dec

Google Lens

Czech → English

Task 8

The frequency of hemophilia A is approx

Select one or more options:

a 1:500-1:1000 in men

Zen

1:200-1:400 in men

15000-1:10000 u

mužů

d 1: 100 000 - 1: 500 000

u roses and zen

e 1: 200 u mužů a žen

Comm

Viewers

suggest

Com

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

15:00 Tue, 6 Dec

Google Lens

Czech → English

Type 2 allergic reaction

Select one or more options:

a. it is caused by excessive production of IgE

b. it is called without remote hypersensitivity

c. it is caused by antibodies exclusively IgG

d. it is directed against the antigens carried by the Hal membrane of your body's cells, which causes hemolytická anémie, thrombocytopenia

type 2 = antigen

Com

Viewers

suggestion

Comm

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Anemia and at the same time found significant reticulocytosis is a typical manifestation

Select one or more options:

- [ ] and pernicious anemia
- [ ] b. infiltration of bone pulp by nådor
- [x] AIHA (autoimmune hemolytic anemia)
- [ ] d Fanconi anemia

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

6 Dec

Google Lens

Czech → English

Udhm 2

Prothrombin time (PT) increases in case

Select one or more options:

- Thrombocytopenia
- Marfanou's syndrome
- Vitamin K deficiency.
- Lack of factor VII
- Hemophilia A

Com

Viewers

suggestion

Yes No

Translated text

Home

Select all

Listen

Share

Download Czech

14:46 Tue, 6 Dec

Google Lens

Czech → English

Reticulocytes may be in the blood increased by

Select one or more options:

- [ ] a. chronic myeloid leukemia
- [x] b. in a patient after acute severe bleeding
- [ ] c. aplastic anemia
- [x] d. intravascular hemolysis

Task 2

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

14:45 Tue, 6 Dec

Google Lens

Czech → English

Task 2

Atopy means

Select one or more options:

- [ ] body condition after the administration of crede protein
- [x] changing the activity of the subject in the sense of pathological hypersensitivity
- [ ] systemic flower after a single parenteral administration
- [x] the number of antigen-specific and anti-antibodies in the premutation of Kantaka

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

14:43 Tue, 6 Dec

Google Lens

Czech → English

Task 1

It applies to pernicious anemia

Select one or more options:

- [ ] And they were among the hemolytic anemias
- [x] is the result of an autoimmune disease
- [ ] in the peripheral blood there are plutomic microcytes
- [x] *there is a deficiency of vitamin 812*

Task 2

Atopy means

Select one or more options:

- [ ] *[illegible]*

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

14:43 Tue, 6 Dec



Translated text
Send to Translate Home
Select all
Listen
Share
Download Czech
Did you find these results useful?
Yes
No

14:41 Tue, 6 Dec

Google Lens

Czech → English

moodle.upol.cz

Select condition(s) or diseases with increased risk of hemolysis:

Select one or more options:

- [x] Advanced burns affecting 60% aunt
- [x] Bacterial sepsis
- [x] State of disseminated intravascular coagulation
- [ ] Transfusion of ABO incompatible Rh-negative erythrocytes to Rh-negative to the patient
- [x] Transfusion of AC ompatib/nich Rh-negative erythrocytes to an Rri-positive patient

Task 7

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

14:40 Tue, 6 Dec

Google Lens

Czech → English

Task 9

In von Willebrand's disease, it is the first to be affected

Select one or more options:

- [ ] fibrin polymerization
- [ ] platelet degranulation
- [x] adhesion of platelets to subendinite collagen
- [ ] Prothrombin sctivation

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Autoimmune reactions against cell surface receptors

Select one or more options:

3. It is described as a reaction type III hypersensitivity

It can cause Graves' disease

It can cause myasthenia gravis

d. It is a typical cause of anaphylactic reaction

It is described as an action. hypersensitivity type V. which is a subtype of hypersensitivity reaction type rll

Abhina

Comment o

Viewers of this suggestions.

Comment

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

Google Lens

Czech → English

Task 3

Megaloblastic anemia

atrophic gastritis

Select one or more options:

- [ ] and is a consequence of Fe deficiency
- [ ] b arises as a result overproduction of hepcidin
- [x] accompanies atrophic gastritis
- [ ] d arises as a result of reduced production of hepcidin
- [x] It is the result of vitamin B12 and tetrahydrofolic acid deficiency

Abhina

Comment or a

Viewers of this fi
suggestions.

Comment

Task 4

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

14:36 Tue, 6 Dec

Google Lens

Czech → English

Which disorder of (1) primary hemostasis is most likely in patients with severe liver cirrhosis with congestive splenomegaly?

thrombocytopenia

Select one or more options:

- Immune thrombocytopenia
- Reduced or missing megakaryocytes in the bone marrow
- Thrombocytopenia due to platelet sequestration in an enlarged spleen
- Leukemia
- Hemolytic uremic Syndrome

Viewers suggest

Com

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

14:35 Tue, 6 Dec

Google Lens

Czech → English

Select the correct statement about hemophilia

Linked to X

A. Select one or more options:

- [ ] If the patient has approximately 10% Vill factor, this means a high physics of spontaneous bleeding
- [x] APTT prolonged
- [ ] This is a disease linked to the Y chromosome, which is why men can have it
- [ ] PT is teasing
- [ ] Typickel tou getechiatni cishi and purpura.

Task 9

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

14:34 Tue, 6 Dec

Google Lens

Czech → English

Select the correct statement(s) about hematopoietic stem cell transplantation

Select one or more options:

- [ ] a. The most common type of transplantation is xenotransplantation
- [ ] b. Peripheral blood can be a source of hematopoietic stem cells
- [x] High border compatibility is important for a successful transplant. HLA donor and recipient
- [ ] Kasthi marrow is the only source of hematopoietic stem cells
- [ ] Donor stem cells or their progeny can ignite recipient cells

Com

Viewers suggesti

Com

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

14:33 Tue, 6 Dec

Google Lens

Czech → English

A 55-year-old patient with a history of cirrhosis of the liver related to alcohol abuse is admitted to the hospital for multiple episodes of vomiting blood over a 2-hour period. This is the 3rd hospitalization with similar symptoms last year, previous episodes were related to Jicnu varices. The patient is pale, his arterial blood pressure is 60/30 mm Hg, pulse rate 150 beats per minute, respiratory rate 22/min, hemoglobin level 60 g/L. Select the correct statement(s) about the patient:

Select one or more options:

- [x] The patient probably has ascites
- [ ] b Cirrhosis of the liver may have developed after the first bleeding due to hypoxia of the liver
- [ ] Probably the lung bleeding is not related to cirrhosis of the liver
- [x] Treatment of a bleeding patient is complicated by possible disorders of hemostasis
- [ ] The patient is High Risk úmrtí

Abhinav Is

Comment or ad

Viewers of this file suggestions.

Comment

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

mood Czech → English

Task 8

Choose the incorrect statement

for iron deficiency

Select one or more options:

- a. *is accompanied by low transferrin saturation*
- b. it is accompanied by a high concentration of serum ferritin
- c. leads to microcytic and hypochromic anemil
- d. *can arise from frequent bleeding*

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

Google Lens

Czech → English

Task 6

In the pathogenesis of type 4 allergy apply

cytotoxic lymphocytes

Select one or more options:

- [x] a. cytotoxic lymphocytes ✓
- [ ] b. class antibodies (gM and lgG
- [ ] c. lgD and lgE class antibodies
- [ ] d. circulating immunocomplexes

penetrating the extravascular space

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

10:06 Mon, 5 Dec

Google Lens

Czech → English

02.CZ

2:02 p.m

moodle.upol.cz

Task 4

Protein C

Select one or more options:

- [x] after activation, it hydrolyzes accelerators (coagulation factors Va, FVIIIa)
- [x] is vitamin K dependent factor
- [ ] is activated by tissue factor
- [ ] dje acute phase protein
- [x] it is produced in the liver

Task 5

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Task 3

The most common cause of sideropenic anemia in our population is

Select one or more options:

- vegetarianism
- chronic bleeding
- malabsorpce
- d. lack of iron in diet

Abhinav Is

Comment or ad

Viewers of this file suggestions

Comment

Task 4

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

Google Lens

Czech → English

The von Willebrand factor is important

Select one or more options:

- [ ] Prevention of hyperactivation of protein C, thereby inhibiting fibrinolysis
- [x] Remembrance of the contact of platelets with collagen after damage to the vessel wall
- [ ] It reduces the activity of leukocytes during fibrinolysis
- [ ] Activation of antithrombin II, which prevents blood coagulation
- [ ] Protection of factor Vi from degradation in plasma

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Ulona 1

On the pathogenesis of hepatic encephalopathy

do not participate:

Select one or more options:

- [ ] a. Phenol
- [ ] b. Merkaptany
- [ ] c. Ammonia
- [ ] d. Uric acid

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Dilation of the pupils (mydriasis) and sweating are induced in anaphylactic shock

Select one or more options:

- a by the effect of leukotrienes LCT4 and LCD4
- b effect of histamine
- activation of the sympathetic nervous system
- d by the effect of bradykinin

Abhina

Comment or a

Viewers of this f suggestions.

Comment

Task 3

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

09:39 Mon, 5 Dec

Google Lens

Czech → English

It is characteristic of a type 2 allergic reaction

antibodies on surface cells

Select one or more options:

- [ ] a. polyclonal B-cell activation
- [x] b. the presence of antibodies on the surface of cells
- [ ] c. Tissue infiltration by eosinophilic granulocytes
- [ ] d. tissue infiltration by basophilic granulocytes

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

09:51 Mon, 5 Dec

Google Lens

Czech → English

Paroxysmal nocturnal hemoglobinuria (PNH)

Select one or more options:

- [ ] in the pathogenesis of their symptoms, the drop in blood pH during sleep and the activation of complement is applied
- [ ] it is caused by the lack of some protective > proteins on the erythrocyte membrane
- [ ] it arises as a result of excessive production of complement components
- [ ] it arises in the absence of spectrin in the membrane of Verytrocytes

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Task B

Parenchymal bleeding in
manifest hemophiliacs can be stopped

Select one or more options:

- [ ] Infusion of the missing coagulation factor
- [ ] b. long-term compression or tamponade.
- [ ] c transfusion of preserved blood
- [ ] and by the administration of blood clots plates

Abhin

Comment o

Viewers of this
suggestions.

Comment

Task 9

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

09:49 Mon, 5 Dec

Google Lens

Czech → English

Steatorhea

Select one or more options:

- [ ] a. It is characteristic of peptic ulcer
- [ ] it is common in patients with severe disorders of the exocrine function of the pancreas
- [ ] cje characteristic for idiopathic proctocolitis
- [ ] is characterized containing 5 g or more fat per 24 hours in stool
- [ ] is characteristic of malabsorption syndrome

Abhina

Comment o

Viewers of this suggestions.

Comment

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

09:46 Mon, 5 Dec

Google Lens

Czech → English

Utuna 9

The simultaneous decrease in plasma ferritin and rise in plasma transferrin is explainable

Select one or more options:

- [ ] a. a disorder of liver proteosynthesis
- [ ] b. inflammation
- [x] c. iron deficiency
- [ ] d. hemolysis

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

09:43 Mon, 5 Dec

Google Lens

Czech → English

Which of the following inflammatory mediators is most important in the pathogenesis of anaphylactic shock?

Select one or more options:

- a. Histamine
- b. Noradrenaline
- c. Somatostatin
- d. Serotonin
- e. Gastrin

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Prothrombin time can be used for diagnosis

Select one or more options:

- [ ] a. the number and function of platelets
- [ ] b. external pathways of the coagulation cascade (tissue factor pathway)
- [ ] c. internal pathways of the coagulation cascade
- [ ] liver function

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

Google Lens

Czech → English

It applies to ferritin

Select one or more options:

- [ ] and, it is not present in the blood
- [x] b. is a protein in which the storage form of iron is stored
- [x] when iron is sufficient, it is present in the reticuloendothelial system
- [ ] d. binds iron in ferroform

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Graft-versus-host reaction

Select one or more options:

- [x] and — extinct species 2 population can contribute to the eradication of the pathological clone of Leukemic cells
- [ ] b. is the reaction of the recipient against the transplanted organ
- [ ] c — can complicate autologous transplantation
- [x] can complicate allogeneic transplantation

Translated text

09:39 Mon, 5 Dec

Google Lens

Czech → English

Typical changes in coagulation screening tests in von Willebrand disease are (1):

bleeding – aPTT

Select one or more options:

- ☐ and. decreased Quick index and increased Duke bleeding time
- ☐ b. increased PT and decreased Duke bleeding time
- ☐ c. elevated aPTT and elevated D-dimery
- ☐ increased aPTT and increased bleeding time
- ☐ there is no answer correct

Abhinav Is

Comment or add

Viewers of this file c
suggestions.

Comment

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

09:32 Mon, 5 Dec

Google Lens

Czech → English

The pathogenesis of hepatic encephalopathy involves:

Select one or more options:

a. Endotoxin

b. Phenol

c. Mercaptans

d. Cytokininy

ALL

'pathogenesis'

Send to Translate Home | Copy text | Copy to computer | Search | Listen

Did you find these results useful? Yes No

Google Lens

Czech → English

Task 8

Complications of transfusion are not included

Select one or more options:

- [ ] a. risk of infection
- [ ] b risk of overloading the organism
- [x] risk of increased bleeding
- [ ] risk of lung damage

includes

Task 9

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

09:29 Mon, 5 Dec

Google Lens

Czech → English

Tissue factor

Select one or more options:

a. is an activator of fibrinolysis

b. is expressed under physiological conditions in the *endothelium*

initiates in a complex with negatively charged phospholipids the external pathway of activation of the plasma coagulation system

is expressed during inflammation

activated

monocytech

Abhin

Comment abou

Viewers of thi
suggestions.

Comment

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

Google Lens

Czech → English

Task 6

The Coombs test is used to determine

antibodies- RBC

Select one or more options:

- [x] autoantibodies against erythrocytes
- [ ] b. bleeding time
- [ ] c. absorption of vitamin B12
- [ ] d. levels of coagulation factors

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

09:28 Mon, 5 Dec

Google Lens

Czech → English

Task 4

Primary biliary cirrhosis is:

Select one or more options:

and disease from autoimunity

b. liver damage as a result of the action of chemotoxic agents

c. no answer is correct

d. Liver damage as a result of the effects of alcohol

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Tuesday 1

Coagulation factor XII deficiency

Select one or more options:

- [ ] manifests itself in a prolongation of the thrombin time
- [ ] is incompatible with life
- [ ] it's called hemophilia A
- [x] there are no speeches
- [ ] manifested by prolongation of the activated partial thromboplastin time (OPTT)

Abhinav J

Comment or a

Viewers of this fil
suggestions.

Comment

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Done

Task 1

Score out of 1.00

What are the most common hereditary hypocoagulation disorders (choose 1 answer)? Select one or more options :

a. Factor V mutation (Leyden)

→ hypercoagulation

b. Hemophilia A and von Willebrand disease

c. Thrombocytopenia associated with hypersplenism

d Coagulopathy related to cirrhosis

E. Thrombocytopenia associated with leukemia

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Task 2

Done

Score out of 1.00

Disorder of iron metabolism in chronic anemia

diseases is caused by (2)

Select one or more options:

a. increased production of hepcidin in the liver

b. accelerated transfer of iron from hepatocytes to bile

c. slowed iron export from macrophages

d. lack of iron in food

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

Google Lens

Czech → English

Task 5

Done

Score out of 1.00

Microcytic and/or hypochromic anemias

include:

Select one or more options:

a. Folic acid deficiency anemia

b. Sideroblastic anemia

c. Pernicious anemia

d. Iron deficiency anemia

e. Anemia due to lead poisoning

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

Google Lens

Czech → English

Task 7

Done

Score out of 1.00

When examining blood laboratories, in patients with moderate bacterial gram-positive infection (staphylococci, streptococci, pneumococci), it is usually assumed: Select one or more options:

a. Agranulocytosis

b. Lymphocytosis

c. Polycytémie

d. Eosinofilie

e. Neutrophia

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

Google Lens

Czech → English

Utoha 4

Which laboratory parameter is LEAST important for the diagnosis of disseminated intravascular syndrome coagulation?

Select one or more options:

- [ ] a. Prothrombin Time (PT)
- [ ] b. Albumin concentration
- [ ] c. D-dimery
- [ ] d. Fibrinogen
- [ ] e. Number of plates

Task 5

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Task 1

It is NOT among the etiological factors of liver cirrhosis

hepatitis A

Select one or more options:

- [ ] and hepatitis C
- [ ] b. Wilson's disease
- [ ] c. alpha-1-antitrypsin deficiency
- [ ] hepatitis A

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Task 2

The mean erythrocyte volume (MCV) is

in aplastic anemia

normal

Select one or more options:

- [ ] a. reduced
- [x] b. normal
- [ ] c. 70-83 femtoliters
- [ ] d. increased

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

Google Lens

Czech → English

Task 3

It does not include diseases

accompanied by eosinophilia

Select one or more options:

- [ ] and allergic conditions
- [ ] b. tetanus
- [ ] c. bronchial asthma
- [ ] d parasitic diseases

Abhina

Comment or

Viewers of this fi
suggestions.

Comment

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

16:09 Sun, 4 Dec

Google Lens

Czech → English

Can a boy have hemophilia if his parents do not have clinical manifestations of the disease?

Select one or more options:

- a. No, because it is a hereditary disorder
- b. Yes, because the severity of the disorder increases with each new generation
- c. Yes, because it can have a new mutation
- d. Yes, because the mother can be heterozygous and have no signs of hemophilia
- e. No, because both parents are healthy

Abhinav Jethi

Comment or add other

Viewers of this file can see suggestions.

Comment Cancel

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Task 8

Hemophilia A and B differ:

Select one or more options:

- [ ] and APTT over time
- [ ] The level of factor VIII and factor IX
- [ ] c. By way of inheritance
- [ ] d. PT or INR over time
- [ ] e. Main clinical symptoms

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

16:07 Sun, 4 Dec

Google Lens

Czech → English

Choose the correct statement(s).

Select one or more options:

- [ ] Bechterev's disease (ankylosing spondylitis) has a high association with a specific HLA type
- [ ] MHC II molecules are expressed on antigen presenting cells
- [ ] MHC I molecules are expressed on all nucleated somatic cells.
- [ ] d. MHC class 1 molecules are found on the surface of erythrocytes.
- [ ] MHC class 1 molecules are found only on the surface of cells

Imanit's system

Q+

Translated text


16:06 Sun, 4 Dec
Google Lens
Czech → English
Task 9
They contribute to the formation of ascites in live
failure
Select one or more options:
increased blood flow
in the splanchnic area
6. secondary disorder
aldosterone metabolism
c hyperbilirubinemia
d. relative hyperalbuminemia
during paradoxical dehydration
Complete the tour
Translated text
Send to Translate Home
Select all
Listen
Share
Download Czech
Did you find these results useful?
Yes
No

16:06 Sun, 4 Dec

Google Lens

Aplastic anemi Czech → English

Select one or more options:

and

It can develop with the administration

of certain antibiotics,

e.g. chloramphenicol

b. Patients do not need an infusion

treatment

c. Parvovirus B19 infection

accounts for 90-95

9% of severe aplastic anemia

It is mostly an acquired

disease that arises from

an unclear cause

During antitumor treatment,

it was mostly

transient!!

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

Google Lens

Czech → English

Choose the correct statement about jaundice

Select one or more options:

Hemolytic icterus can occur with extensive hematomas

b. The most common cause of jaundice in newborns is Crigler and Najjarüv syndrome

In posthepatalning Jaundice at all obstruction, the urine is dark and we demonstrate there urobilinogen and bilirubin

in Gilbert's syndrome unconjugated bilirubin predominates

U. hepatâlnino icterus is urine dark and we promote ram uroluinogen and bilirubin

Viewers suggesti

Comm

Comm

ranslated text

Google Lens

Czech → English

Task 3

Manifest sideropenic anemia will not manifest itself

Select one or more options:

- by lowering the level hemoglobin
- b. lowering the level of serum iron
- c. lowering the ferritin level
- macrocytosis

Task 4

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

After an acute respiratory infection of a 39-year-old male patient, the laboratory examination showed: 80*10^9 platelets, INR 1.1, aPPT 28 sec, Duke bleeding time 4 min 30 sec. The patient is most likely to:

Select one or more options:

- [ ] He acquired the disorder of hemophilia A due to the formation of antibodies against VIII
- [ ] b. Må vitamin K deficiency due to antibiotic treatment
- [ ] He must be hospitalized in the ICU for a high risk of life-threatening bleeding
- [ ] d. Is the mother heterozygous for hemophilia A or B
- [ ] Most likely, he has no clinical manifestations of hemostasis disorders

Com

Viewers

suggestio

Com

ranslated text

Google Lens

Czech → English

Task 1

Which of the following diseases has an autoimmune origin?

Select one or more options:

- [ ] and Hemophilia
- [x] Systemic lupus erythematosus
- [ ] c. Hashimoto's thyroiditis
- [ ] d. Down syndrome
- [x] Crohn's disease

Task 2

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

15:47 Sun, 4 Dec

Google Lens

Czech → English

Task 9

Hepatic encephalopathy

Select one or more options:

- [ ] and occurs in patients with hyperuricemia
- [ ] is a syndrome characterized by disorders of consciousness and neuromuscular activity
- [ ] venica in patients with hepatocellular failure
- [ ] dje syndrome characterized by impaired consciousness and hemolysis

Complete the tour

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

moodle.upol.cz

can

Google Lens

Czech → English

Task 8

Allogeneic bone marrow transplantation includes

related, unrelated

Select one or more options:

- [x] transplantation of cells of a relative
- [ ] b transplantation of own cells
- [ ] transplantation of cells from another animal species
- [x] transplantation of unrelated cells

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Oluh 7

Hereditary hemochromatosis is caused by iron absorption

Select one or more options:

- [x] excessive absorption of iron in the duodenum
- [ ] liver disorder with reduced transfer of iron into bile
- [ ] accelerated breakdown of erythrocytes
- [ ] d. urinary iron excretion

disorder

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

Google Lens

Czech → English

Task 6

Manifestations of disorders of primary hemostasis

do NOT include (1):

Select one or more options:

- [ ] a. Petechiae and ecchymoses
- [ ] b. Enteroragie
- [ ] Bleeding into the joints
- [ ] d. Hematuria
- [ ] E Epistaxis

Task 7

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

15:31 Sun, 4 Dec

Google Lens

Czech → English

Task 5

It does not belong to the manifestations of

decompensated liver cirrhosis

Select one or more options:

- [ ] a. dehydration
- [ ] b. jaundice
- [ ] c. encephalopathy
- [ ] d. prolonged prothrombin .time

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Utoha 4

Which laboratory parameter is LEAST important for the diagnosis of disseminated intravascular syndrome coagulation?

Select one or more options:

- [ ] a. Prothrombin Time (PT)
- [ ] b. Albumin concentration
- [ ] c. D-dimery
- [ ] d. Fibrinogen
- [ ] e. Number of plates

Albumin Concentration

Task 5

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Choose the correct statement(s)
about macrocytic anemia

Select one or more options:

a. Their symptom may be
subicterus

They can be caused
by chronic autoimmune
gastritis

c. They are usually caused
by Fe deficiency

May be caused by
ileal resection (due to
decreased Fe reabsorption)

They can be caused
by alcoholism

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

15:28 Sun, 4 Dec

Google Lens

Czech → English

Task 6

Arthus and Raynaud's phenomenon

are examples of an allergic reaction

Select one or more options:

- a. 1st type
- b. 3rd type
- c. 2 types
- d. 4. t PU

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful?

Yes No

Google Lens

Czech → English

Task 5

Select factors dependent on vitamin K:

Select one or more options:

- [ ] and III, IV
- [x] IX,X
- [x] Cas proteins
- [ ] d. heparin
- [ ] II, VII

9/10

2.7

2,7,9,10 Cf proteins

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Task 4

Acquired vasculopathies include:

Select one or more options:

- [ ] Glanzman thrombasthenia
- [ ] b. Marfan syndrome
- [x] Henoch-Schönlein purpura
- [x] Cushing's syndrome
- [ ] e. Ehlersův-Danlosův syndrome

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

15:22 Sun, 4 Dec
Utaha S

Google Lens

Czech → English

Select the correct statement(s) about hematopoietic stem cell transplantation:

Select one or more options:

- [x] It is an important method of treatment for Hodgkin's disease and Wiskott-Aldrich syndrome
- [x] Recipients require immunosuppressive drugs to reverse Graft versus Host Disease
- [x] Indications for transplantation include aplastic anemia, sickle cell anemia, thalassemil
- [ ] d. It is often performed in patients with sideropenic anemia
- [ ] It is most often performed in patients with lymphoproliferative disorders by kukemia disorders

Translated text

15:21 Sun, 4 Dec

Google Lens

Czech → English

These changes usually do NOT occur in liver failure

Select one or more options:

- [x] a. polyuria due to reduced aldosterone breakdown
- [ ] b bleeding disorders
- [ ] c gynecomastia in men
- [ ] d. hyperkinetic circulation

Task 2

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

15:10 Sun, 4 Dec

Google Lens

Czech → English

all Vodafone CZ

90%

moodle.upol.cz

Task 9

The principle of alleviating the symptoms of hepatic encephalopathy with lactulose is:

Select one or more options:

- [ ] a. reducing the production of toxic substances by inducing diarrhea
- [ ] D. no answer is correct
- [ ] cv prevention of asterixis manifestations
- [ ] in lowering the level of ammonia in the blood

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Task 8

It is not used in the pathogenesis of hepatic encephalopathy

Select one or more options:

- a. urea
- b. ammonia
- c portocaval anastomoses
- d. bleeding into the digestive tract

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Shift to the left in the differential

blood count

Select one or more options:

- [ ] and is present at vitamin B12 deficiency
- [ ] b indicates the presence of overripe blood elements
- [ ] which indicates the absence of progenitors of the beaten line
- [ ] it is observed in acute bacterial inflammation

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Task 6

Risk factors for the development of thromboembolic disease include

Select one or more options:

- [ ] and excess protein Ca S
- [ ] protein Ca S deficiency (C+S)
- [ ] prothrombin gene mutation
- [ ] increased HDL level cholesterol

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

15:05 Sun, 4 Dec

Google Lens

Czech → English

Prolongation of aPTT can induce

Select one or more options:

- [x] a. heparin therapy
- [ ] b. Factor Leyden mutation IN
- [x] c. factor XI deficiency
- [x] d. hemophilia A
- [x] e. DIC patients

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Sun, 4 Dec

Google Lens

Maybe 3

Czech → English

Choose the correct statement(s) about disorders of the immune system

Select one or more options:

- [x] a. Malnutrition can cause secondary immunodeficiency
- [ ] b. HIV infection primarily damages neutrophils
- [x] Bradykinin and leukotrienes are important components in the pathogenesis of a type hypersensitivity reaction
- [x] Eosinophilia is common in allergy
- [ ] e. Selective IgA deficiency is an example of secondary immunodeficiency

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

15:01 Sun, 4 Dec

Google Lens

Czech → English

Task 2

It does not apply to anemia

Select one or more options:

- [ ] a. with anemia, the content is reduced oxygen in the arterial curve
- [x] b. with anemia, the partial pressure of oxygen in the arterial blood is reduced
- [ ] c. to the anemic Petri syndrome, pallor, fatigue, decreased physical performance. gasping for damage, tachycardia
- [ ] d. is defined as discounted antu circulating Bytrocyte or Vencentrade hemoglobin

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

15:01 Sun, 4 Dec

Google Lens

Czech → English

A decrease in the plasma concentration of haptoglobin is indicative of

Select one or more options:

- [x] and. iron deficiency
- [ ] b. excess iron
- [ ] Cextravascular hemolysis
- [x] intravascular hemolysis

haptoglobin decrease = intravascular hemolysis

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Task 2

Inefficient hematopoiesis (inefficient erythropoiesis) is part of the pathogenesis Select one or more options:

- [x] myelodysplastic syndrome
- [ ] neonatal anemia anemia
- [ ] cine deficiency prythropoietin
- [x] megaloblastic anemia

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

Google Lens

Czech → English

Choose the correct statement about

hypersensitivity reactions

Select one or more options:

AND type hypersensitivity reaction is the cause of bronchial asthma

B. for the mechanism The development of glomerulonephritis often results in a hypersensitivity type reaction

Systemic lupus erythematosus is a type (1) hypersensitivity reaction

D Weakness of the hypersensitivity of the pool is associated with the level of the moose

E intermediate reaction type e sensitive cells

Abhinav Jeth

Comment or add

Viewers of this file c suggestions.

Comment

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

14:57 Sun, 4 Dec

Google Lens

Czech → English

Task 3

Leukemoid reaction is

increased WBC

Select one or more options:

- [x] a. reactive state se increased number leukocytes and immature forms in peripheral blood
- [ ] b. the initial stage of leukemia
- [ ] c. end stage chronic myeloid leukemia
- [ ] d. characterized by a very low activity of alkaline phosphatase in granulocytes

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

14:56 Sun, 4 Dec

Google Lens

Czech → English

Task 9

An increase in reticulocytes above the upper physiological limit of 1.5% occurs (2)

Select one or more options

hemolytic anemias
hypersplenism

- [ ] and. in myelodysplastic syndrome
- [ ] b in aplastic anemia after transfusion erythrocyte masses
- [x] in hemolytic anemias
- [x] due to hypersplenism

Abhina

Comment o

Viewers of this suggestions.

Comment

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No

14:55 Sun, 4 Dec

Google Lens

Czech → English

Task 1

Methemoglobin

Select one or more options:

a. is hemoglobin's defective glob hov retexci

b. contains trivalent iron

c. occurs during hypoxia

4. contains ferrous iron

Task 2

Translated text

Send to Translate Home | Select all | Listen | Share | Download Czech

Did you find these results useful? Yes No