Task 10

Hypochloremic metabolic alkalosis

Select one or more options:

- [ ] a. It develops in patients with diarrhea
- [x] b. It develops in patients with vomiting ✔
- [ ] c. It develops in patients with a NaCl intake of 5-8 g / day
- [ ] d. It develops in patients with reduced absorption of HCO3- in the proximal tubules of the kidneys
- [ ] e. It develops in patients with long-term starvation

The correct answer is: It develops in patients with vomiting.

The intrinsic factor complex with cobalamin (vitamin B12) is absorbed

Select one or more options:

- [ ] a. by paracellular transport in the jejunum and ileum
- [ ] b. facilitated diffusion in the ileum
- [x] c. using receptors in the area of the brush border of the ileum
- [ ] d. in the cells of the gastric pylorus

27% 5:36 p.m

Task 5 Points 0.50/1.00

Choose the correct statement about stress starvation

Select one or more options:

- [ ] a. Ketone bodies are often elevated
- [ ] b. Severe emotional stress (e.g. the death of a partner) is one of the most important causes
- [ ] c. It is characterized by intense catabolism of proteins
- [x] d. Glucose level often increases✓
- [ ] e. Insulin levels are usually decreased

The correct answers are:
It is characterized by intense catabolism of proteins,
Glucose level often increases

71%
Sesbijpg.2-490
11:29
Google Lens
צ'כית ← אנגלית
jpg.2-485
jpg.2-484
jpg.2-490M
jpg.2-489
jpg.2-488
jpg.2-495
jpg.2-494
jpg.2-493 M
02.CZ 11:26 a.m 91%
moodle.upol.cz

Task 5

The following statements may be true about hypokalemia

Select one or more options:

- a. may be related to high aldosterone secretion
- b. may be associated with metabolic acidosis
- c. can even cause paralytic ileus
- d. can occur in heart failure with significant swelling
- e. a tall spiked T wave is prominent on the EKG

תרגום
שליחה לדף הבית של Translate
בחירת הכול
האזנה
האם התוצאות האלה מועילות?
כן
לא

You doing good. U R our Iron dome

Task 5 Points 0.00 / 1.00

Choose the correct statement about simple starvation:

Select one or more options:

- [x] a. During the first 10-12 hours, the body receives energy mainly due to lipolysis X
- [ ] b. On day 14, the patient will likely have metabolic alkalosis due to proteolysis
- [ ] c. Approximately 4-5 kilograms of body muscle will be lost during the first 7 days of fasting
- [x] d. After a few days, energy metabolism decreases✓
- [ ] e. Adult patients can usually survive for more than 4 weeks if they consume only water

The correct answers are: Adult patients can usually survive for more than 4 weeks if they consume only water After a few days energy metabolism decreases

Task 6 Points 0.25/1.00

Select conditions associated with hypercholesterolemia

Select one or more options:

- ✓a. Diabetes mellitus type II✓
- [ ] b. Hypothyroidism
- [ ] c. Anorexia nervosa
- [ ] d. Nephrotic syndrome
- [ ] e. Hepatocellular carcinoma with cachexia

The correct answers are: Diabetes mellitus type II, Anorexia nervosa, Nephrotic syndrome, Hypothyroidism

moodle.vpol.cz

facebook

Task 1000.50/1.00

Chicken with hepatomas are characterized by:

Select one or more options:

- An enlarged liver with fatty infiltrates
- A protruding flash ✓
- Low resistance to infections ✓
- Low number of eggs laid
- Pitting edema with swelling of ankles and feet

The correct answers are: Protruding flash, Low resistance to infections, Pitting edema with swelling of ankles and feet

Seiversi Pravichuanosti developed metabolic alkalosis due to Silance

stomach and intestines. Probezny test 22 (contents of macadan cut down from prednasek uzvracenest & 2-5.11. 2020, online

15.30 e. The patient probably does not have acid-base balance disorders

The correct answer is: Most likely, the patient has a combination of metabolic acidosis and metabolic alkalosis.

Problem 4 Points 0.00 / 1.00

Choose the correct statement about simple starvation:

choose one or more a. possibility:

During the first For 10-12 hours, the body receives energy mainly thanks to lipolysis

The 14th day the patient probably has metabolic alkalosis due to the proteolysis of

will be After a few days, the energy metabolism decreases ✓

d. During approximately 4-5 Ce are lost during the first 7 days X kilograms of body muscle

of fasting Adult patients can usually survive for more than 4 weeks if they consume only water

The correct answers are: Adult patients can usually survive for more than 4 weeks if they consume only water After a few days energy metabolism decreases

2020-ZS-PFY/VCA11-Binding full story practical

e. Adult patients can usually survive for more than 4 weeks if they consume only water

The correct answers are: Adult patients can usually survive for more than 4 weeks if they consume only water After a few days, energy metabolism slows down

Task 5 Number of points out of 1.00

What are the clinical manifestations of hypokalemia in skeletal muscle, heart and GIT?

on the heart - leads to bradycardia and arrhythmias, on the EKG I observe - a flat T wave, a U wave appears, widened QRS GIT - there is reduced peristalsis, it can lead to paralytic ileus

muscle wasting, weakness,

4:06 p.m

Select the typical symptoms of vitamin deficiency:

Select one or more options:

- → Lack of vitamin A leads to night blindness, xerophthalmia, keratinization of the skin and mucous membranes✓✓
- b. Vitamin D deficiency increases the risk of mood swings and depression
- c. Vitamin C deficiency significantly increases the risk of thrombosis
- d. Lack of vitamin D increases the risk of muscle weakness, muscle spasms
- e. Long-term use of broad-spectrum antibiotics can lead to vitamin K deficiency and an increased risk of bleeding

The correct answers are: Vitamin A deficiency leads to night blindness, xerophthalmia, keratinization of the skin and mucous membranes, Vitamin D deficiency increases the risk of muscle weakness, muscle spasms, Vitamin D deficiency increases the risk of mood swings and depression, Long-term use of broad-spectrum antibiotics can lead to vitamin K deficiency and to increase the risk of bleeding

2020-ZS.PEY/VCA11-17: Pathophysiology of esophagus, stomach and intestines: Continuous test & 2 (content of practical exercises 4-6 and lectures no. 4-6) The correct answer is: The patient has a high risk of developing respiratory acidosis due to hypoventilation.

Problem 3 Points 1.00/1.00

Major vomiting will lead to

Select one or more options:

- a. metabolic alkalosis due to compensatory hyperkalemia
- b. metabolic acidosis due to large losses of chlorides
- c metabolic acidosis due to compensatory formation of HCL
- metabolic alkalosis due to a large loss of hydrogen ions in the acidic gastric juice ✓
- Hypochloremia ✓

The correct answers are: metabolic alkalosis due to a large loss of hydrogen ions in the acid gastric state, Hypochlorémil

Task 4 Score out of 1.00

4:05 p.m.

Task 8

0.00/1.00 points

Intermittent fasting can mean that a person:

Select one or more options:

- [ ] a. Consumes only 1 type of food each week (e.g. 1 week apples, 2 week bread, 3 week cheese...)
- [ ] b. Consumes a low-protein diet for 1 day, then a low-fat diet for 1 day, then a low-carb diet for 1 day X
- [ ] c. Consumes food for 6 hours a day and does not consume food for 18 hours
- [ ] d Consumes food for 8 hours a day and does not consume food for 16 hours
- [ ] e Consumes only 1 type of food each day (e.g. apples 1 day, bread the next day, cheese 3 days...)

The correct answers are: Consumes food for 8 hours a day and does not consume food for 16 hours, Consumes food for 6 hours a day and does not consume food for 18 hours

Task 2/LED

A 12-year-old boy developed decompensated type 1 diabetes mellitus due to an insufficient dose of insulin. His glucose level is 20.9 mmol/L. Blood pH is 7.1. the patient is severely dehydrated. Select all correct statements about the patient:

Select one or more options:

- a. Most likely, the patient has tachycardia due to dehydration and activation of the sympathetic nervous system
- b. 5 most likely the patient's blood osmolarity is high
- c. Most likely, the patient has a normal or decreased anion gap
- d. Most likely, the patient is dehydrated due to osmotic diuresis ✓
- e. Most likely, the patient has an elevated level of ketone bodies in the plasma ✓

The correct answers are: Most likely the patient has tachycardia as a result of dehydration and activation of the sympathetic nervous system, most likely the patient has dehydration as a result of osmotic diuresis, 5 most likely the patient's blood osmolarity is high, Most likely the patient has an increased level of ketone bodies in the plasma

Task 3 from 100/100

What are the clinical manifestations of hypokalemia in the bones, muscles, heart and GIT?

hypokalemia in KS leads to muscle weakness and can also cause rhabdomyolysis.

VGIT causes constipation and paralytic ileus

Heart: prolongs repolarization (action potential), which manifests itself in a prolonged PT-interval, presence of wave U , and leads it karytmim.

search term

Problem 9 Points 0.33/1.00

Choose the correct statement(s) about hyperkalemia

Select one or more options:

- and Associated with hyperaldosteronism
- b. It is a side effect of insulin therapy
- LC May cause arrhythmia
- May cause paresthesia and muscle weakness
- It is typical for patients with end-stage renal insufficiency ✓

The correct answers are: Ie typical for patients with end-stage renal insufficiency. May cause paresthesia and muscle weakness, May cause arrhythmia

Problem 10 Points 1.00/1.00

Ztráty vody a of solutes during excessive sweating lead to

Sepetendi along lecture no. 4-6). AND

e. The patient is most likely hypokalemic

The correct answers are: The patient most likely has proximal tubular acidosis The patient most likely has hypokalemia The urine HCO3- level is most likely to be high

Task 2 Points 0.00 / 1.00

A 12-year-old boy developed decompensated type 1 diabetes mellitus due to an insufficient dose of insulin. His glucose level is 20.9 mmol/L, blood pH is 7.1, and the patient is severely dehydrated. Select all correct statements about the patient:

Select one or more options:

- The patient most likely has a normal or reduced anion gap X
- b. The patient is most likely dehydrated due to osmotic diuresis
- Most likely, the patients blood osmolarity is high ✓
- d. Most likely the patient has tachycardia due to dehydration and activation of the sympathetic nervous system
- Most likely, the patient has an elevated level of ketone bodies in the plasma ✓

The correct answers are: Most likely the patient has tachycardia due to dehydration and activation of the sympathetic nervous system system, Most likely the patient is dehydrated due to osmotic diuresis, Most likely the blood osmolarity patients high. Most likely, the patient has an elevated level of ketone bodies in the plasma

Task 9 Number of points out of 1.00

Give 4 examples of swelling caused by increased hydrostatic pressure.

right heart weakness

venous insufficiency: varices on DKK, immobilization, myopathy, muscle paralysis

venous obstruction: phlehothrombosis, oppression by tumor

Task 10 Score out of 1.00

Tissue lactate acidosis may NOT be caused

Select one or more options:

sepsi s MODS

hypoglycaemia

binding of lactate metabolism in the liver

lack of oxygen in the tissues from circulatory causes

lack of oxygen in respiratory diseases

given expression

Hypothyroidism ✓

e. Type II diabetes mellitus ✓

The correct answers are: Diabetes mellitus type II, Anorexia nervosa, Nephrotic syndrome, Hypothyroidism

Problem 7 Points 1.00 / 1.00

Hypochloremia is seen in

Select one or more options:

edematous conditions associated with dilutional hyponatremia ✓

6. excessive losses of HCO3- from the organism (GIT, kidneys)

c. isotonic dehydration

d. hypertonic dehydration

metabolic alkalosis during vomiting✓

The correct answers are: edematous states associated with dilutional hyponatremia, metabolic alkalosis with vomiting

14:17 LTE+ 91
moodle.upol.cz/mod/quiz/re 5 :

Task 7

0.00 / 1.00 points

Choose the correct statement about stress starvation

Select one or more options:

- [ ] a. Glucose level is often elevated
- [x] b. Severe emotional stress (e.g. the death of a partner) is one of the most important causes of X
- [x] C. Insulin levels are usually decreased X
- [x] d. It is characterized by intensive protein catabolism✓
- [ ] e. Ketone bodies are often elevated

The correct answers are: It is characterized by intense catabolism of proteins, Glucose level often increases

Task 8

0.50 / 1.00 points

Task 3

Choose the correct statement(s) about iron:

Select one or more options:

- [x] a. Microcytic anemia, dysphagia, atrophic gastritis can be manifestations of iron deficiency ✔
- [x] b. Iron deficiency often occurs in patients with chronic blood loss ✔
- [ ] c. Iron can accumulate in the cells of the pancreas, liver, heart in case of multiple transfusions
- [ ] d. Iron overload never happens
- [ ] e. Iron is absorbed mostly in the terminal (distal) ileum

The correct answers are: Iron can accumulate in the cells of the pancreas, liver, heart in case of multiple transfusions, Iron deficiency often occurs in patients with chronic blood loss, Microcytic anemia, dysphagia, atrophic gastritis can be manifestations of iron deficiency

University of Palacký UISU X

review.php?attempt=1840248&cmid=337462

2020-25 Area iyi

fiu and stew Weight test 2 (content of practical exercises 46ed4/ wed 22-5.11.2008 online 15:50

A 52-year-old patient, a chronic alcoholic, is admitted to the hospital with signs of liver cirrhosis, acute alcohol intoxication, and vomiting. His albumin level is 10 g (N-35-50 g/L), pH 7.38, CL-90 mmol/L, PaCO2-5.5 kPa (40 mm Hg). Choose the 1 most appropriate statement about the patient:

Select one or more options: The

- patient probably does not have acid-base balance disorders
- Most likely, the patient developed metabolic alkalosis due to vomiting
- Based on the available information, no conclusion can be made about the acid-base balance of patient X
- DdS most likely the patient has a combination of metabolic acidosis and respiratory alkalosis
- OS. the patient most likely has a combination of metabolic acidosis and metabolic alkalosis

The correct answer is: 5 most likely the patient has a combination of metabolic acids and metabolic alkaloids

Task 250/1.00

A 12-year-old boy developed type 1 diabetes mellitus decompensation due to an insufficient dose of insulin. His glucose level is 20.9 mmol/L, the pH of coffee is 7.1, the patient is severely dehydrated. Select all correct statements about the patient

Select one or more options:

- Most likely, the patient has tachycardia due to dehydration and activation of the sympathetic nervous system
- Most likely, the patient's blood osmolarity is high
- Most likely, the patient has a normal or reduced anion gap
- 5 most likely the patient has dehydration due to osmotic dimly✓
- The patient has an elevated level of cartilaginous cells ✓

Problem 2 Under 1.00

A 12 year-old boy developed decompensated type 1 diabetes mellitus due to an insufficient dose of insulin. His glucose level is 19 mmol/L, blood pH is 7.1. The patient has fluid dehydration. Select all the correct statements about the patient

Select one or more options

- Most likely, the patient has tachycardia as a result of dehydration and activation of the sympathetic nervous system
- Most likely, the patient's blood osmolarity is high
- Most likely, the patient has a normal or reduced anion limit
- The patient is most likely dehydrated due to osmotic diuresis
- The patient probably has an increased die of ketones in the flame

Task 3

as the roots are manifestations of hypokalemia by the bone of the ruling stock, on the GIT)

and sits down

Task 8

Choose the correct statement(s) about vitamin B12:

Select one or more options:

- [ ] a. The recommended daily intake of B12 is 70 - 150 mg for adults
- [x] b. Vitamin B12 deficiency leads to hyperhomocysteinemia with an increased risk of developing endothelial damage ✔
- [ ] c. B12 is mostly reabsorbed in the duodenum
- [x] d. Vitamin B12 deficiency can lead to macrocytic anemia and nerve demyelination ✔
- [ ] e. Gastric resection leads to B12 reabsorption disorders

The correct answers are: Gastric resection leads to B12 reabsorption disorders, Vitamin B12 deficiency leads to hyperhomocysteinemia with an increased risk of developing endothelial damage, Vitamin B12 deficiency can lead to macrocytic anemia and nerve demyelination

Task 10

Hypochloremic metabolic alkalosis

Select one or more options:

- [ ] a. I develop in patients with diarrhea
- [x] b. It develops in patients with vomiting
- [ ] c. It develops in patients with a NaCl intake of 5-8 g/day
- [ ] d. It develops in patients with reduced absorption of HCO_3^- in the proximal tubules of the kidneys
- [ ] e I develop in patients with prolonged starvation

The correct answer is: I develop in patients with vomiting.

Task 5

Under what pathological conditions (4) can hypernatremia occur?

Increased supply of sodium -enterally,

increased RAA activity

renal insufficiency - reduced excretion

increased supply of physio, solution

He commented:

Task 6

The following characteristics are not used for the classification of edema:

Select one or more options:

- [ ] a. Acquired and [illegible]
- [ ] b. Hyperosmolar and hypoosmolar
- [ ] c. Local and generalized
- [ ] d. Primary and secondary lymphedema
- [ ] e. Pitting a non-pitting

The correct answers are: Hyperosmolar and hypoosmolal, Acquired and inherited

14:17 LTE+ 91

The correct answer is: The patient is at high risk of developing respiratory acidosis due to hypoventilation.

Task 6

0.00 / 1.00 points

The following characteristics are NOT used to classify edema:

Select one or more options:

- [x] a. Pitting a non-pitting ✗
- [x] b. Acquired and inherited✓
- [ ] c. Hyperosmolar and hypoosmolar
- [ ] d. Local and generalized
- [ ] e. Primary and secondary Lymphedema

The correct answers are: Hyperosmolar and hypoosmolar, Acquired and inherited

Task 7

0.00 / 1.00 points

Choose the correct statement(s) about ulcer disease:

Select one or more options:

- [x] A. Increased amounts of *pepsinogen* or proteinases are present in many patients with gastric ulcers AND
- [x] B. Nonsteroidal anti-inflammatory drugs are significant risk factors for ulcer formation
- [ ] C. The influence of the autonomic nervous system on the development of ulcers is negligible
- [ ] D. Helicobacter pylori is an important factor in the development of duodenal ulcer, but does not play a role in the development of gastric ulcers
- [x] E. Increased production of HCL is a significant factor in the development of duodenal ulcers

10:461 53

Task 9

Nonsteroidal antirheumatic drugs (e.g. indomethacin, acetylsalicylic acid) have an ulcerogenic effect by

Select one or more options:

- [x] a. they reduce bicarbonate secretion
- [x] b. they cause inhibition of prostaglandin formation with subsequent reduced quantity and quality of mucus
- [ ] c. they increase the production of prostaglandins
- [ ] d. they activate proteinases
- [x] e. they penetrate by diffusion into the cells of the gastric mucosa

Complete the tour

Choose the correct statement(s) about ulcer disease:

Select one or more options:

- [x] AND Increased amounts of pepsinogen or proteinases are present in many patients with gastric ulcers
- [x] 3. Nonsteroidal anti-inflammatory drugs are significant risk factors for ulcer formation
- [ ] C. The influence of the autonomic nervous system on the development of ulcers is negligible
- [ ] D. Helicobacter pylori is a significant factor in the development of duodenal ulcer, but does not play a role in the development of gastric ulcers
- [x] E. Increased HCL production is a significant factor in the development of duodenal ulcers

12:35 p.m 71%
Test No. 2 2021: N...
moodle.upol.cz

e. bicarbonates pass from plasma to erythrocytes

Task 4

Causes of hypokalemia include:

Select one or more options:

- Rotavirus infection of the GIT in infants with diarrhea and vomiting
- b. Adrenal cortex failure
- c. Hypoaldosteronismus
- Hyperaldosteronismus
- e. Advanced renal failure

12:34 p.m 71%

Task 2

Malabsorption syndrome

Select one or more options:

- [ ] and. it is usually accompanied by a tendency to develop metabolic alkalosis as a result of diarrhea
- [x] b. it is usually accompanied by diarrhea
- [ ] c. It is most often related to stomach disorders
- [x] d. may present with hypochromic or megaloblastic anemia
- [ ] E. it is usually accompanied by obstinate constipation

12:35 p.m 71%1
X Test No. 2 2021: N...
moodle.upol.cz

Task 5

Complications of peptic ulcer disease include:

Select one or more options:

- [x] A. Anemia due to occult (hidden) bleeding
- [ ] B. Dehydration due to frequent vomiting
- [ ] C. Chronic pancreatitis
- [x] D. Stomach cancer
- [x] E. Perforation

Palpation is like saying

thermometer or stethoscope

12:35 pm 4G 71%

X Test No. 2 2021: N...

moodle.upol.cz

e. bicarbonates pass from plasma to erythrocytes

Task 4

Causes of hypokalemia include:

Select one or more options:

a. Rotavirus infection of the GIT in infants with diarrhea and vomiting

b. Adrenal cortex failure

c. Hypoaldosteronismus

d. Hyperaldosteronismus

e. Advanced renal failure

d. is influenced by the current state of renal function

Task 8

It poses a fatal risk in patients with mechanical ileus

Select one or more options:

a. weight loss

b. resorption of bacterial toxins

c. Bile secretion disorder

d. hyperhydratace

e. dehydration

12:36 p.m 71%

Task 7

It applies to potassium

Select one or more options:

- [ ] and decreases with tissue breakdown.

burns, hemolysis

- [ ] b. in metabolic

alkalosis, potassium is

released from binding to

intracellular phosphates

and proteins

- [x] which is dependent

on a

sufficient energy supply for

the proper function of

the Na+/K+ ATPaza

- [x] is influenced by the

current state of renal function

T-Mobile CZ 22:23 15%
moodle.upol.cz

Task 2

The most typical disorder of lower esophageal sphincter relaxation caused by loss of nerve cells in the esophagus is called (choose 1 best answer)

Select one or more options:

- [ ] a. Esophageal varices
- [ ] b. Peptic ulcer
- [ ] C. Gastroesophageal Reflux Disease (GERD)
- [x] d. Achalasie
- [ ] e. Barrett's esophagus

Task 3

moodle.upol.cz

Task 5000/1.00

Tissue lactic acidosis may NOT be caused

Select one or more options:

- a. binding of lactate metabolism in the liver
- b. lack of oxygen in respiratory diseases
- c. hypoglykemii
- d. sepsi s MODS
- e. lack of oxygen in the tissues from circulatory causes ✗

The correct answer is hypoglykemii

Task 600/100

Anion gap is calculated from plasma concentration values

Select one or more options:

- [Na] [K]-[HCO3 -]
- (Na+) [K][Ca] [HCO3-]-[C-]
- [Na+] + [K+] - [HCO3-]

Task 5 Points 0.67/1.00

Choose the correct statement(s) about SIADH

Select one or more options:

- a. Swelling of the lower extremities is common
- b. One of the causes of SIADH is pneumonia
- In case of incorrect treatment, pontine demyelinosis can occur✓✓
- May cause hyponatremia✓✓
- It is associated with a reduced level of ANP

The correct answers are: It can cause hyponatremil, In the case of incorrect technique, it can lead to pontine demyelination. One of the causes of SIADH is pneumonia

Task 6 0.30/1:00

what meta regular menstruation from 12 to 13 years

Problem 7 Points 0.00 / 1.00

A 2.5-year-old girl developed diarrhea, most likely due to a rotavirus infection. Her temperature is 37.8 °C, during the examination she sweats, her skin is pink. She vomited several times and had diarrhea 5-6 times every day for 2 days, her heart rate is 130 beats per minute (N 80-130 beats per minute), arterial blood pressure 100/65 mm Hg (still normal), responds to simple questions and communicates with the doctor and her mother. Choose the 1 most appropriate statement about the best way to treat the patient:

Select one or more options:

- and Needs immediate hospitalization with 0.9% NaCl infusion therapy
- b. Needs immediate hospitalization with infusion therapy with 5% glucose solution
- It can be treated at home with advice to drink more sweet tea and X fruit juice
- *It can be treated at home with advice to drink oral rehydration drinks*
- He needs immediate hospitalization, but infusions cannot be given until the doctor receives the results of the laboratory test.

The correct answer is: It can be treated with the recommendation of drinking oral rehydration drinks.

T-Mobile CZ 22:23 15%
moodle.upol.cz

Task 3

An acute form of intestinal pseudo-obstruction can NOT occur as a consequence

Select one or more options:

- [ ] a. myocardial infarction
- [ ] b. surgical procedure
- [ ] c. potassium depletion
- [x] d. sodium depletion

Task 4

Hyperchloremická acidóza

T-Mobile CZ 22:23 15%
moodle.upol.cz

Choose the correct statement(s) about simple starvation:

Select one or more options:

- [ ] a. During the first 10-12 hours, the body *receives energy mainly* thanks to lipolysis
- [x] b. Adult patients can usually survive for more than 3 weeks if they consume only water
- [x] c. After a few days, the energy metabolism decreases
- [ ] d. During the first 7 days of fasting, approximately 5-8 kilograms of body muscle will be lost
- [ ] e. On day 14, the patient will likely have metabolic alkalosis due to proteolysis

T-Mobile CZ 22:23 15%
moodle.upol.cz

Task 1

Causes of increased volume of interstitial fluid and edema formation

include:

Select one or more options:

- and. reduced filtration pressure in the arterial part of the capillary
- b. increased oncotic pressure in *the capillary*
- c. action of histamines and related substances
- d. insufficient lymphatic circulation

T-Mobile CZ 10:24 p.m 15%
moodle.upol.cz

Task 9

Respiratory acidosis

is characterized

Select one or more options:

- [ ] a. rise in pCO2
- [ ] b. a decrease in pCO2
- [ ] c. May develop due to hyperventilation under stress
- [ ] d. increased excretion of K+ in the urine
- [ ] e. increased excretion of HCO3- into the urine

Complete the tour

T-Mobile CZ 22:23 15%
moodle.upol.cz

Task 7

Choose the correct statements about gastroesophageal reflux disease:

Select one or more options:

- [ ] A. often leads to achalasia
- [x] B. high prevalence in the adult population of developed countries
- [x] C. can be treated by inhibiting gastric juice secretion
- [ ] D. the highest incidence is in newborns and leads to pylorospasm
- [x] E. associated with this is Barrett's esophagus

T-Mobile CZ 22:23 15%
moodle.upol.cz

Task 7

Choose the correct statements about gastroesophageal reflux disease:

Select one or more options:

- [ ] A. often leads to achalázia
- [x] B. high prevalence in the adult population of developed countries
- [x] C. can be treated by inhibiting the secretion of gastric juice
- [ ] D. the highest incidence is in newborns and leads to pylorospasm
- [x] E. Barrett's jicen is associated with it

T-Mobile CZ 22:23 15%
moodle.upol.cz

Choose the correct statement(s) about simple starvation:

Select one or more options:

- [ ] a. During the first 10-12 hours, the body receives energy mainly thanks to lipolysis
- [x] b. Adult patients can usually survive for more than 3 weeks if they consume only water
- [x] c. After a few days, the energy metabolism decreases
- [ ] d. During the first 7 days of fasting, approximately 5-8 kilograms of body muscle will be lost
- [ ] e. On day 14, the patient will likely have metabolic alkalosis due to proteolysis

T-Mobile CZ 10:24 p.m. 15%
moodle.upol.cz

Task 9

Respiratory acidosis

is characterized

Select one or more options:

- [ ] by rising pCO_2
- [ ] b. a decrease in pCO_2
- [ ] c. May develop due to hyperventilation under stress
- [ ] d. increased excretion of K^+ in the urine
- [ ] e. increased excretion of HCO_3^- into the urine

Complete the tour

T-Mobile CZ 22:23 15%
moodle.upol.cz

Task 8

Major diarrhea will lead to

Select one or more options:

- [ ] a. metabolic acidosis due to compensatory formation of HCL
- [ ] b. metabolic alkalosis due to bicarbonate loss
- [ ] c. metabolic alkalosis due to compensatory hyperkalemia
- [x] d. metabolic acidosis due to bicarbonate loss
- [x] e. dehydration

Task 9

Nonsteroidal antirheumatic drugs (e.g. indomethacin, acetylsalicylic acid) have an ulcerogenic effect by

Select one or more options:

- [x] they reduce bicarbonate secretion
- [x] they cause inhibition of prostaglandin formation with subsequent reduced quantity and quality of mucus
- [ ] c. they increase the production of prostaglandins
- [ ] d. they activate proteinases
- [x] penetrate the cells of the gastric mucosa by diffusion

Complete the tour

T-Mobile CZ 22:23 15%
moodle.upol.cz

Task 8

Major diarrhea will lead to

Select one or more options:

- [ ] a. metabolic acidosis due to compensatory formation of HCL
- [ ] b. metabolic alkalosis due to bicarbonate loss
- [ ] c. metabolic alkalosis due to compensatory hyperkalemia
- [x] d. metabolic acidosis due to bicarbonate loss
- [ ] e. dehydration

4:06 p.m 零路 83%

Task 8

Constipation

Select one or more options:

- [ ] a. can occur with hyponatremia
- [ ] b. arises in hyperthyroidism
- [x] c. can occur during dehydration
- [x] d. is a negative factor that may contribute to portosystemic encephalopathy in advanced liver cirrhosis
- [x] e. risk factors include older age

||| ○ <

4:04 p.m AND. 83%

Task 2

Acute depletion of extracellular fluid volume by diuretic therapy (non-carbonic anhydrase-inhibiting diuretics) leading to loss of water and NaCl

Select one or more options:

- [ ] a. does not affect the acid-base balance
- [ ] b. can cause acidosis from dehydration
- [ ] c. leads to bicarbonate losses by the kidneys
- [x] d. may cause alkalosis from dehydration

Task 3

4:04 p.m 83%

Task 2

Acute reduction of *extracellular fluid volume by diuretic* therapy (non-carbonic anhydrase-inhibiting diuretics) leading to water and NaCl losses

Select one or more options:

- [ ] axoneme affects the acid-base balance
- [ ] b. can cause acidosis from dehydration
- [ ] c. leads to bicarbonate losses through the kidneys
- [x] d. can cause alkalosis from dehydration

Task 3

16:04 83%

Task 1

Hirschsprung's disease
(megacolon congenitum)

Select one or more options:

- [x] and is a congenital disease in which ganglion cells in the intestinal plexus are missing
- [x] b. is manifested by diarrhea due to insufficient digestion of fats
- [ ] c. is manifested by organic constipation due to stagnation of intestinal contents
- [ ] d. is an intestinal manifestation of congenital progressive myodystrophy

III ○ <

11:41

Google Lens

02-SK

Czech → English

Task 9

Point 0.67/1.00

Choose the correct statement/s about SIADH

Select one or more options:

- alt is associated with a reduced level of ANP
- bIn case of incorrect treatment, it can lead to pontine demyelination ✓
- May cause hyponatremia ✓
- d.Swelling of the lower extremities is common
- eOne of the causes of SIADH is pneumonia

The correct answers are: It can cause hyponatremia, In case of incorrect treatment it can lead to pontine demyelination, One of the causes of SIADH is pneumonia

Translated text

Send to Translate Home

Select all

Listen

Are these results useful?

YES

NO

11:41

Google Lens

02-SK

Czech → English

Task 9

Point 0.67/1.00

Choose the correct statement/s about SIADH

Select one or more options:

- alt is associated with a reduced level of ANP
- bIn case of incorrect treatment, it can lead to pontine demyelination ✓
- May cause hyponatremia ✓
- d.Swelling of the lower extremities is common
- eOne of the causes of SIADH is pneumonia

The correct answers are: It can cause hyponatremia, In case of incorrect treatment it can lead to pontine demyelination, One of the causes of SIADH is pneumonia

Translated text

Send to Translate Home

Select all

Listen

Are these results useful?

YES

NO

11:41

14:17 A

Google Lens

Czech → English

The
has a hig
acidosis as a result of hypoventilation.

Task 6

0.00/1.00 point

The following characteristics are NOT used to classify edema:

Select one or more options:

- Pitting a non-pitting X
- Acquired and Inherited✓
- c. Hyperosmolar and hypoosmolar
- d. Local and generalized
- e Primary and secondary lymphedema

The correct answers are:
Hyperosmolar and hypoosmolar,
Acquired and inherited

Task 7

Points 0.00/1.00

Translated text

Send to Translate Home | Select all | Listen

Are these results useful? YES NO

Task 2

Which acid-base balance disorder develops primarily in patients with long-term simple starvation

Select one or more options:

- [ ] A. Combined metabolic and respiratory acidosis
- [x] B. Metabolic acidosis
- [ ] C. Respiratory acidosis
- [ ] D. Metabolic alkalosis
- [ ] E. Respiratory alkalosis

11:35 moodle.upol.cz/mod/q

Google Lens (57%)

Czech → English

Choose the correct statement/s about SIADH

Select one or more options:

- [x] May cause hyponatremia ✓
- [ ] b It is associated with a reduced ANP level
- [ ] eOne of the causes of SIADH is pneumonia
- [ ] d Swelling of the lower extremities is common
- [ ] e In case of incorrect treatment, it can lead to pontine demyelination

The correct answers are: It can cause hyponatremia, In the case of incorrect treatment, it can lead to pontine demyelination.

One of the causes of SIADH is pneumonia

Translated text

Send to Translate Home | Select all | Listen

11:33 20% 7:
moodle.upol.cz/mod/q
Google Lens
Czech → English

Role/

In the case of salicylate poisoning, it is used in the development of an acid-base disorder

Select one or more options:

- [ ] aprüjem - metabolic alkalosis
- [x] b metabolic acidosis ✓
- [ ] respiratory center depression -respiratory acidosis
- [ ] d. irritation of the respiratory center, respiratory alkalosis
- [ ] e formation of ketone bodies metabolic acidosis

The correct answers are:
irritation of the respiratory center, respiratory alkalosis
, metabolic acidosis

Translated text
Send to Translate Home
Select all
Listen

11:40

Google Lens

Czech → English

Task 7

Points 0.00/1.00

Select the processes that are common in the pathogenesis of gout:

Select one or more options:

- a Myocardial infarction is often the initial sign of gout
- b Hyperemia and severe pain in the affected joint during exacerbation ✓
- c High level of blood urea X
- d Uric acid crystals precipitate in the joint cavity
- e With the progression of gout, tophi appear on the hands, fingers, elbows and ears ✓

The correct answers are: Uric acid crystals precipitate in the joint cavity , Hyperemia and severe pain in the affected joint during exacerbation, As gout progresses, tophi appear on the hands, fingers , elbows and ears

Task 8

0.00/1.00 points

Hypochloremia is seen in

Select one or more options:

- a excessive losses of HCO3- from the organism (GIT, kidneys)
- b metabolic alkalosis during vomiting ✓
- c hypertonic dehydration ✓
- d. edematous conditions associated with dilutional hyponatremia
- e isotonic dehydration X

The correct answers are: edematous states associated with dilutional hyponatremia, metabolic alkalosis during vomiting

Translated text

Send to Translate Home | Select all | Listen

Are these results useful? YES NO

11:40

Google Lens

Czech → English

Task 5 Points 0.50/1.00

A 12-year-old boy developed decompensated type 1 diabetes mellitus due to an insufficient dose of insulin. His glucose level is 20.9 mmot/L blood pH is 7.1, the patient is severely dehydrated. Select all correct statements about the patient:

Select one or more options:

- a Most likely the patient has tachycardia due to dehydration and activation of the sympathetic nervous system
- b Most likely, the patient has a normal or reduced anion gap
- c Most likely, the patients blood osmolarity is high ✓
- d Most likely, the patient is dehydrated due to osmotic diuresis
- e Most likely, the patient has an increased level of ketone bodies in the plasma✓

The correct answers are: Most likely the patient has tachycardia due to dehydration and activation of the sympathetic nervous system, Most likely the patient has dehydration due to osmotic diuresis Most likely the patient's blood osmolarity is high, Most likely the patient has an elevated level of plasma ketone bodies

Task 6 Point 1.00/1.00

Anion gap is calculated from plasma concentration values

Select one or more options:

- a [Na+] +[K+] - [HCO3-]
- b Na] + [K+] - [Cl-]
- c |On [K]-[HC03-]
- d [Na[K+] + [Ca] + [HCO3-]-[Cl-1
- e [Na)-[Cl--HC031 ✓

The correct answer is: [Na+1-(Cl-1-HC031

Translated text

Send to Translate Home | Select all | Listen

Are these results useful? YES NO

11:40

Google Lens

Czech → English

Task 3 Points 1.00/1.00

A 52-year-old patient, a chronic alcoholic, is admitted to the hospital with signs of liver cirrhosis, acute alcohol intoxication and vomiting . His albumin level is 10 g/L (N 35-50 g/L), pH-7.38 mmol/L Cl = 90 , PaCO2-5.3 kPa (40 mm Hg). Choose the 1 most appropriate statement about the patient

Select one or more options:

- a The patient probably does not have acid-base balance disorders
- Ob Based on the available information no conclusion can be made about patient's acid-base balance
- c Most likely, the patient developed metabolic alkalosis due to vomiting
- ☑ d Most likely, the patient has a combination of metabolic acidosis and metabolic alkalosis ✓
- e Most likely, the patient has a combination metabolic acidosis and respiratory alkalosis

The correct answer is: Most likely, the patient has a combination of metabolic acidosis and metabolic alkalosis

Task 4 Point 1.00/1.00

What are the clinical manifestations of hypokalemia in the skeletal muscle , heart and GIT?

muscle weakness, convulsions to tetany, contraction disorders

heart - rhythm disorders, changes on the ECG, even cardiac arrest GIT - the result of hypokalemia can be diarrhea; reduced intestinal motility to paralytic ileus

Comment:

Translated text

Send to Translate Home | Select all | Listen

Are these results useful? YES NO

11:39

Google Lens

Czech → English

Task 1

Points 0.50/1.00

Choose the correct statement(s) about hyperchloremic acidosis

Select one or more options:

- O a. May be caused by vomiting
- b. The anion gap is usually normal
- ✓ It can be caused by income
- d.It is a type of respiratory acidosis

Oh It can be caused by a panic attack

The correct answers are: May be caused by diarrhea Anion gap is usually normal

Task 2

Points 1.00/1.00

Tissue lactic acidosis may NOT be caused

Select one or more options:

- ✓ hypoglycemia

With lack of oxygen in respiratory diseases

- e sepsis with MODS
- d binding of lactate metabolism in the liver

Oe. lack of oxygen in the tissues due to circulatory causes

The correct answer is: hypoglycemia.

Translated text

Send to Translate Home | Select all | Listen

Are these results useful? YES NO

11:33 26% 17:56

Google Lens

Czech → English

Task 10

Odds 0.50/1.00

Intermittent fasting can mean that a person:

Select one or more options:

- [ ] and Consumes only 1 type of food each week (e.g. apples 1 week, bread 2 weeks, cheese 3 weeks...)
- [ ] b Consumes a low-protein diet for 1 day, then a low-fat diet for 1 day, then a low-carb diet for 1 day
- [ ] c. Consumes only 1 type of food every day (e.g. 1 day apple, 2nd day bread , 3rd day cheese...)
- [ ] d. He eats 6 hours a day and does not eat for 18 hours
- [x] He eats for 8 hours a day and does not eat for 16 hours ✓

Translated text

Send to Translate Home | Select all | Listen

11:33

2-714.png Google Lens

Czech → English

13:12 67%

Task 3

Choose the correct statement(s) about simple starvation:

Select one or more options:

- [ ] A. prolonged starvation can be irreversible due to degeneration of the GIT mucosa and pancreas
- [x] B. during the first 10-12 hours, the body receives energy mainly through lipolysis
- [ ] C. patients who drink water but do not consume food usually die from kidney failure that develops due to high protein catabolism
- [ ] D. the main problem leading to clinical symptoms is vitamin C deficiency
- [x] E. After three to five days, lipids are the main source of energy

Translated text

Send to Translate Home | Select all | Listen

Are these results useful? YES NO

12:36 p.m 71%

Task 9

The lack of bicarbonates arises as a result

Select one or more options:

- [ ] a. hypernatremie
- [ ] b. often due to inadequate bicarbonate intake
- [ ] c. hyperaldosteronismu
- [x] d. renal tubular acidosis
- [x] e. diarrhea

Complete the tour

12:36 p.m 71%

Task 9

The lack of bicarbonates arises as a result

Select one or more options:

- [ ] a. hypernatremie
- [ ] b. often due to inadequate bicarbonate intake
- [ ] c. hyperaldosteronismu
- [x] d. renal tubular acidosis
- [x] e. diarrhea

Complete the tour

02 CZ02 SK 10147% 13:57
moodle.upol.cz/mod/qu

Task 5

They do NOT belong to alkalizing agents

Select one or more options:

- [ ] and sodium lactate
- [ ] NH 4 Cl
- [ ] bicarbonate
- [ ] sodium citrate

Task 6

Write down the question number according to the question list and the answer to that question. You do not need to rewrite the wording of the question.

O2 - CZ|O2 - SK 10147% 13:57

Task 4

Hiatal hernia

Select one or more options:

- [ ] and. It often leads to cirrhosis of the liver
- [x] b. It can manifest as heartburn
- [x] c. It can lead to the development of strangulation hernia or obturation
- [ ] d. It is often manifested by diarrhea
- [ ] E. It is one of the important causes of celiac disease

Task 5

It does NOT belong to alkalizing agents

O2 - CZ|O2 - SK 10147% 13:57

Task 7

Lactose intolerance in adults is caused by

Select one or more options:

- [ ] a. excessive splitting of lactose into glucose and galactose in the intestine
- [x] b. lactose hydrolysis disorder in the intestine
- [ ] c. a malfunction of the mechanisms ensuring the transport of lactose and galactose through the intestinal wall
- [ ] d. a disorder of lactose phosphorylation in the intestinal wall

Task 8

Select an important cause/important

O2 - CZ|O2 - SK 10147% 13:58

Task 8

Select the important cause(s) of hypernatremia:

Select one or more options:

- [x] Profuse sweating in patients with fever
- [ ] B. Sterile wound on the forearm without signs of SIRS
- [ ] CSIADH
- [ ] D. Type 2 diabetes mellitus, fully compensated state

Task 9

Select the processes that are common in the pathogenesis of gout:

O2 - CZ|O2 - SK 10147% 13:58

Task 8

Select the important cause(s) of hypernatremia:

Select one or more options:

- [x] A. Profuse sweating in patients with fever
- [ ] B. Sterile wound on the forearm without signs of SIRS
- [ ] C. SIADH
- [ ] D. Type 2 diabetes mellitus, fully compensated

state

Task 9

Select the processes that are common in the pathogenesis of gout:

b

O2 - CZ|O2 - SK 10147% 13:57

moodle.upol.cz/mod/qu 7

Task 5

They do NOT belong to alkalizing agents

Select one or more options:

- [ ] and sodium lactate
- [ ] NH4C
- [ ] bicarbonate
- [ ] sodium citrate

Task 6

Write down the question number according to the question list and the answer to that question. You do not need to rewrite the wording of the question.

4:04 p.m
54
moodle.upol.cz/mod/qui:O)
2

Task 2

Acidemia

Select one or more options:

- [ ] a. leads to hypokalemia
- [ ] b. leads to hypokalemia
- [x] leads to irritation of receptors of the respiratory center
- [x] shifts the cystic-hemoglobin dissociation curve to the right

Task 3

16:050 54
moodle.upol.cz/mod/qui: 2

hyperaldosteronismem

- [x] e. It is typical for patients with end-stage renal insufficiency

---

Task 9

Profuse vomiting leads to

Select one or more options:

- [ ] a. potassium retention
- [ ] b. metabolic acidosis
- [x] c. hypochloremia
- [ ] d. potassium depletion

(apartment

Complete the tour

TO

O2 - CZ|O2 - SK 10147% 13:57

Task 7

Lactose intolerance in adults is caused by

Select one or more options:

- [ ] and. by excessive splitting of lactose into glucose and galactose in the intestine
- [x] b. lactose hydrolysis disorder in the intestine
- [ ] c. disruption of the mechanisms ensuring the transport of lactose and galactose through the intestinal wall
- [ ] d. a disorder of lactose phosphorylation in the intestinal wall

Task 8

Select an important cause/important

16:04 83%

Task 1

Hirschsprung's disease
(megacolon congenitum)

Select one or more options:

- [x] and. is a congenital disease in which the ganglion cells in the intestinal plexus are missing
- [x] b. is manifested by diarrhea due to insufficient digestion of fats
- [ ] c. is manifested by organic constipation due to stagnation of *intestinal contents*
- [ ] d. is an intestinal manifestation of congenital progressive myodystrophy

16:04 D ⊖ O 54
moodle.upol.cz/mod/qui: 2

Task 2

Acidemia

Select one or more options:

- [ ] a. leads to hypokalemia
- [ ] b. leads to hypocalcemia
- [x] c. leads to irritation of receptors of the respiratory center
- [x] d. shifts the oxygen-hemoglobin dissociation curve to the right

Task 3

< TO

4:04 p.m 54

Task 4

Select the typical symptoms of vitamin deficiency:

Select one or more options:

- [ ] a. Vitamin C deficiency significantly increases the risk of thrombosis
- [x] Long-term use of wide-spectrum antibiotics can lead to vitamin K deficiency and increase the risk of bleeding
- [x] Lack of vitamin A leads to squint blindness, xerophthalmia, keratinization of the skin and mucous membranes
- [ ] Lack of vitamin D increases the risk of muscle weakness, muscle cramps
- [ ] e. Lack of vitamin D increases the risk of night blindness, xerophthalmil

4:05 p.m
moodle.upol.cz/mod/qui:O)

hyperaldosteronismem

e. It is typical for patients with end-stage renal insufficiency

Task 9

Profuse vomiting leads to

Select one or more options:

and potassium retention

b. metabolic acidosis

hypochloremia

d. Drastic depletion

(apartment

Complete the tour

4:04 p.m AND. 83%

Task 7

Administering insulin infused with glucose has this effect on potassium

Select one or more options:

- [ ] a. none
- [ ] b. potassium tends to rise because insulin reduces urinary potassium excretion
- [x] potassium will drop because insulin causes potassium to enter the cells
- [ ] d. potassium rises because insulin causes potassium to leak out of the cells

Task 8

4:04 p.m 83%

Task 7

Administering insulin infused with glucose has this effect on potassium

Select one or more options:

- [ ] and. none
- [ ] b. potassium tends to rise because insulin reduces urinary potassium excretion
- [x] c. potassium will drop because insulin causes potassium to enter the cells
- [ ] d. potassium rises because insulin causes potassium to leak out of the cells

Task 8

4:05 p.m 83%


Task 8


Constipation

Select one or more options:


- [ ] a. can occur with hyponatremia
- [ ] b. arises in hyperthyroidism
- [x] c. can occur during dehydration
- [x] d. is a negative factor that may contribute to portosystemic encephalopathy in advanced liver cirrhosis
- [ ] e. risk factors include older age


70%
Google Lens
X Collected Questions - Goog
צ'כית ← אנגלית
Collected questions


































תרגום
שליחה לדף הבית של Translate
בחירת הכול
האזנה
האם התוצאות האלה מועילות?
כן
לא

Task 7

Ulcerative changes in the oral cavity

Select one or more options:

- [ ] a. do not occur in infectious diseases
- [x] b. can be manifested in allergic diseases
- [x] c. occur in states of deficiency
- [x] d. may develop in patients with Crohn's disease

Task 4

Secondary hyperaldosteronism

Select one or more options:

a. reduces the ratio of Na/K concentrations in urine

b. leads to metabolic acidosis

c. increases the concentration ratio of Na/K in urine

d. leads to metabolic alkalosis

e. can lead to an increase in arterial blood pressure

Task 4

Secondary hyperaldosteronism

Select one or more options:

- [x] a. reduces the concentration ratio of Na/K in urine
- [ ] b. leads to metabolic acidosis
- [x] c. increases the Na/K concentration ratio in the urine
- [x] d. leads to metabolic alkalosis
- [x] E. can lead to an increase in arterial blood pressure

Task 2

Which acid-base balance disorder develops primarily in patients with long-term simple starvation

Select one or more options:

- [ ] A. Combined metabolic and respiratory acidosis
- [x] B. Metabolic acidosis
- [ ] C. Respiratory acidosis
- [ ] D. Metabolic alkalosis
- [ ] E. Respiratory alkalosis

Task 8

Crohn's disease

Select one or more options:

- [ ] and. It is an acute inflammatory disease of the colon
- [x] D. Extraintestinal complications include joint inflammations, canker sores
- [x] C. Granulomatous inflammation penetrates the entire wall
- [x] d. It most often affects the small intestine (ileitis)
- [ ] E. It usually starts with involvement of the anus

4:21 p.m

X

Test #2 2021: Suddenly...
moodle.upol.cz

Task 1

Select the processes that are common in the pathogenesis of gout:

Select one or more options:

- [ ] a. High level of urea in the blood
- [x] b. As the gout progresses, tophi appear on the hands, fingers, elbows and ears
- [x] c. Uric acid crystals precipitate in the joint cavity
- [ ] d. *Myocardial infarction is* often the initial sign of gout
- [x] e. Hyperemia and severe pain in the affected joint during exacerbation

Task 2

4:21 p.m.

Test #2 2021: Suddenly...
moodle.upol.cz

Task 1

Select the processes that are common in the pathogenesis of gout:

Select one or more options:

- [ ] c. High level of urea in the blood
- [x] As the gout progresses, tophi appear on the hands, fingers, elbows and ears
- [x] Uric acid crystals precipitate in the joint cavity
- [ ] d. Myocardial infarction is often the initial sign of gout
- [x] Hyperemia and severe pain in the affected joint during exacerbation

Task 2

Task 7

Ulcerative changes in the oral cavity

Select one or more options:

- [ ] and. do not occur in infectious diseases
- [x] b. can be manifested in allergic diseases
- [x] c. occur in states of deficiency
- [x] d. can develop in patients with Crohn's disease

4:22 p.m

Task 4

Causes of normochloremic metabolic acidosis do NOT include

Select one or more options:

- [ ] a. starvation
- [ ] b. alcohol poisoning
- [ ] c. salicylate poisoning
- [x] d. diabetes insipidus

Task 5

Plasma concentration of chlorides is increased at

Select one or more options:

- [ ] AND _____
- [ ] B. prolonged vomiting
- [x] with severe diarrhea
- [ ] D. respiratory alkalosis
- [x] E. renal tubular acidosis

malignant lymphoma

Task 3

Lactase deficiency

Select one or more options:

- [x] e. causes intolerance to milk and some dairy products
- [ ] b. causes Crohn's disease
- [x] *some patients have* diarrhea and pain after milk and milk products, some have minimal problems
- [ ] d. causes animal fat *intolerance*
- [ ] E. causes intolerance to foods containing gluten

Task 8

Crohn's disease

Select one or more options:

- [ ] and. It is an acute inflammatory disease of the colon
- [x] b. Extraintestinal complications include joint inflammations, canker sores
- [x] c. Granulomatous inflammation penetrates the entire wall
- [x] d. It most often affects the small intestine (ileitis)
- [ ] e. It usually starts with involvement of the anus

malignant lymphoma


Task 3

Lactase deficiency

Select one or more options:

- [ ] a. causes intolerance to milk and some dairy products
- [ ] b. causes Crohn's disease
- [ ] c. some patients have diarrhea and pain after milk and milk products, some have minimal problems
- [ ] d. causes animal fat intolerance
- [ ] E. causes intolerance to foods containing gluten

71% 11:30 יום ד׳, 1 בנוב׳
Google Lens
X Collected Questions - Goog
צ׳כית ← אנגלית
Sesb jpg.2-504



02.CZ 2:38 p.m 36%
moodle.upol.cz

Task 7


5% dextrose intravenously distributes in a few hours mainly to:

Select one or more options:

- [ ] a. transcellular fluids
- [x] b. intravascular compartment
- [x] c. extracellular compartment
- [x] d. intracellular compartment
- [ ] e. interstitial compartment

תרגום
שליחה לדף הבית של Translate
בחירת הכול
האזנה
האם התוצאות האלה מועילות?
כן
לא

71% 11:30 יום ד׳, 1 בנוב׳
Google Lens
X Collected Questions - Goog
צ׳כית ← אנגלית
Sesb jpg.2-502



02.CZ 2:38 p.m 36%
moodle.upol.cz

Task 8


Early postprandial syndrome (dumping syndrome) develops most often as a result

Select one or more options:

- [x] a. partial gastrectomy with loss of pyloric function
- [ ] b. vagotomy
- [ ] c. total resection of the duodenum
- [ ] d. total resection of the stomach

תרגום
שליחה לדף הבית של Translate
בחירת הכול
האזנה
האם התוצאות האלה מועילות?
כן
לא

71% 11:30 יום ד׳, 1 בנוב׳
Google Lens
X Collected Questions - Goog
צ׳כית ← אנגלית
Sesb jpg.2-500



02.CZ 2:39 p.m 36%
moodle.upol.cz

Task 9


In the event of a metabolic disorder of the acid-base balance, events leading to its adjustment take place in this program


Select one or more options:

- [ ] a. compensation, buffer reaction, correction
- [ ] b. buffer reaction, correction, compensation
- [x] c. buffer reaction, compensation, correction
- [ ] d. correction, compensation, buffer reaction

תרגום
שליחה לדף הבית של Translate
בחירת הכול
האזנה
האם התוצאות האלה מועילות?
כן
לא

71% 11:30 יום ד', 1 בנוב'
Google Lens
X Collected Questions - Goog
צ'כית ← אנגלית
Sesb jpg.2-498





תרגום
שליחה לדף הבית של Translate
בחירת הכול
האזנה
האם התוצאות האלה מועילות?
כן
לא

77% 5:44 p.m

Task 10

Points 1.00 / 1.00

The mother of an 8-month-old boy comes to the doctor complaining that her son does not seem well after she stopped breastfeeding him approximately 2 weeks ago and began gradually giving him a regular diet, including some baked goods. He had intermittent diarrhea, decreased appetite, stomach pain and bloating for several days. He has lost approximately 0.5 kg of weight in the last 2 weeks. He had no vomiting and no fever during the past month. Before that, he seemed to be a generally healthy child. Most likely, the patient has:

Select one or more options:

- [ ] a. Lactose intolerance
- [ ] b. Phenylketonuria
- [ ] C. Cystic fibrosis
- [x] d. Celiac disease ✓
- [ ] e. Fructose intolerance

14:17 Θ Δ

LTE+ 91

Task 5

Points 1.00 / 1.00

A 65-year-old patient was hospitalized with symptoms of acute respiratory disease. He has a history of smoking for 25 years and obstructive pulmonary disease for 10 years. Within hours, PCR analysis showed that he was positive for COVID-19 and, according to a lung X-ray, he had symptoms of pneumonia. His respiratory rate is 23/min, heart rate 105 beats/min, arterial blood pressure 130/80 mm Hg, t = 38.3 °C, leukocyte count 3.9*10^9/L.

Choose the correct statement about the patient considering the greatest possible complications in the next 2 days:

Select one or more options:

- [x] a. The patient is at high risk of developing respiratory acidosis due to hypoventilation✓
- [ ] b. The patient is at high risk of developing dehydration due to kidney damage
- [ ] c. The patient is at high risk of developing metabolic alkalosis due to diarrhea
- [ ] d. The patient is at high risk of developing metabolic acidosis due to vomiting

4:05 p.m

Trial length	15 min. 58 seconds
Stamp	5.25 from possible 10.00 (53%)

Task 1 0.00/1.00 points

Choose the correct statement(s) about SIADH

Select one or more options:

- Swelling of the lower extremities is common X
- D. May cause hyponatremia ✓
- In CV case of incorrect treatment, pontine demyelination can occur
- d. One of the causes of SIADH is pneumonia
- e It is associated with a reduced level of ANP

The correct answers are: It can cause hyponatremia, In the case of incorrect treatment, pontine demyelination can occur, One of the causes of SIADH is pneumonia

The correct answer is: can lead to acute liver failure.

Task 14

Correctly

Points 1.00/1.00

Which of the following systems is not a buffering system of the human body?

Select one or more options:

- a. protein system
- b. sodium chloride system✓
- c. hemoglobin system
- d. bicarbonate system
- e. phosphate system

The correct answer is: sodium chloride system.

Hyperchloremic acidosis

224

Select one or more options:

- a. is characterized by an increased anion gap
- b. it is characterized by increased chloride levels and decreased bicarbonate levels
- c. may be due to impaired bicarbonate absorption in the proximal tubule
- d. cannot be caused by a diarrheal disease

It applies to Crohn's disease

Select one or more options:

- [ ] a. the sigmoid and rectum are most often affected
- [ ] b. the terminal ileum and the adjacent cecum are most often affected
- [ ] c. H. pylori is the most important etiological factor
- [ ] d. the inflammation spreads continuously, the mucosa is affected diffusely

77% 17:42
moodle.upol.cz/mod/qu 5

Problem 3 Points 0.50/1.00

Select conditions associated with hypercholesteroiemia

Select one or more options:

- [ ] a. Nephrotic syndrome
- [ ] b. Hepatocellular carcinoma with cachexia
- [x] c. Type diabetes mellitus II ✓
- [ ] d. Anorexia nervosa
- [x] e. Hypothyroidism ✓✓

The correct answers are:
Diabetes mellitus type II, Anorexia nervosa, Nephrotic syndrome, Hypothyroidism

Task 4 Points 1.00/1.00

The correct answer is: it monitors the emergence and development of a disease state in organs and tissues.

Problem

10 Incorrect

They will score 0.00/1.00

Select the incorrect statement about blood sodium concentration.

Select one or more options:

- [ ] a. can be lowered in patients with primary hyperaldosteronism
- [ ] b. is regulated by the renin-angiotensin-aldosterone system
- [ ] c. may be reduced in patients with syndrome of inappropriate secretion of antidiuretic hormone
- [ ] d. hyponatremia can lead to cerebral edema
- [x] e. is regulated by atrial natriuretic peptide X

The correct answer is: it can be lowered in patients with primary hyperaldosteronism.

Task 11

Correctly

They will score 1.00/1.00

Rising carbon dioxide concentrations and decreased blood pH due to chronic obstructive pulmonary disease indicate:

Task 10 100/100

[redacted] year old boy comes to the doctor and complains that her son does not seem to be well after he ingested [redacted] ago. [redacted] reports a normal diet, including some peanuts. For several days he had intermittent diarrhea, decreased appetite [redacted]. During the last 2 weeks he has lost approximately 0.5 kg of weight. He has not vomited or had a fever. [redacted] healthy child. Most likely, the patient has

Select one or more options:

- a. [redacted]
- b. Don't rush the fruits
- c. CeliAH ✓
- d. Lactor intolerance
- e. Cystic fluorosis

[redacted]

URS of the Packho-USU University x

Task 6050/1.00

I am characterized by Ditis kwashiorkor

Select one or more options:

- A protruding flash ✓
- Low resistance to infections. ✓
- Enlarged liver with fatty infiltrates
- Cd Pitting edema with swelling of ankles and feet
- Lack of dietary carbohydrates

The correct answers are: Wynivajicin bichem, Low resistance in infectient jerys by such indiring Piting ey
ankle and feet

Task 701.00/1.00

Shypochlorémil we will meet at

Select one or more options

- isotonic dernydration
- metabolic akabras have begun ✓
- Hypertonic dehydration
- edematous conditions associated with dubs hyponatremia ✓

ee funny expression

What are the clinical manifestations of hypokalemia in skeletal muscle, heart and G11?

on the heart - leads to bradycardia and arrhythmias, on the EKG I observe - a flat T wave, a U wave appears, widened QRS GIT is reduced peristalsis, can lead to paralytic ileus

muscle - muscle weakness, weakness,

Problem 6 Points 1.00 / 1.00

Select conditions associated with hypercholesterolemia

Select one or more options:

- Anorexia nervosa ✓
- Nephrotic syndrome ✓
- Hypothyroidism ✓
- Hepatocellular carcinoma with cachexil
- Type II diabetes mellitus ☐ ✓

Same afternoon

Mertal anorexia. Notrotic syndrome, Hypothyroidism

2000-25 PENCALL

The patient is more likely to have tachycardia due to dehydration and activation of the sympathetic nervous system

The correct answers are: Most likely the patient has tachycardia due to dehydration and activation of the sympathetic nervous system, Most likely the patient has dehydration due to osmotic diuresis. Most likely the patient's blood osmolarity is high, most likely the patient has an elevated level of plasma ketone bodies

Problem 3 Points 1.00/1.00

A 52-year-old patient, a chronic alcoholic, is admitted to the hospital with signs of liver cirrhosis, acute alcohol intoxication, and vomiting. His albumin level is 10 g/L (N = 35-50 g/L), pH = 7.38, CL = 90 mmol/L, PaCO2 = 5.3 kPa (40 mm Hg). Choose the 1 most appropriate statement about the patient:

Select one or more options:

a. The patient most likely has a combination of metabolic acidosis and respiratory alkalosis

Most likely, the patient has a combination of metabolic acidosis and metabolic alkalosis✓

C. Based on the available information, no conclusion can be made about the patient's acid-base balance

d. The patient most likely developed metabolic alkalosis due to vomiting

Be The patient probably does not have acid-base balance disorders

The correct answer is: Most likely, the patient has a combination of metabolic acidosis and metabolic alkalosis.

Task 7

Select the typical symptoms of vitamin deficiency:

Select one or more options:

- [ ] a. Long-term use of broad-spectrum antibiotics can lead to vitamin K deficiency and an increased risk of bleeding
- [ ] b. Vitamin D deficiency increases the risk of mood swings and depression
- [ ] c. Lack of vitamin A leads to night blindness, xerophthalmia, keratinization of the skin and mucous membranes ✔
- [ ] d. Lack of vitamin D increases the risk of muscle weakness, muscle spasms
- [ ] e. Vitamin C deficiency significantly increases the risk of thrombosis

The correct answers are: Lack of vitamin A leads to night blindness, xerophthalmia, keratinization of the skin and mucous membranes, Lack of vitamin D increases the risk of muscle weakness, muscle spasms, Lack of vitamin D increases the risk of mood swings and depression, Long-term use of broad-spectrum antibiotics can lead to a lack of vitamin K. and to increasing the risk of bleeding

---

Task 8

Describe the pathogenesis of ketoacidosis

Pathogenesis of ketoacidosis: Glucose reserves have been depleted (starvation or inability to use glucose - lack of insulin). as a source of energy, fats are split (lipolysis). MK are formed, which are metabolized to acetyl Coa by Beta oxidation - acetyl Coa enters the CC, however, due to the large amount of acetyl Coa, ketoacids (acetate, acetoacetate, B - hydroxy butyrát) are formed. They can be a source of energy or they are excreted from the body

12:36 p.m 71%

Task 7

It applies to potassium

Select one or more options:

- [ ] and decreases with tissue breakdown, burns, hemolysis
- [ ] b. in metabolic alkalosis, potassium is released from binding to intracellular phosphates and proteins
- [x] c. it is dependent on a sufficient energy supply for the proper function of the Na+/K+ ATPase
- [ ] TO d. it is influenced by the current state of renal function

12:34 p.m 71%

Task 2

Malabsorption syndrome

Select one or more options:

- [ ] a. it is usually accompanied by a tendency to develop metabolic alkalosis as a result of diarrhea
- [x] b. it is usually accompanied by diarrhea
- [ ] c. It is most often related to stomach disorders
- [x] d. it can manifest as hypochromic or megaloblastic anemia
- [ ] E. it is usually accompanied by obstinate constipation

Task 3

During blood flow through systemic capillaries

Select one or more options:

- [ ] the dissociation curve of hemoglobin oxygen saturation shifts to the left
- [ ] b. chlorides pass from erythrocytes to plasma
- [x] c. chlorides pass from plasma to erythrocytes
- [x] d. pH decreases
- [ ] e. bicarbonates pass from plasma to erythrocytes

Problem 2 Points 1.00/1.00

A 65-year-old patient was hospitalized with symptoms of acute respiratory disease. He has a history of smoking for 25 years and obstructive pulmonary disease for 10 years. Within hours, PCR analysis showed that he was positive for COVID-19 and, according to a lung X-ray, he had symptoms of pneumonia. His respiratory rate is 23/min, heart rate 105 beats/min, arterial blood pressure 130/80 mm Hg, t- 38.3 °C, leukocyte count 3.9*10^9/L. Choose the correct statement about the patient considering the greatest possible complications in the next 2 days:

Select one or more options:

- [ ] a. The patient is at high risk of developing respiratory alkalosis due to hypoventilation
- [ ] b. The patient is at high risk of developing dehydration due to kidney damage
- [ ] c. The patient is at high risk of developing metabolic alkalosis due to diarrhea
- [ ] The patient is at high risk of developing metabolic acidosis due to vomiting
- [x] The patient has a high risk of developing respiratory acidosis due to hypoventilation ✓

The correct answer is: The patient is at high risk of developing respiratory acidosis due to hypoventilation.

Duodenal ulcer is a multifactorial disease. However, it turned out that 592 plays a significant role in its creation

Select one or more options:

- [x] a. increased output of gastrin
- [ ] b. reduction of vagus tone
- [x] c. smoking
- [ ] d. dysplasia of duodenal epithelium caused by hypersecretion of digestive enzymes
- [x] chronic Helicobacter pylori infection

Task 2

Done

Score out of 1.00

Decreased plasma protein levels

Select one or more options:

a. activates the renin-angiotensin-aldosterone system

b. will decrease potassium excretion

c. will increase the transfer of water from the blood to the interstitium

d. Helps treat ascites

e. will increase sodium excretion

I don't know

D 26% 1 5:36 p.m

Problem 9 Points 0.50/1.00

Describe the pathogenesis

of ketoacidosis

ketoacidosis - the presence of a large amount of ketone bodies in the blood

- this increased amount is, for example, during starvation, diabetes mellitus

1

- ketone bodies are formed in the liver from acetyl-CoA

Comment:

Task 10 0.50/1.00 points

Intermittent

fasting can mean

that a person:

O2 - SK 4:56 p.m @ 51%
moodle.upol.cz - Anonymous window

thrombosis

C. Vitamin A deficiency leads to night blindness, xerophthalmia, keratinization of the skin and mucous membranes✓✓✓

d. Vitamin D deficiency increases the risk of mood swings and depression

e. Long-term use of broad-spectrum antibiotics can lead to vitamin K deficiency and increase the risk of bleeding✓✓

The correct answers are:

Vitamin A deficiency leads to night blindness, xerophthalmia, keratinization of the skin and mucous membranes, Vitamin D deficiency increases the risk of muscle weakness, muscle spasms, Vitamin D deficiency increases the risk of mood swings and depression, Long-term use of broad-spectrum antibiotics can lead to vitamin Ka deficiency increasing the risk of bleeding

moodle.upol.cz

Facebook

Two correct answers are: Uric acid crystals are produced in the joint cavity. Hyperemia and severe pain in the affected joint during

Task 9 Score 1.00

Give 4 examples of swelling caused by increased hydrostatic pressure:

blood stasis in heart failure

vein thrombosis by tumor

100/100 task

Select one or more options:

- Zvětšenými játry s tukovými infiltráty
- with a growing stomach ✓
- Low susceptibility to infections ✓
- Lack of dietary carbohydrates

Task 3

During blood flow through systemic capillaries

Select one or more options:

- [ ] a. the dissociation curve of hemoglobin oxygen saturation shifts to the left
- [ ] b. chlorides pass from erythrocytes to plasma
- [x] c. chlorides pass from plasma to erythrocytes
- [x] d. pH decreases
- [ ] e. bicarbonates pass from plasma to erythrocytes

77% 5:43 p.m

Task 9

Points 1.00 / 1.00

Choose the correct statement about simple starvation:

Select one or more options:

- [x] a. Adult patients can usually survive for more than 4 weeks if they consume only water ✓
- [ ] b. Approximately 4-5 kilograms of body muscle will be lost during the first 7 days of fasting
- [x] c. After a few days, the energy metabolism decreases ✓
- [ ] d. On day 14, the patient will likely have metabolic alkalosis due to *proteolysis*
- [ ] e. During the first 10-12 hours, the body receives energy mainly thanks to lipolysis

14:17 LTE+ 91
moodle.upol.cz/mod/quiz/re 5 :

--> ΓΙΖΙΚΟ ILHS

Comment:

Task 4

0.50 / 1.00 points

Major vomiting will lead to

Select one or more options:

- [ ] a. metabolic acidosis due to large losses of chlorides
- [x] b. metabolic alkalosis due to a large loss of hydrogen ions in the acidic gastric juice✓
- [ ] c. Hypochloremia
- [ ] d. metabolic acidosis due to compensatory formation of HCL
- [ ] e. metabolic alkalosis due to compensatory hyperkalemia

The correct answers are:
metabolic alkalosis due to a large loss of hydrogen ions in the acidic gastric juice,
Hypochloremia

Task 1

What changes are typical of severe dehydration caused by sweating (uncompensated by drinking)?

Select one or more options:

- [ ] a. Tachycardia due to activation of the parasympathetic nervous system
- [ ] b. Decreased plasma Na + concentration due to loss of Na + through sweat
- [ ] c. No answer is correct
- [ ] d. Increased amount of Na + in the body due to increased secretion of ANP (atrial natriuretic peptide)
- [x] e. Increased arterial blood pressure due to activation of the sympathetic nervous system

The correct answer is: None of the answers are correct.

The intrinsic factor complex with cobalamin (vitamin B12) is absorbed

Select one or more options:

- [ ] a. by paracellular transport in the jejunum and ileum
- [ ] b. facilitated diffusion in the ileum
- [x] c. using receptors in the area of the brush border of the ileum
- [ ] d. in the cells of the gastric pylorus

Task 1

0.00 / 1.00 points

Chronic lead poisoning resulted in damage to the renal proximal tubules in a young patient. Select the correct statement(s) regarding the patient, assuming he has no other damage to the body due to lead poisoning:

Select one or more options:

- Most likely, the level of HCO3- in the urine will be high✓
- Most likely, the patient has proximal tubular alkalosis
- Most likely, the patient has proximal tubular acidosis
- Most likely, the patient has hyperkalemia and hypernatremia X
- The patient is most likely hypokalemic

The correct answers are: The patient most likely has proximal tubular acidosis. The patient most likely has hypokalemia. Most likely, the level of HCO3+ in the urine will be high

Important risk factors for a hiatal hernia include:

Select one or more options:

that's how it is in times gone by

- [x] a. Cough
- [x] b. Pregnancy
- [ ] c. Lifting heavy objects
- [x] d. Obesity
- [ ] e. Age over 70 years

moodle.upol.cz

With the osmolarity of the patient's blood vysuk. Shepti measures the patient will log the Sholin of hate cells from ... of the ...
... 2020, online 15:30 ... Facebook ... Úvodní strana
... 2020 ... pathophysiology of the cns: pharynx and atlas Current test: 2 (content of practical exercises 4-6 and further 4-6) Test 2-5/11 2020 ... online 15:30

Problem 2 B 1.00/1.00

A 52 year-old patient, a chronic alcoholic, is admitted to the hospital with signs of liver cirrhosis, acute alcohol intoxication and vomiting. His albumin level is 18 g/L (N 35-50 g/L), pH 7.38, Cl- 90 mmol/L, paCO2 5.3 kPa (40 mm Hg). Choose the 1 most appropriate statement about the patient.

Select one or more options:

a. Most likely, the patient has a combination of metabolic acidosis and respiratory alkalosis

b. Most likely, the patient developed metabolic alkalosis due to vomiting

c. Most likely, the patient has a combination of metabolic acidosis and metabolic alkalosis ✓

d. Based on the available information, no conclusion can be made about the patient's acid-base balance

e. The patient probably does not have acid-base balance disorders

At first, Shepti responds with probabilities that the patient is a combination of metabolic acidity and metabolic al...

Task 3035/1.00

In chronic lead poisoning data in a young patient to renal proximal tubule damage. Select the correct/applicable one... the patient, provided that he has no other damage to the organism as a result of lead poisoning.

Select one or more options:

5. most likely the patient has hypokalemia

77% 17:42

Anorexia nervosa, Nephrotic

syndrome, Hypothyroidism

Problem 4 Points 1.00/1.00

Anion gap is calculated

from plasma concentration values

Select one or more options:

- [ ] a. [Na+] + [K+] - [HCO3-]
- [ ] b. [Na+] + [K+] - [HCO3-]
- [x] c. [Na+] - [Cl] - [HCO3-]
- [ ] d. [Na+] + [K+] + [Ca] + [HCO3-] - [CL-]
- [ ] e. [Na+] + [K+] - [CL-]

The correct answer is: [Na+] - [CL-] - [HCO3-].

4:06 p.m

Task 9

0.00/1.00 points

A 65-year-old patient was hospitalized with symptoms of acute respiratory disease. He has a history of smoking for 25 years and obstructive pulmonary disease for 10 years. Within hours, PCR-assisted analysis showed that he was positive for COVID-19 and, according to a lung X-ray, he had symptoms of pneumonia. His respiratory rate is 23/min, heart rate 105 beats/min, arterial blood pressure 130/80 mm Hg, t = 38.3 °C, leukocyte count 3.9*10^9/L. Choose the correct statement about the patient considering the greatest possible complications in the next 2 days:

Select one or more options:

- and The patient is at high risk of developing respiratory acidosis due to hypoventilation
- b. The patient is at high risk of developing metabolic alkalosis due to diarrhea
- c. The patient is at high risk of developing metabolic acidosis due to vomiting
- d. The patient is at high risk of developing respiratory alkalosis due to hypoventilation

2020-25-PFV/KCAH 77, Pathophysiology of the esophagus, stomach and intestines: Continuous test No. 2 (content of practical exercises 4-6 and lectures & 4-6). The correct answers are: It can cause hyponatremia. In case of incorrect technique, it can reach the pontine demyelinosis, One of the causes of SIADH is pneumonia

Task 6 Points 0.50/1.00

During a visit to the pediatrician, a fourteen-year-old wild girl is diagnosed with anorexia nervosa. Select the symptoms that helped the doctor make this diagnosis:

Select one or more options:

- Ascites due to hypoproteinemia
- Body weight below 5th percentile
- Body height above 95th percentile
- BMI 21.5 kg/m2
- Absence of periods for more than 3 months after having had regular periods since age 12 to 13 ✓

The correct answers are Body weight below 5 percent, Absence of menstruation for more than 3 months after, had regular menstruation from 12 to 13 years

What is typical of celiac disease?

Select one or more options:

- [x] and. gluten-induced autoimmune response
- [ ] b. lactose-induced autoimmune response
- [ ] c. is often associated with other autoimmune diseases
- [x] d. following a lifelong strict gluten-free diet is the best treatment
- [ ] e. reduced number of lymphocytes in the intestinal mucosa

i don't know

4:05 p.m

Comment:

Task 3

Points 1.00/1.00

Choose the correct statement about stress starvation

Select one or more options:

- a. Insulin levels are usually decreased
- b. It is characterized by intense protein catabolism✓
- c. Severe emotional stress (e.g. death of a partner) is one of the most important causes
- d. Glucose level often increases✓✓
- e. Ketone bodies are often elevated

The correct answers are: It is characterized by intense protein catabolism. The glucose level often increases

Lactic acidosis

Select one or more options:

a. it arises during tissue hypoxia

b. it arises from the accumulation of ketone bodies

c. it is formed by the accumulation of oxoacids

d. it arises as a result of the need to renew NAD+ when there is insufficient 02 supply to the mitochondria

e. is formed by the conversion of lactose into lactate

Task 3 Points 0.00/1.00

Select the processes that are common in the pathogenesis of gout:

Select one or more options:

- [x] a. High blood urea X
- [x] ✓b. As gout progresses, tophi appear on the hands, fingers, elbows and ears ✓
- [ ] c. Myocardial infarction is often the initial sign of gout
- [ ] d. Hyperemia and severe pain in the affected joint during exacerbation
- [x] e. Uric acid crystals precipitate in the joint cavity✓

The correct answers are: Uric acid crystals precipitate in the joint cavity, Hyperemia and severe pain in the affected joint during exacerbation, As gout progresses, tophi appear on the hands, fingers, elbows and ears

Task 4 Points 0.50/1.00

Hypochloremia is seen in

Select one or more options:

- [ ] a. excessive losses of HCO3- from the organism (GIT, kidneys)
- [ ] b. edematous conditions associated with dilutional hyponatremia
- [ ] c. hypertonic dehydration
- [x] d. metabolic alkalosis during vomiting✓
- [ ] e. isotonic dehydration

The correct answers are: edematous conditions associated with dilutional hyponatremia, metabolic alkalosis with vomiting

Task 8

Transcellular fluid is formed, among other things:

Select one or more options:

- [ ] a. lymph
- [x] b. glandular secretions
- [ ] c. blood plasma
- [ ] d. fluid in the cartilage a bones

d. is influenced by the current state of renal function

Task 8

It poses a fatal risk in patients with mechanical ileus

Select one or more options:

- a. weight loss
- D. resorption of bacterial toxins
- C. disorder of bile secretion
- d. hyperhydratace
- e. dehydration

14:17N

moodle.upol.cz/mod/quiz/re

Task 1

0.67/1.00 points

Choose the correct statement(s) about hyperkalemia

Select one or more options:

- [x] a. May cause arrhythmia✓
- [ ] b. It is a side effect of insulin therapy
- [ ] c. Associated with hyperaldosteronism
- [ ] d. May cause paresthesia and muscle weakness
- [x] e. It is typical for patients with end-stage renal insufficiency✓

The correct answers are: It is typical for patients with end-stage renal insufficiency, *It can cause paresthesia* and muscle weakness, It can cause arrhythmia

Task 2

Bodů 0,67 / 1,00

Choose the correct statement(s) about SIADH

26% 5:36 p.m.

Problem 9 Points 0.50/1.00

Describe the pathogenesis

of ketoacidosis

ketoacidosis - the presence of a

large amount of ketone bodies in the

blood

- this increased

amount is, for example, during

starvation, diabetes mellitus

1

- ketone bodies are

formed in the liver from acetyl-CoA

Comment:

Task 10 0.50/1.00 points

Intermittent

fasting can mean

that a person:

AND

* [© 77% 17:42

Task 5

0.00 / 1.00 points

A 52-year-old patient, a chronic alcoholic, is admitted to the hospital with signs of liver cirrhosis, acute alcohol intoxication, and vomiting. His albumin level is 10 g/L (N = 35-50 g/L), pH 7.38, CL = 90 = mmol/ PaCO2 = 5.3 kPa (40 mm Hg). Choose the 1 most appropriate statement about the patient:

Select one or more options:

✓a. Most likely, the patient has a combination of metabolic acidosis and metabolic alkalosis✓

b. Based on the available information, no conclusion can be made about the patient's acid-base balance

c. Most likely developed in the patient

12:35 p.m 71%
Test No. 2 2021: N...
moodle.upol.cz

Task 5

Complications of peptic ulcer disease include:

Select one or more options:

- [x] A. Anemia caused by occult (hidden) bleeding
- [ ] B. Dehydration due to frequent vomiting
- [ ] C. Chronic pancreatitis
- [x] D. Stomach cancer
- [x] E. Perforation

Task 1 Points 0.00/1.00

What changes are typical of severe dehydration caused by sweating (uncompensated by drinking)?

Select one or more options:

- a. Tachycardia due to activation of the parasympathetic nervous system
- b. Decreased plasma Na + concentration due to Na + loss through sweat
- c. No answer is correct
- d. Increased amount of Na + in the body due to increased secretion of ANP (atrial natriuretic peptide)
- e. Increased arterial blood pressure due to activation of the sympathetic nervous system X

The correct answer is: None of the answers are correct

Celiac sprue (celiac disease)

Select one or more options:

gliadin intolerance is induced

The prevalence in the Czech Republic is approximately 1:250-300 in the entire age spectrum

C. The basis of treatment is a gluten-free diet only during periods of difficulty

d. is caused by lactose intolerance

immune factors are important in the etiopathogenesis of the disorder

it's immune intolerance, so I don't know

O2 - SK 4:56 p.m @ 51%
moodle.upol.cz - Anonymous window

Task 3

0.00 / 1.00 points

Intermittent fasting can mean that a person:

Select one or more options:

- [x] a. Consumes only 1 type of food every day (e.g. apples 1 day, bread the next day, cheese 3 days...) X
- [ ] b. Consumes food for 8 hours a day and does not consume food for 16 hours
- [x] C. Consumes only 1 type of food each week (eg apples 1 week, bread 2 weeks, cheese 3 weeks...) X
- [ ] d. Eats a low-protein diet for 1 day, then a low-fat diet for 1 day, then a low-carb diet for 1 day
- [ ] e. Consumes food for 6 hours a day and does not consume food for 18 hours

And after school audios and ser Continuous test 2 arah of practical exercises 4-6 and pre-test 242

Task 5 Point 067/1.00

Choose the correct statement(s) about SIADH

Select one or more options:

- a. Swelling of the lower extremities is common
- b. One of the causes of SIADH is pneumonia
- c. In case of incorrect treatment, pontine demyelination can occur✓✓
- d. May cause hyponatremia✓
- e. It is associated with a reduced level of ANP

Spne answers are Can hyponatremil work, in the wrong case it can lead to ponting demyesnation. One of the oficio SADH is pneumonte

Task 60/140

Task 3

Points 1.00 / 1.00

Explain the pathogenesis of 3 symptoms of acute cholecystitis

pruritus- itching caused by increased bile acids pain in the right subcostal area- caused by bile congestion

acholic stools, steatorrhea - caused by a lack of bile in the intestine

Comment:

almost

Spre dpowd are the most likely that the patient has a tachycardia during dehydration and activation of the symje of the probability that the patient is dehydrated due to osmotic diuresis 5 the most likely to be related to the patent is high

Task 3100/1.00

A 2-year-old patient, a chronic alcoholic, is admitted to the hospital with signs of liver cirrhosis, acute alcohol intoxication, and [illegible] level = 1 g/L, rN-35-50 g/L, pH-7.38, Cl-90 mmol/L, PaCO2-5.3 kPa (40 mm Hg). Choose the 1 most appropriate statement about the patient.

Select one or more options:

- Most likely, the patient has a combination of metabolic acidosis and respiratory alkalosis
- The patient most likely has a combination of metabolic acidosis and metabolic alkalosis✓
- Based on the available information, no conclusion can be made about the patient's acid-base balance
- Most likely, the patient developed metabolic alkalosis *due to* vomiting
- The patient probably does not have acid-base balance disorders

It is likely that the patent has a combination of metabolic acidosis and metabolic alkalosis

Problem 6 Point 0.67/1.00

Select the important cause(s) of hypernatremia:

Select one or more options:

D&SIADH

- [x] a. Increased production of aldosterone ✓
- [ ] c. Glucocorticoid deficiency

From Diabetes insipidus

- [x] b. Increased sweating in patients with fever✓✓

The correct answers are: Increased production of aldosterone, Diabetes insipidus, Increased sweating in patients with fever

Renal tubular acidosis of the proximal type

Select one or more options:

- [ ] A. due to chloride reabsorption disorders, hypochloremia occurs
- [x] B. it leads to an increase, often to an alkaline pH of the urine
- [ ] C. does not result in bicarbonate losses through the kidneys
- [x] D. has a normal anion gap