

ORGANICS: THE BIO-SYSTEMS ARCHITECTURE MANUAL

**Living Substrates, Secondary Metabolite Chemistry, Regenerative Biomass
& Ecological Remediation**

Author: **Marie-Soleil Seshat Landry** • Principal Systems Architect

Organizations: **Landry Industries** (LandryIndustries.ca) • **Marie Landry's Spy Shop**
(marielandryspyshop.com)

Scope: Applied Pedology, Rhizosphere Microbiology, Organic Synthesis, Fermentation Bio-
Inputs, Mycoremediation


FIELD DIRECTIVE: "Organic" is not merely an agricultural certification or consumer marketing label. In industrial and biological engineering, *organics* represents the thermodynamics of living carbon architectures: closed-loop trophic cascades, rhizosphere electrochemistry, complex enzymatic polymerization, and biological defense mechanisms. This manual establishes standard operating procedures for engineering living soils, synthesizing bio-inputs, extracting bioactive secondary metabolites, and remediating contaminated biomes.

DOCUMENT INDEX

1. Foundations of Organic Systems: Carbon Dynamics & Soil Electrochemistry
2. Rhizosphere Microbiology: The Soil Food Web Engine
3. Biochemical Soil Building: Mineral Fractions, Humic Substances & CEC
4. Secondary Plant Metabolites: Terpenoids, Phenolics, Alkaloids & Defense Chemistry
5. Regenerative Fermentation Protocols: KNF, JADAM & Liquid Inoculants
6. Composting Dynamics: Aerobic Thermal In-Vessel vs. Static Fungal Humification
7. Bioremediation: Mycoremediation, Heavy Metal Hyperaccumulation & Soil Cleansing
8. Biomaterials & Closed-Loop Industrial Organics (Mycelium, Chitin, Bio-Polymers)
9. The Master Field Recipe & Formulation Repository
10. Quality Assurance, Chromatography & Biological Verification Checklist

1. FOUNDATIONS OF ORGANIC SYSTEMS: CARBON DYNAMICS & ELECTROCHEMISTRY

Organic systems operate on the energetic storage and release of solar photons mediated by the carbon atom. The unique tetravalent bonding capacity of carbon allows the construction of complex biochemical polymers—cellulose, lignin, proteins, lipids, and humic macromolecules—that serve as the thermodynamic currency of terrestrial ecosystems.



In living substrates, soil is not a passive mechanical anchor; it is an active electrochemical matrix. Cation-Exchange Capacity (CEC), oxidation-reduction potential (Eh), and pH govern the availability of nutrients to roots. In an organic matrix, minerals are bound not in harsh synthetic salts, but inside organo-mineral complexes chelated by humic and fulvic polyacids.

Electrochemical Metric	Conventional Inert System	Living Organic Matrix	Functional Implication
CEC (meq / 100g)	2 – 8 (Sand / Rockwool)	30 – 120 (Living Compost / Humus)	High buffering capacity; prevents leaching of Ca, Mg, K, and NH4.
Soil Redox (Eh)	+400 to +600 mV (Oxidized)	+150 to +300 mV (Biologically Active)	Optimum electron availability for microbial respiration without anaerobiosis.
pH Dynamics	Strictly fluctuating (5.5 – 6.5)	Self-buffering (6.2 – 6.8)	Microbial enzymes buffer the rhizosphere independently of input wash water.
Carbon Stabilization	< 0.5% organic matter	5% to 25% stable organic matter	Permanent moisture retention; glomalin-bound micro-aggregates.

2. RHIZOSPHERE MICROBIOLOGY: THE SOIL FOOD WEB ENGINE

The rhizosphere—the immediate millimeters surrounding plant roots—is the most metabolically intense biological zone on the planet. Plants invest between 20% and 40% of their total photosynthetically derived carbohydrates into the soil as root exudates. This is a deliberate biological procurement strategy: plants exchange exudates with specialized microbial guilds in return for solubilized minerals, systemic resistance elicitors, and moisture.

The Four Functional Trophic Guilds

- **1. Plant Growth-Promoting Rhizobacteria (PGPR):** Genera such as *Bacillus*, *Pseudomonas*, and *Azotobacter*. They fix atmospheric nitrogen, synthesize indole-3-acetic acid (IAA) to drive root expansion, and produce siderophores—chelating compounds that bind and deliver insoluble iron (Fe^{3+}) directly to roots.
- **2. Mycorrhizal Fungi (AMF):** Arbuscular mycorrhizal fungi (e.g., *Glomus intraradices*, *G. mosseae*) penetrate root cortical cells, extending the root system's surface area by 1,000x through microscopic hyphal networks. They produce **glomalin**, a hydrophobic glycoprotein responsible for building stable soil crumb structure and long-term carbon sequestration.
- **3. Saprophytic & Nematophagous Fungi:** Decomposers (e.g., *Trichoderma harzianum*, *Chaetomium*) that dismantle complex lignocellulosic residues into humic monomers while parasitizing parasitic nematode eggs and pathogenic fungi like *Fusarium* and *Pythium*.
- **4. Protozoa & Bacterial-Feeding Nematodes:** Amoebae, flagellates, and ciliates consume bacteria and excrete excess nitrogen in plant-available ammonium (NH_4^+) forms directly adjacent to root hairs—the primary mechanism of the "microbial loop".

3. BIOCHEMICAL SOIL BUILDING: MINERALS, HUMUS & CEC

A balanced organic substrate balances physical aeration, moisture retention, biological habitat, and a comprehensive mineral reserve. Under the Albrecht system of Base Saturation, optimal soil health requires a targeted ratio of exchangeable basic cations attached to the soil colloid:

Cation Colloid	Ideal Albrecht Saturation Range	Physiological Role in the Organism	Primary Organic Amendment Source
Calcium (Ca ²⁺)	65% – 70%	Cell wall integrity, pectin synthesis, soil flocculation	Agricultural Gypsum (CaSO ₄), Wollastonite (CaSiO ₃), Calcite
Magnesium (Mg ²⁺)	10% – 12%	Central atom of chlorophyll, ATP enzyme activation	Dolomite Lime (use selectively), Sul-Po-Mag, Epsom Salts
Potassium (K ⁺)	3% – 5%	Osmotic regulation, stomatal mechanics, carbohydrate transport	Langbeinite, Glaucinite (Greensand), Kelp Meal, Wood Ash
Sodium (Na ⁺)	< 1.5%	Non-essential trace ion; high values disrupt soil aggregates	Mitigated via gypsum flushes and humic acid leaching
Hydrogen (H ⁺)	10% – 15%	Maintains balanced soil acidity for trace mineral dissolution	Supplied naturally via organic acid exudates and rainfall

The Humic Fraction Architecture


Humified matter comprises three distinct fractions classified by molecular weight and solubility:

- Fulvic Acids:** Low molecular weight (1,000 – 10,000 Da), pale yellow, soluble in water across all pH ranges. Excellent natural chelator for foliar delivery; capable of penetrating plant cell walls directly.
- Humic Acids:** Medium-to-high molecular weight (50,000 – 100,000 Da), dark brown/black, soluble in alkaline solutions but precipitates in acid. Forms the primary foundation of soil CEC and moisture retention.

- **Humin:** Massive molecular weight, insoluble across all pH ranges. Resists microbial degradation for centuries, forming the physical backbone of ancient organic soils.

4. SECONDARY PLANT METABOLITES: DEFENSE & EXTRACTION CHEMISTRY

While primary metabolites (carbohydrates, amino acids, structural lipids) are required for baseline plant survival, **secondary metabolites** govern the plant's interface with environmental stress, ultraviolet radiation, and herbivorous threats. In organic production systems, avoiding synthetic systemic insecticides forces the plant to ramp up secondary metabolic synthesis, resulting in higher nutritional value, elevated terpene concentrations, and medicinal potency.



Clean Extraction & Processing Dynamics

Extracting secondary botanical metabolites without petrochemical solvent contamination relies on thermodynamic principles:

- **Supercritical Fluid Extraction (SFE - CO2):** At temperatures exceeding 31.1°C and pressures above 73.9 bar, carbon dioxide enters a supercritical state, behaving

simultaneously like a liquid and a gas. Adjusting temperature and pressure parameters selectively isolates volatile terpenes from heavier waxes and polar polyphenols.

- **Hydrodynamic Cavitation & Ultrasonic Extraction (UAE):** High-frequency sound waves (20 kHz to 40 kHz) create microscopic vacuum bubbles in an aqueous or ethanol substrate. The collapse of these bubbles mechanically ruptures plant cell walls, releasing intracellular secondary compounds without heat degradation.
- **Solventless Ice-Water Mechanical Separation:** Freezes non-glandular trichome stalks and capitalizes on differing specific gravities to collect intact glandular heads through graduated micron sieves (25µm to 220µm) without chemical transformation.

5. REGENERATIVE FERMENTATION PROTOCOLS: KNF, JADAM & INOCULANTS

Industrial chemical fertilizers can be replaced using biologically cultured inputs derived from indigenous microbial ecosystems. Originating from Korean Natural Farming (KNF) and JADAM methodologies, these closed-loop protocols allow operators to culture local microbiomes on site:

Protocol 01: Indigenous Microorganisms (IMO-1 through IMO-4)

KNF Standard • Fungal Propagation

1. IMO-1 Collection:

- Cook organic short-grain rice to a firm consistency.
- Place 2 inches of rice in a cedar box covered with breathable washi/paper.
- Bury the box 2 inches deep beneath an undisturbed virgin forest leaf litter canopy (oak, cedar, or bamboo grove).
- Incubate for 3 to 5 days until dense, brilliant white fungal mycelium covers the rice. Avoid black or dull grey molds.

2. IMO-2 Preservation:

- Weigh the IMO-1 rice. Add an equal 1:1 weight of unrefined brown sugar.
- Pack tightly into a glass jar, leaving 1/3 headspace. Cover with breathable cloth.
- Osmotic pressure extracts microbial fluids and stabilizes the culture into a shelf-stable shelf ferment (12 months shelf life).

3. IMO-3 Bulking (Solid-State Fermentation):

- Dilute IMO-2 (1:500) in clean water containing fermented plant juice and humic acid.
- Mix thoroughly into a 50/50 blend of wheat bran and rice hulls until 65% moisture is reached.
- Pile the substrate 30-40 cm high on a dirt floor. Monitor core heat: keep below 50°C (122°F) by turning daily.
- Fermentation is complete when temperature stabilizes and substrate is bound in white hyphal carpets.

4. IMO-4 Soil Integration:

- Blend IMO-3 with virgin local topsoil at a 1:1 ratio.
- Rest for 7 days under cover to acclimate microorganisms to soil minerals before field application.

Protocol 02: JADAM Microbial Solution (JMS) High-Volume Inoculant

Low-Cost Anaerobic/Aerobic Suspension

Materials for 500 Liters of Aqueous Inoculant:

- 500 Liters non-chlorinated water (dechlorinate via bubbler or ascorbic acid).
- 1.5 kg boiled potatoes (mashed to liquid slurry as carbohydrate fuel).
- 500 grams leaf-mold humus gathered from local fertile forest floor.
- 500 grams unrefined rock sea salt (supplying trace mineral spectrum).

Procedure:

1. Suspend the mashed potatoes and forest leaf mold in fine mesh brew sock S.
2. Dissolve the sea salt directly into the water barrel.
3. Submerge the brew bags and hand-massage vigorously to extract microbes and starches into the solution.
4. Cover the barrel and seal loosely. Incubate at 20°C–25°C.
5. Growth curve monitoring: At 24 to 72 hours, a foam circle forms on the surface, reaching a dense peak.
6. APPLICATION WINDOW: Apply immediately when the foam ring is at maximum density before it collapses.

Dilute 1:10 with water for soil drenches to deliver diverse microbial inoculants.

6. COMPOSTING DYNAMICS: THERMOPHILIC VS. STATIC FUNGAL HUMIFICATION

Composting transforms raw biological residues into stable organic matter. The physical and microbiological profile of the finished compost is determined by the decomposition method used:

Parameter	Thermophilic Windrow Composting	Static Fungal Bioreactor (Johnson-Su)	Vermicomposting (Earthworms)
Predominant Organisms	Thermophilic bacteria (<i>Actinobacteria</i>)	Saprophytic & mycorrhizal fungi	Epigeic worms (<i>Eisenia fetida</i>) + microflora
Thermal Profile	55°C – 68°C (Pasteurizing)	Mesophilic throughout (< 35°C after day 5)	Strictly mesophilic (15°C – 25°C)
Turning Frequency	High (turned 5–15 times)	Zero turns (static aeration chimneys)	No turns; passive worm migration
Fungal:Bacterial Ratio	0.1:1 to 0.5:1 (Bacterial dominant)	4:1 to 10:1 (Fungal dominant)	1:1 to 2:1 (Balanced)
Maturation Timeline	6 to 12 weeks	10 to 12 months	12 to 16 weeks
Primary Target Application	High-nitrogen vegetable crops	Perennial soils, orchards, mycorrhizal regeneration	Seedling starts, foliar liquid extraction


The Johnson-Su Bioreactor Construction Parameters

To produce biologically diverse, fungal-dominant compost without turning, build a static bioreactor:

- Form Construction:** Build a cylindrical cage (2 meters tall, 1 meter diameter) using wire fencing lined with landscaping fabric.
- Aeration Chimneys:** Insert four to six 4-inch perforated PVC pipes vertically from the floor pallet to the top of the cage. These ensure oxygen reaches the center via convection, eliminating the need for turning.
- Moisture Maintenance:** Maintain 70% moisture saturation through automated micro-drippers running for 60 seconds once daily.
- Maturation:** Allow to process undisturbed for a full 365 days. The finished material will have converted into a dark, rich, biologically active humic paste.

7. BIOREMEDIATION: MYCOREMEDIATION & PHYTOREMEDIATION

Industrial contaminants, petrochemical spills, and toxic heavy metal accumulations can be systematically remediated using the enzymatic machinery of fungi (mycoremediation) and the bioaccumulation capacity of hyperaccumulating plants (phytoremediation).



Heavy Metal Hyperaccumulation Matrix

Unlike organic petroleum hydrocarbons, heavy metals (Pb, Cd, As, Ni) cannot be degraded into simpler molecules. They must be extracted from the soil matrix via plant root uptake and translocation into harvestable aerial biomass:

Contaminant Ion	Hyperaccumulating Species	Bioaccumulation Index	Disposal / Processing Mechanism
Lead (Pb ²⁺)	Brassica juncea (Indian Mustard)	Up to 1.5% dry leaf mass	Incineration to bio-ore ash; vitrification
Arsenic (As ³⁺ /As ⁵⁺)	Pteris vittata (Brake Fern)	Up to 2.2% dry frond mass	Controlled ashing and closed-loop smelting extraction
Cadmium (Cd ²⁺)	Helianthus annuus (Sunflower)	Exceeds 100 mg/kg dry biomass	Pyrolysis to biochar; immobilizes ions in carbon cages
Radionuclides (Cs-137, Sr-90)	Cannabis sativa L. (Industrial Hemp)	Deep taproot translocation	Gasification inside closed hazardous containment vessels

8. BIOMATERIALS & CLOSED-LOOP INDUSTRIAL ORGANICS

Modern organics extends beyond agriculture into advanced structural materials, replacing petroleum-derived polymers with biological alternatives:

1. Mycelium Composite Structural Cores

Agricultural byproducts (hemp hurd, sawdust, oat hulls) are inoculated with the vegetative mycelium of bracket fungi such as *Ganoderma sessile* or *Fomes fomentarius*. Over 7 to 14 days in darkness, the fungal hyphae branch through the substrate, binding it into an interconnected composite matrix.

- **Thermal Pressing & Desiccation:** Once fully colonized, the material is baked at 90°C to denature the fungus, halting growth. Compression pressing yields lightweight, fire-retardant structural panels with acoustic and thermal insulation values exceeding expanded polystyrene (EPS).

2. Chitin & Chitosan Deacetylation

Chitin—the structural polymer found in crustacean shells and insect exoskeletons (e.g., black soldier fly pupae)—is deacetylated using sodium hydroxide (or enzymatic processing via chitin deacetylases) into **chitosan**.

- **Industrial Functions:** Chitosan forms non-toxic, antimicrobial packaging films, acts as an organic flocculant for settling suspended particles in industrial wastewater, and triggers an immune response in plants by simulating insect predation, stimulating natural pest defenses.

9. THE MASTER FIELD RECIPE & FORMULATION REPOSITORY

Formulation 01: Super Soil Substrate (1 Cubic Yard Master Blend)

Landry Industries
Specification

Base Medium:

- 1/3 Sphagnum Peat Moss (or washed coco coir)
- 1/3 Aeration Grade Perlite / Pumice / Lava Rock (3/8" grading)
- 1/3 Johnson-Su Compost & Thermal Earthworm Castings (50/50 blend)

Mineral & Biological Amendments (Per Cubic Yard):

- 8 kg Glacial Rock Dust (paramagnetic broad trace minerals)
- 4 kg Agricultural Gypsum (calcium sulfate – unpelletized)
- 2 kg Wollastonite (calcium silicate – plant available silica)
- 2 kg Basalt Meal (high paramagnetic volcanic force)
- 1.5 kg Crustacean Meal (cracked crab/shrimp shell – chitin source)
- 1.5 kg Kelp Meal (Ascophyllum nodosum – natural cytokinins & trace potassium)
- 1 kg Alfalfa Meal (natural triacontanol growth hormone)
- 500 grams Steamed Bone Meal (slow-release biological phosphorus)
- 250 grams Pure Leonardite-derived Humic Acid Powder

Conditioning Phase:

Moisten blend with rainwater inoculated with JMS or compost tea until 60% field capacity is reached.

Cover with a tarp and allow to "cook" for 30 to 45 days, allowing the microbial flora to break down the raw amendments before planting.

Formulation 02: Actively Aerated Compost Tea (AACT) Brewer

50-Gallon Foliar & Drench Protocol

Equipment:

- 55-gallon food-grade drum.
- High-output commercial diaphragm air pump (minimum 80 LPM).
- Large-pore air diffuser (fine-bubble airstones clog rapidly with bio-films).
- 400-micron nylon brew bag.

Brew Ingredients:

- 50 Gallons de-chlorinated clean spring water (Temp: 18°C – 22°C).
- 2.5 kg Fresh Fungal-Dominant Compost / Worm Castings.
- 60 mL Unsulphured Blackstrap Molasses (bacterial carbohydrate fuel).
- 60 mL Liquid Kelp Extract (fungal food and trace minerals).
- 30 mL Liquid Humic Acids (hyphal growth stimulant).
- 30 grams Micronized Soluble Fish Hydrolysate (cold-processed enzymatically).

Operational Process:

1. Dissolve molasses, kelp, humic acid, and fish hydrolysate into the water column.
2. Turn on the air pump and verify vigorous dissolved oxygen agitation (target > 8.0 mg/L DO).
3. Place castings/compost inside the nylon mesh bag and submerge into the bubbling vortex.
4. Aerate continuously for 24 to 36 hours.
5. Verification: A rich, yeasty, forest-floor aroma and a thick head of biological foam indicates a successful brew.
6. Apply within 4 hours of air cut-off; do not seal in airtight containers.

10. QUALITY ASSURANCE, CHROMATOGRAPHY & FIELD CHECKLIST

Ensuring inputs meet safety and potency standards requires systematic testing:

Pfeiffer Circular Chromatography

A field-deployable qualitative assay used to evaluate living biological systems:

- A soil or compost sample is extracted in a 1% sodium hydroxide (NaOH) solution.

- The solution is wicked onto circular filter paper (Whatman No. 1) sensitized with silver nitrate (AgNO₃).
- **Reading the Pattern:** Dead soils produce flat, dark, concentric bands. Healthy soils produce intricate radiating patterns with feathered edges and warm copper-gold tones, reflecting the integration of minerals, organic matter, and microbial activity.

Standard Operating Procedure Verification Checklist

- [] **Base Saturation Audit:** Albrecht exchange balance verified (Ca: 65–70%, Mg: 10–12%, K: 3–5%).
- [] **Water Quality Check:** Total dissolved solids (TDS) < 150 ppm; chlorine and chloramines neutralized.
- [] **Biological Inoculation:** Substrates primed with local forest-derived mycorrhizae or diverse IMO cultures.
- [] **Heavy Metal Screening:** Compost stocks screened via ICP-MS for cadmium, lead, arsenic, and mercury below safety limits.
- [] **Aeration Balance:** Substrate maintains at least 30% air-filled porosity at field capacity to prevent anaerobiosis.
- [] **Pesticide Screening:** Bio-inputs verified clean of persistent synthetic pyridine and pyralid herbicides.
- [] **Extract Stability:** Aerated biological teas applied within 4 hours of brewing to protect aerobic microbes.