One density surface, two kinds of data

A joint spatial model for marine mammal eDNA and visual line-transect surveys

Eiren Jacobson and the MMARINeDNA team

SMM 2026

Two ways to count whales


	Visual line transects	eDNA
What is recorded	Groups seen, their size and distance from the line	DNA copies in a filtered water sample
What it tells us about density	Absolute: animals per km^2, after correcting for detection	Relative: copies depend on shedding, decay and transport
Where it works	Daylight, calm seas, animals at the surface	Any weather; samples at depth; species that are hard to see
Where it struggles	Rare or cryptic species, rough seas	Turning copies into animals

Project goals

- **One density surface per species**, informed by both data streams at once
- **Calibrate eDNA against animals:** let visual data set the scale that links DNA copies to density
- **Fill gaps:** eDNA where there is little visual effort, or for species rarely seen
- **Honest uncertainty:** each data source keeps its own observation model
- **Test on simulations first**, where the truth is known

The data

Two surveys, one coast



Visual line transects (2018 CCES)


- 7,561 km of effort in 1,641 segments
- Perpendicular distances, group sizes, Beaufort sea state

eDNA (2019)

- 551 water bottles, 0–500 m depth
- **qPCR** for Pacific hake
- **Metabarcoding** (MarVer1) for all vertebrates


Study area: north of 35°N, a 690 × 1,770 km box

Humpback whale: easy to see, rare in eDNA




- 548 groups detected within 6.5 km of the line
- Reads in 36 of 551 bottles
- Visual data are rich; eDNA hits are sparse but broadly match the sighting hot spots

Pacific white-sided dolphin: weak in both



- 89 groups detected within 4.6 km of the line
- Reads in 60 of 551 bottles
- Few sightings, but large groups (up to hundreds)
- This is where combining sources should help most

Pacific hake: the eDNA anchor



- qPCR detects hake in 527 of 551 bottles
- Metabarcoding reads in 447 bottles
- Not surveyed visually
- Abundant and eDNA-rich, so it pins down the **shared eDNA noise terms** (bottle-to-bottle variation, non-target DNA)

The model

The core idea: one latent field per species

$$\log \lambda_s(\mathbf{x}) \;=\; \underbrace{\mu_s}_{\text{intercept}} \;+\; \underbrace{f_s(X, Y)}_{\text{spatial GP}} \;+\; \underbrace{h_s(Z_{\text{bottom}})}_{\text{depth spline}} \qquad \text{animals per km}^2$$

eDNA: expected copies in bottle i

$$\mu_{is} = \lambda_s(\mathbf{x}_i) \times c_s \times v_i \times \omega_s(z_i)$$


conversion factor c_s, litres filtered v_i, water-column offset ω_s

Visual: expected groups on segment j

$$\mathbb{E}[n_j] = \frac{\lambda_s(\mathbf{x}_j)}{\mathbb{E}[\text{group size}_s]} \times 2L_j \times \text{ESW}_j$$

segment length L_j, effective strip half-width ESW_j


The latent field: a Hilbert-space Gaussian process



- **Anisotropic GP:** separate length-scales east–west and along the coast
- **HSGP approximation:** the GP is a weighted sum of fixed sine waves (up to 800), so the sampler fits hundreds of weights instead of an $N \times N$ covariance


- **Bottom depth:** smooth spline, 4 df
- **Priors from the survey design only:** sample spacing, domain size and basis budget, never the counts
- Fitted in **Stan**

eDNA observation model




- **Bottle copies** $C_{is} \sim \mathrm{NB}(\mu_{is}, \theta)$: bottles vary even at the same density
- **Non-target "junk" DNA** J_i, with its own spread θ_{junk}
- **qPCR** (hake): detection with probability $1 - e^{-\kappa C}$; Ct falls linearly with $\ln C$
- **Metabarcoding**: reads split among species and junk in proportion to copies; zero-inflated beta-binomial
- Replicates share their bottle's copies

Visual observation model



- **Encounters:** groups per segment are Poisson with mean $\lambda / \mathbb{E}[s] \times 2L \times \text{ESW}$
- **Distances:** half-normal or hazard-rate detection, truncated at w; $\log \sigma = \log \sigma_0 + \beta_{\text{size}}(s - \bar{s}) + \beta_{\text{bft}}(\text{Bft} - 3)$
- **Group size:** modelled, with size-bias correction
- Same field, units and priors as the eDNA side
- $g(0) = 1$ assumed (for now)

The joint model



- **One field per species** feeds both observation models
- **Switches** turn either likelihood off: eDNA-only, visual-only and joint fits share priors, basis and starting values, and differ only in data
- With one side switched off, it reproduces the single-source model exactly
- **Conversion factors** c_s can be **estimated** for the cetaceans

Why combining should help

- **eDNA alone** sees only $\lambda_s \times c_s$, so it can map *relative* density but cannot separate density from the conversion factor
- **Visual data** fix the absolute scale of λ_s, so the joint model can **estimate** c_s: copies per unit of animal density
- **eDNA** adds information where visual effort is thin or sightings are few, sharpening the surface
- **Shared noise terms**, learned mostly from hake, carry over to the rare whale signals

Simulation study

Simulation design

Species	eDNA	Visual detections	Role
Pacific hake	qPCR + metabarcoding	not surveyed	eDNA-rich anchor
Humpback whale	metabarcoding	~800–1,500	rare in eDNA, conspicuous at the surface
Pacific white-sided dolphin	metabarcoding	~100	weak in both

- Study box north of 35°N, 690 × 1,770 km
- 200 eDNA stations; 35 east–west transects in 10 km segments
- True length-scales: 50 km east–west, 150 km along the coast

- Four fits per dataset: **eDNA-only, visual-only, joint, joint with c_s free**
- Scored on field R^2, 95% interval width and coverage, and recovery of c_s

Results

Placeholder: simulation results to come (joint comparison now running on the server, seeds 2–5)

- Figure: posterior mean against true log density, by species (rows) and fit (columns), from `outputs/joint/compare_s<SEED>/compare_joint.png`
- Table: field R^2 (95% interval width) per species and fit, from `compare_field.csv`
- Conversion-factor recovery for humpback and PWSD (joint, c_s free), from `compare_recovery.csv`

Species	eDNA only	Visual only	Joint	Joint, c_s free
Hake	...	not surveyed	...	...
Humpback	...	...	...	...
PWSD	...	...	...	...

Next steps

- **Real data:** fit the joint model to the 2018 sightings and 2019 eDNA
- **eDNA inputs:** qPCR Ct values and standard curves; volumes filtered
- **Detection:** Beaufort covariate in the joint model's visual side; $g(0)$
- **Different years:** the surveys are a year apart, so the joint fit assumes a stable density surface
- **eDNA transport and decay:** where the DNA is is not quite where the animal is

Thank you

https://github.com/MMARINeDNA/eDNAVisualJointModel

Riutort-Mayol *et al.* (2023) Practical Hilbert space approximate Bayesian Gaussian processes for probabilistic programming. *Statistics and Computing* 33:17. Buckland *et al.* (2001) *Introduction to Distance Sampling*. Oxford University Press.