Apollo Imperial HOSPITALS

DIAGNOSTIC REPORT *ID12609508463*

MRN	: IHL1037895	Order No	: ID12609508463
Patient Name	: M MOHSIN CHOWDHURY	Collected	: 09/09/26 12:25 AM
Age / Gender	: 59Y 1M 29D / Male	Received	: 09/09/26 12:51 AM
Ref. Doctor	: DR. DANIA RAHMAN	Reported	: 09/09/26 02:03 AM
Report No	: 12609142170	BED No	: 2-DB-33
LAB ID	: 12609876178	Status	: FINAL

HAEMATOLOGY

COAGULATION PROFILE

Parameter	Result	Unit	Flag	Bio.Ref.Interval	Method
PROTHROMBIN TIME					
Control Value (PT)	12.5	Sec			Photo Optical
Patient value (PT)	11.9	Sec		10.00 - 14.00	Photo Optical
Prothrombin Ratio (PR)	0.95				
International Normalized Ratio(INR)	1.04				
Index	105.04	%			

Interpretive Guide

Intended Use
Screening to identify a deficiency of 1 or more of the clotting factors of the extrinsic coagulation system (I, II, V, VII, X) due to hereditary deficiency or acquired conditions such as liver disease, vitamin K deficiency, or a specific factor inhibitor.
Monitoring patients on oral anticoagulant therapy to maintain a patient in a safe therapeutic range

Clinical Information
This assay is used to monitor oral anticoagulant therapy to maintain a patient in safe therapeutic range In the absence of oral anticoagulant therapy, a prolonged prothrombin time indicates deficiency of 1 or more factors(I, II, V, VII or X) or the presence of a coagulation inhibitor.The prothrombin time is also reported as the INR (international Normalized Ratio), based on the ISI (International Sensitivity Index) assigned to the thromboplastin and coagulometer. This assay is used for monitoring "stable" oral anticoagulation. A mixing test of patient and normal plasma (1:2) can be performed, if indicated, to differentiate coagulation factor from inhibition.

Interpretation
Prolongation of the prothrombin time (PT) can occur as a result of deficiency of 1 or more coagulation factors(acquired or congenital in origin) , or the presence of an inhibitor of coagulation such as heparin, a lupus anticoagulant , a "nonspecific" inhibitor such as a monoclonal immunoglobulin, or a specific coagulation factor inhibitor.
PT Mixing study, using equal volume patient and normal pool plasma, may be performed on specimens with a prolonged PT to assist in differentiating coagulation factor deficiencies from coagulation inhibitors. Correction of the PT mix to within the normal reference range usually indicates a coagulation factor deficiency. If the prolonged PT is due to an inhibitor,the PT mix typically fails to correct a prolonged PT.

Cautions
Prothrombin time (PT) mixing studies have no utility when the PT is normal.
Lipemic specimens may interfere with the instrument clot detection mechanism.

Legend: H-High, HC-High Critical, L-Low, LC- Low Critical

---------- End of Report ----------

Powered by: MYSOFT Ltd.

MR. ROBIN DAS
DMT, B.Sc. in Medical Technology
Medical Technologist

DR. RAHAT ANJUM
MBBS, MCPS(CL.PATH), M.PHIL(PATH)
ASSOCIATE CONSULTANT,BMDC. Reg .No :A-44688

Estate of Imperial Hospital | Zakir Hossain Road, Pahartali, Chattogram, Bangladesh.
+88 09612 247 247 | www.aihlbd.org | info@aihlbd.org