

# THE IMPRINT OF WATER-COLUMN RESONANCE ON SECONDARY MICROSEISMIC SOURCE SPECTRA

M Meschede<sup>1</sup>, É Stutzmann<sup>1</sup>, V Farra<sup>1</sup>, M Schimmel<sup>2</sup>, F Ardhuin<sup>3</sup>


<sup>1</sup> Institut de Physique du Globe de Paris, Paris 75005, France

<sup>2</sup> Institut of Earth Sciences Jaume Almera - CSIC, 08028 Barcelona, Spain


<sup>3</sup> University of Brest, CNRS, IRD, Ifremer, Laboratoire d'Océanographie Physique et Spatiale, IUEM



# THE SECONDARY MICROSEISMIC FREQUENCY BAND




# P-WAVE GENERATION FROM INTERACTING OCEAN WAVES




inspired by Gualtieri et al. (2014, 2015) and Ardhuin et al. (2013)

# SECONDARY MICROSEISM PRESSURE MODELS



[model by Ardhuin et al. (2011), distributed by IFREMER]

# SECONDARY MICROSEISM PRESSURE MODELS




[model by Ardhuin et al. (2011), distributed by IFREMER]



## BUILDING A DATASET





1 year of continuous data per array





P-wave (30-90 degree) coverage

- five arrays (North-America)
- combined timespan of five years (2006, 2008, 2011, 2015, 2015)
- daily synthetic and observed beams for 3 models (1809 beams as a function of slowness and frequency)
- this presentation:  
spectral characteristics of the strongest-sources



# FROM PRESSURE MODEL TO SYNTHETIC BEAM



COMPARISON OF BACKPROJECTED BEAMS




California array



Alaska array




# THE STRONGEST-SOURCE DISTRIBUTION




## SELECTING COMPARABLE SOURCES

observed source time and location ...






... match with model R



... don't match with model R

- On 26% (467/1809) days the strongest-sources match
- non-matching sources are mostly microseisms as well

# THE SPECTRAL DOUBLE PEAK




- Spectral double peak in observation and models with resonance
- Amplitudes are predicted within factor of 0.5 - 6 (1 std)
- peak frequencies are either low or high




# ASSOCIATING SPECTRAL SHAPES WITH GEOGRAPHICAL REGIONS

## low frequency cluster



## high frequency cluster


## mid frequency cluster



- Spectral Shape and Geographical Region are related
- Low frequency peak is broader than the high frequency one
- High frequency spectra are better modeled





# SOURCE SITE RESONANCE




- resonance when  $2h/\lambda + 1/2 = n$
- $h = 1/4\lambda, 3/4\lambda, 5/4\lambda, \dots$
- $n = 1$  (shallow ocean) and  $n = 2$  (deep ocean) most important for secondary microseisms

## SOURCE SITE RESONANCE MAP



- $n=1$  in shallow ocean [cross]
- no resonance from 0.10 - 0.24Hz [circle]
- $n=2$  in deep ocean [triangle]

## CONCLUSIONS



- secondary microseism P-wave spectra are strongly affected by resonance effects
- bathymetry favours resonance at 0.15 and 0.20Hz (first and second harmonic)
- ocean wave model can reasonably well predict the spectra