

# He Is Everywhere

Input file:            **standard input**  
Output file:           **standard output**  
Time limit:            1 second  
Memory limit:         256 megabytes

Real Madrid's past coach **Carlo Ancelotti** has stopped asking **Camavinga** where he wants to play.

**Why?** Because the answer is simple: **yes of course coach**

Left-back ? Yes.. Midfilder? Yes.. Right-wing? Yes.. Go get me some gum ? Yes. meanwhile **Bellingham** mostly stays where he should, **Vinicus** mostly stays on the wing.

Before the match, the coach writes down Camavinga's official position because of the official operations. but during the match, the analyst keeps spotting him all over the field.

Your mission is simple.. given  $N$  positions the analyst recorded for Camavinga during the match and his official position  $P$ . If Camavinga played in **any position different** from the official one, then he was **"everywhere"**.

## Input

The first line contains two integers  $N$  and  $P$  ( $1 \leq N \leq 20$ ) the number of positions that the analyst recorded for Camavinga during the match and his official position.

The second line contains  $N$  strings. Camavinga recorded positions during the match.

## Output

If Camavinga played in any position different from his official one, then print **"he is everywhere"** or print **"something wrong, Camavinga is always everywhere"** if he only plays in his official position.

## Examples

| standard input         | standard output                                 |
|------------------------|-------------------------------------------------|
| 4 CB<br>CB CB CB CB    | something wrong, Camavinga is always everywhere |
| 5 LM<br>CB GK SS CB LM | he is everywhere                                |