




# PHY 110 Engineering Physics

---

Lecture 1  
UNIT 2 – laser

# Lasers and applications:


- Fundamentals of laser- energy levels in atoms
- Radiation matter interaction
- Absorption of light
- Spontaneous emission of light
- Stimulated emission of light
- Population of energy levels
- Einstein A and B coefficients
- Metastable state
- Population inversion,
- Resonant cavity
- Excitation mechanisms
- Nd – YAG
- He-Ne Laser
- Semiconductor Laser
- lasing action
- Properties of laser
- Applications of laser: holography



# History:



- **1917-** Einstein predicted the possibility of Stimulated emission.
- **1954-** C.H.Towens developed MASER (Microwave Amplification through Stimulated emission)
- **1958-** A.Schawlow and C.H.Towens extended principle of Maser to light.
- **1960-** T.H.Maiman built first laser.

**LASER**( **L**ight **A**mplification through **S**timulated **E**mission of **R**adiation)



# Energy levels in an atom:

Every atom has a ground level with minimum energy (Ground state,  $n=1$ ). All the levels above it is called excited states.



# Radiation-Matter interaction

9

In physics, **radiation** is the emission or transmission of energy in the form of waves or particles (photon) through space or through a material medium

light is nothing but radiation, thus will act as either a wave or a particle (photon); both of which are important in its interaction with **matter**.


**matter** consists of various types of particles, each with mass and size. The most familiar examples of material particles are the *electron, the proton and the neutron*. Combinations of these particles form atoms; bonding of atoms molecules and that extend we have matter...

When a photon travel through a material three different processes occur, namely, **absorption, spontaneous emission and stimulated emission**. We study these 3 processes now...

# 1. ABSORPTION

10

Suppose an atom is in the lower energy level  $E_1$ . A photon of energy  $h\nu = (E_2 - E_1)$  interacts with the atom and gets annihilated. So atom actually absorbs that photon and goes to the next allowed level  $E_2$  and this transition is called absorption transition or induced absorption



$A + h\nu = A^*$  Where A is an atom in the ground state and  $A^*$  is an excited

# 1. ABSORPTION

11

The **probability** of an absorption transition ( $P_{12}$ ) is proportional to photon density  $\rho(\nu)$

$$P_{12} \propto \rho(\nu) \quad P_{12} = B_{12} \rho(\nu)$$

$B_{12}$  is known as the Einstein coefficient for induced absorption, and indicates the probability of occurrence of an induced transition from  $1 \rightarrow 2$ .  $B_{12}$  characteristics of an atom as per the properties of  $E_1$  and  $E_2$ .

If  $N_1$  and  $N_2$  are the population of atoms at  $E_1$  and  $E_2$  levels in the material, then **Rate of absorption transitions** ( $R_{\text{abs}}$ ) is defined as

$$R_{\text{abs}} = -\frac{dN_1}{dt} = \frac{dN_2}{dt}$$



## 1. ABSORPTION

$R_{\text{abs}}$  is also equal to the product of probability of absorption transition and number of atoms at the energy level 1

$$R_{\text{abs}} = P_{12} N_1 = B_{12} \rho(\nu) N_1$$

i.e. absorption transition rate,  $R_{\text{abs}}$  when the  $N_1$  is more or photons density is more, absorption transitions from level 1  $\rightarrow$  level 2 are more

As it is expected, due to this absorption transition  $N_1$  decreases and  $N_2$  increases. But under normal conditions  $N_2$  can not be greater than  $N_1$ ...


**That we will see later in Population inversion for lasing action**



## 2. SPONTANEOUS EMISSION

13

For this to happen there should be atoms in the excited state  $E_2$



But in the excited state  $E_2$ , atoms cannot stay longer. In about  $10^{-8}$  seconds atoms come back to the lower level  $E_1$  by releasing a photon of energy  $h\nu = (E_2 - E_1)$ . This photon emission without any external aid is called **spontaneous emission**.

## 2. SPONTANEOUS EMISSION

14

Spontaneous emission is the reverse process of induced absorption transition and can be expressed as  $A^* \rightarrow A + h\nu$

Spontaneous emission is independent of the photon density and depends only on  $E_2$  and  $E_1$  properties. Hence probability of spontaneous transition  $(P_{21})_{\text{Spont}}$  from  $2 \rightarrow 1$

$$(P_{21})_{\text{Spont}} = A_{21}$$

Where  $A_{21}$  is known as the Einstein coefficient for Spontaneous emission, characteristic of the atom. Rate of spontaneous transition  $R_{\text{sp}}$

$$R_{\text{sp}} = P_{21}N_2 = A_{21}N_2$$

## 2. SPONTANEOUS EMISSION

15

In other words,  $A_{21}$  is a measure of the lifetime of atoms at  $E_2$  state against the spontaneous emission and subsequent transition of atom to the  $E_1$ . Note; spontaneous transition is not possible from  $E_1$  to  $E_2$  so

$$A_{12} = 0$$

### Characteristics of the Spontaneous Emission

1. **No way to control** the process from outside
2. Direction of propagation, phase and plane of polarization of emitted photons are random - **Incoherent**
3. Light emitted are **polychromatic** not monochromatic
4. The intensity of the emitted radiation **is proportional to the number radiating atoms(N)**.. i.e excited atoms at level  $E_2$

$I_{\text{tot}} = N I$ , where  $I$  is the intensity of light emitted by one atom.

**Spontaneous emission is what we see in conventional light sources.**

### 3. STIMULATED EMISSION

16

From the previous discussions it is clear rate of absorption transition ( $R_{\text{abs}}$ ) is higher compared to rate of spontaneous transition ( $R_{\text{spont}}$ ) and due to that  $N_2$  can be higher than  $N_1$  with passage of time. But **this wont happen** and an equilibrium is maintained. **But we can make it☺, population inversion, surely not by normal means**


So to account this Einstein came up with the brilliant idea that say that a photon can stimulate an atom in the excited state  $E_2$  to release a photon and come back to the lower energy state  $E_1$ . This mechanism is called stimulated emission and is **depended on the photon density** present in the medium. And like before can be expressed as





### 3. STIMULATED EMISSION

17



Atoms in the excited state interact with the photon in the material medium and without consuming that photon, it get the stimulation to produce another identical one. As a result two photon are generated now. **A different light source with more intensity.**

### 3. STIMULATED EMISSION

18

The probability with which a stimulated transition occurs is given by

$$(P_{21})_{\text{stimu}} \propto \rho(\nu)$$

$$(P_{21})_{\text{stimu}} = B_{21} \rho(\nu)$$

$B_{21}$  is known as the Einstein coefficient for **induced/stimulated** transition, and indicates the probability of occurrence of an induced transition from  $2 \rightarrow 1$ .  $B_{21}$  characteristics of an atom as per the properties of  $E_1$  and  $E_2$ .

Similarly, Rate of stimulated emission ( $R_{\text{st}}$ ) is expressed in terms of photon density and the number of atoms ( $N_2$ ) at the excited state

$$R_{\text{st}} = (P_{21})_{\text{st}} N_2 = B_{21} \rho(\nu) N_2$$

# CHARACTERISTICS OF THE STIMULATED EMISSION

19

- 1) The process **is controllable from outside-** control it through the photon interaction
- 2) Induced and inducing photon propagate in the **same direction**
- 3) Identical to incident photon-**coherent**
- 4) Multiplication of the photon**
- 5) Light amplification**
- 6) Net intensity of light is proportional to the square of the number of atoms (N) radiating light..

$I_{\text{tot}} = N^2 I$ , where  $I$  is the intensity of light emitted by one atom.