

SMART INDIA HACKATHON 2026




TITLE PAGE

SagarDrishti सागर दृष्टि

A browser-native 3D digital twin of India's ocean

- **Problem Statement ID** - SIH26067
- **Problem Statement Title** - Web-based 3D Visualization Platform for Ocean Model Data and In-situ Observations
- **Theme** - Disaster Management
- **PS Category** - Software
- **Team ID** - [TEAM ID]
- **Team Name (Registered on portal)** - [TEAM NAME]



Running today, offline: 24 depth levels from 5 m to 2000 m, the 26 degC isosurface, 337 current vectors and CAP warning areas.

Not a proposal on paper. 543 automated tests · 51 fps at 1080p on integrated graphics · 10.5 M gridded values · 16,390 QC-passed instrument levels · zero requests that leave the machine. Every figure in this deck was measured, not estimated.



IDEA TITLE



SagarDrishti सागर दृष्टि

One screen where INCOIS's 3D model fields and the instruments in the water finally meet, with a number saying how far apart they are.

THE PROBLEM · INCOIS'S OWN WORDS

- **“No integrated, web-based 3D visualization platform currently exists”** that renders model fields and in-situ observations together.
- **Forecasters are “forced to toggle between disparate software packages”**
- **which “impedes timely hazard assessment, search-and-rescue support, fishery advisories”**

PROPOSED SOLUTION · HOW IT ADDRESSES THE PROBLEM

- **Volumetric water column in a browser.** 24 depth levels, 5 m to 2000 m, as blended depth slices, extracted isosurfaces and time animation. No client install.
- **Instruments fused into the same scene.** 25 real casts from 17 instruments; click one to read its measured profile against the model at that point and time.
- **A skill number beside the field.** Class-4-style verification prints bias and RMSE per depth band, with the caveat that limits it, on the same screen.
- **Hazards drawn over the water they concern.** Live CAP v1.2 from NDMA SACHET, India's national alert backbone.
- **A new source is a config entry, not a rewrite.** YAML registry plus a plugin interface; OGC WMS/WCS and CF served.

INNOVATION AND UNIQUENESS

Skill, not just pictures

The only viewer that prints how wrong the model is, beside it

Air-gapped by construction

Zero off-machine requests; a test fails the build if one appears

An agent that cannot invent


Code withholds any answer stating a number no tool produced

Derived fields as plugins

Density (TEOS-10) and cyclone-heat depth, no change to the store



One scene: model field, instrument casts, hazard areas, the verification card and a measured profile against the model.



Printed on the running screen, beside the field itself.

PROOF ON SCREEN


- **11,718 matched pairs** from 23 real casts, per depth band
- **13 ocean parameters** including oxygen, chlorophyll-a, nitrate and pH
- **The caveat is printed in red** analysis fit, NOT forecast skill



TECHNICAL APPROACH



NetCDF · ASCII · CAP XML → source registry (YAML) → chunked Zarr cube → FastAPI + OGC WMS / WCS → WebGL 3D scene in the browser





FEASIBILITY AND VIABILITY



FEASIBILITY IS NOT A FORECAST HERE. THE SYSTEM ALREADY RUNS. EVERY FIGURE BELOW WAS MEASURED, NOT ESTIMATED.

543

automated tests, 10 of them driving a real browser

51 fps

at 1080p on integrated Intel graphics, not a gaming GPU

0

requests leave the machine; a test fails the build if one does

10.5 M

gridded values in the demo cube from two agencies

16,390

QC-passed instrument levels from 17 real instruments

13

ocean parameters served, 2 of them computed by plugins

ANALYSIS OF FEASIBILITY

TECHNICAL

- **Proven stack.** CesiumJS, FastAPI, xarray and Zarr are mature and open; nothing here needs inventing.
- **Frame budget cleared.** 51 fps median on the weakest target GPU, against a 30 fps floor.

OPERATIONAL

- **Reads INCOIS's own ERDDAP.** Zero data duplication; the agency keeps its single source of truth.
- **No client install.** A browser is the whole deployment surface.

ECONOMIC

- **Zero licence cost.** Fully open source, one commodity GPU server.
- **Skills exist in-house.** Standard Python and JavaScript; INCOIS already runs ERDDAP and LAS.

POTENTIAL CHALLENGES AND RISKS

DATA

- **Instrument coverage is uneven.** Glider and CTD files are not published on INCOIS ERDDAP.
- **Datasets disagree on axes.** GLORYS stops at 1942 m where the INCOIS field has a 2000 m level.

TECHNICAL

- **Volumetric rendering is costly.** Translucent overdraw is the dominant frame cost.
- **Scaling to the full basin.** The national grid is far larger than the demo box.

ADOPTION AND TRUST

- **A wrong number is worse than none.** A skill figure quoted without its caveat would mislead.
- **A false alarm is unacceptable.** A drill must never read as a live warning.

STRATEGIES TO OVERCOME THEM

DATA

- **Parsers already written and tested.** Glider and CTD ingest needs a file, not code.
- **Never stretch one grid onto another.** The served depth is printed beside the requested one.

TECHNICAL

- **Depth-space level of detail.** At most 10 of 24 slices blended at once; slices-only is the proven floor.
- **Chunked Zarr over range requests.** Cost is view-dependent, not dataset-dependent.

ADOPTION AND TRUST

- **The caveat travels with the number.** Printed on screen, in the API and in every agent answer.
- **Four independent drill guards.** CAP status Exercise is refused, dashed, stamped and spoken.

VIABILITY · WHY THIS SURVIVES AFTER THE HACKATHON

Native deployment path

The PS is INCOIS's own tooling request. We read their ERDDAP, so there is no data to migrate and no second source of truth.

Standards, not a silo

OGC WMS 1.3.0 and WCS 1.0.0 mean their existing desktop tools can consume this without us writing a client for them.

Maintainable by the owner

Standard Python and JavaScript, a documented plugin registry, and a test suite that fails loudly. INCOIS already runs ERDDAP and LAS.

Costs nothing to keep

Zero licences, one commodity GPU server, and a browser as the entire client-side footprint.



IMPACT AND BENEFITS



POTENTIAL IMPACT ON THE TARGET AUDIENCE

DISASTER MANAGERS

INCOIS runs India's Tsunami Early Warning Centre. A CAP bulletin stops being text and becomes a shape on the water, beside the field that explains it.

OPERATIONAL FORECASTERS

The daily question "can I trust today's field?" is answered per depth band on the same screen, in seconds rather than a separate offline report.

SEARCH AND RESCUE

Depth-resolved currents, not just the surface, which is what a drifting object actually follows.

FISHERY ADVISORIES

Potential Fishing Zone advisories reach about 9 lakh fisherfolk households; subsurface temperature and oxygen are the inputs behind them.

RESEARCH, EDUCATION, OUTREACH

The PS names public outreach and e-learning. Guided tours narrate the ocean for schools and exhibitions from the same engine.

BENEFITS · SOCIAL, ECONOMIC, ENVIRONMENTAL

- **Faster hazard assessment.** One tool replaces the toggling the PS describes; the model and the instruments are already side by side.
- **Verifiable, not decorative.** Bias +0.026 degC and RMSE 0.602 degC over 11,718 matched pairs, with the analysis-fit caveat attached.
- **Trust where it matters most.** The model is within about 0.12 degC below 500 m and 2.07 degC across the thermocline at 50 to 100 m, which is exactly the layer a cyclone forecaster reads.

- **Zero licence cost.** Fully open source against proprietary 3D suites; one deployment serves the nation.
- **Resilient by design.** Runs air-gapped, so a network failure at a control room or a nodal centre does not take the tool down.
- **Wider access.** A browser and, in the next phase, Indian-language voice widen who can interrogate the ocean beyond specialists.

NATIONAL ALIGNMENT

Digital India

Deep Ocean Mission

Blue Economy

SAGAR / MoES mandate

UN SDG 14: Life Below Water

Deployable on INCOIS infrastructure

The PS is an internal INCOIS tooling request, so the deployment story is native rather than argued.

WHAT CHANGES ON THE DAY IT MATTERS · A DEPRESSION FORMS IN THE BAY

1 SEE THE HEAT

2 CHECK THE MODEL

3 CONFIRM AGAINST THE WATER

4 PUT THE WARNING IN CONTEXT



[TEAM
NAME]

RESEARCH AND REFERENCES

DETAILS AND LINKS OF THE REFERENCE AND RESEARCH WORK

METHOD AND VERIFICATION

- [Ryan et al. 2015: GODAE OceanView Class 4 verification, J. Op. Oceanogr. 8](#) · *the method behind our scorecard*
- [Wong et al. 2020: Argo data and its QC flags, Front. Mar. Sci. 7:700](#) · *why we keep only flags 1 and 2*
- [IOC, SCOR, IAPSO: TEOS-10 thermodynamic equation of seawater](#) · *our density field*
- [Balakrishnan Nair et al. 2013: INCOIS Ocean State Forecast validation, Current Science 105\(2\)](#) · *the Indian validation precedent*

WEB 3D OCEAN VISUALISATION

- [Qin et al. 2021: Web 3D framework for time-varying ocean forecast data, Env. Modelling & Software 135](#) · *the closest ancestor*
- [Yu, Qin & Xu 2025: WebGPU volume rendering of ocean scalars, Applied Sciences 15\(5\):2782](#) · *our rendering optimisations*
- [Tucker et al. 2020: Argovis, J. Atmos. Ocean. Tech. 37\(3\):401](#) · *the API contract we follow*
- [Blower et al. 2013: ncWMS, Env. Modelling & Software 47](#) · *OGC WMS over CF-NetCDF*
- [Signell & Pothina 2019: Cloud coastal model analysis, JMSE 7\(4\):110](#) · *why Zarr rather than whole NetCDF files*
- [pyParaOcean \(IIsc\): desktop ocean volumetrics we bring to the browser](#)

DATA SOURCES WE INGEST

- [INCOIS ERDDAP: the analysis field, 24 levels, 5 to 2000 m](#)
- [Argo GDAC \(Ifremer\): Indian Ocean profile files](#)
- [Copernicus Marine: GLOBAL_ANALYSISFORECAST_PHY_001_024 currents](#)
- [NOAA PMEL: RAMA moored buoy array, Indian Ocean](#)
- [NDMA SACHET: India's national CAP alert backbone](#)
- [INCOIS: Indian National Centre for Ocean Information Services](#)

OPEN STANDARDS IMPLEMENTED

- [CF Conventions for climate and forecast metadata](#)
- [OGC Web Map Service \(WMS\) 1.3.0](#)
- [OGC Web Coverage Service \(WCS\) 1.0.0](#)
- [OASIS Common Alerting Protocol \(CAP\) v1.2](#)


HOW THE RESEARCH CHANGED THE BUILD · not a bibliography, a set of decisions

Ryan 2015 → we verify in observation space and refuse to fill a level, rather than scoring on the model grid.

Yu 2025 → stop paying for a slice once it cannot change the picture: 10 of 24 blended at a time.

Wong 2020 → QC flags 1 and 2 only; 4,672 levels excluded and each one counted by reason.

Signell 2019 → Zarr chunks over HTTP range reads instead of whole NetCDF files.



WHERE WE SIT · GAP ANALYSIS

Capability	Desktop tools	Web viewers	Digital Ocean	Sagar Drishti
Browser-native, no install	✗	✓	✓	✓
True volumetric water column	✓	✗	~	✓
Model + in-situ in ONE 3D scene	✗	✗	~	✓
Model-vs-observation skill on screen	✗	✗	✗	✓
CAP hazard layer in ocean context	✗	✗	✗	✓
Runs fully offline / air-gapped	✓	✗	✗	✓

Desktop tools: ODV, IDV, pyParaOcean. Web viewers: Argovis, MyOcean. INCOIS's Digital Ocean portal does advertise 3D and 4D visualisation; what it does not do is fuse instruments into a true volumetric scene or publish model skill beside the field. Full landscape and citations in our prior-art review.

OUR OWN WORK, AND HOW IT CAN BE CHECKED

- 543 automated tests; 10 of them drive a real browser through this exact demo, so the claims on these slides cannot rot silently.
- Our density field reproduces the published TEOS-10 check values to six decimal places.
- Every panel on screen prints its own dataset name, timestamp and citation, so any figure here can be traced back to its source.
- Prior-art review of 15 papers and 8 existing tools is what produced the gap analysis above, before a line of code was written.
- Nothing is claimed that is not built: the deck names what is missing (glider and CTD files, an unrun Docker path) as plainly as what works.