

Proposal: Literature Review & Research Questions

BENNY ROCHWERG, University of Toronto, Canada

ACM Reference Format:

Benny Rochwerg. 2026. Proposal: Literature Review & Research Questions. 1, 1 (March 2026), 2 pages. <https://doi.org/10.1145/nnnnnnn.nnnnnnn>

1 Literature Review & Research Questions

Postoperative hypoxemia is a state that can arise after surgery where blood or tissue oxygen levels become deficient [2]. This state occurs regularly as a post-anesthetic care unit anesthetic complication and has a prevalence as high as 80% worldwide [2].

A previous study used logistic regressions to explore the relationship between early postoperative hypoxemia and variables such as duration of surgery, presence of heart disease, and anesthesia type [5]. Other variables, including body mass index (BMI), sex, and age, were also listed as independent variables in this study [5]. In addition, another study examined the relationship between early postoperative hypoxemia and variables including the presence of chronic disease, BMI, and smoking status using logistic regressions [6]. Furthermore, a different study employed multivariate logistic regression and found that among acute type A aortic dissection patients, BMI and gender were postoperative severe hypoxemia predictors [4]. Additional variables such as operation duration and white blood cell counts before surgery were also included in the analysis [4]. Finally, another study used logistic regression to predict postoperative hypoxemia for patients experiencing acute aortic dissection operation based on variables such as age, BMI, white blood cell count, pH, and hematocrit [3]. Other risk factors such as C-reactive protein (CRP) and lactate were also presented [3].

Although the impacts of several variables on postoperative hypoxemia were examined in these studies, they did not emphasize the use of comprehensive laboratory test results when modeling [3–6].

Consequently, we plan to answer the following research questions in our analysis:

- (1) By using patients' demographic variables (BMI, sex, and age) and latest available preoperative laboratory test results as predictors to model the probability of postoperative hypoxemia with logistic regression following a collinearity check, which variables are significant predictors of postoperative hypoxemia at the 0.05 level of significance?
- (2) By using only the identified significant predictors as the independent variables in another logistic regression model for the probability of postoperative hypoxemia, how accurately can the probability of postoperative hypoxemia be predicted?

Acknowledgments

The ACM Conference Proceedings Primary Article Template [1] was used to prepare this document.

Author's Contact Information: Benny Rochwerg, benny.rochwerg@mail.utoronto.ca, University of Toronto, Toronto, Ontario, Canada.

Permission to make digital or hard copies of all or part of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. Copyrights for components of this work owned by others than the author(s) must be honored. Abstracting with credit is permitted. To copy otherwise, or republish, or post on servers or to redistribute to lists, requires prior specific permission and/or a fee. Request permissions from permissions@acm.org.

© 2026 Copyright held by the owner/author(s). Publication rights licensed to ACM.

ACM XXXX-XXXX/2026/3-ART

<https://doi.org/10.1145/nnnnnnn.nnnnnnn>

A Generative AI

No generative AI tools were used.

References

- [1] Association for Computing Machinery (ACM). 2025. ACM Conference Proceedings Primary Article Template. Retrieved February 25, 2026 from <https://www.overleaf.com/latex/templates/acm-conference-proceedings-primary-article-template/wbvngbjbwpc>
- [2] Yosef Belay Bizuneh, Muluallem Endeshaw Zeleke, Ashebir Mamay Gebru, Destaw Fetene Teshome, Fantahun Ayenew Mekonnen, Dessie Abebaw Angaw, and Amare Belete Getahun. 2025. Global prevalence of postoperative hypoxemia among adult and pediatric surgical patients: a systematic review and meta-analysis. *BMC Anesthesiology* 25, Article 284 (June 2025), 10 pages. doi:10.1186/s12871-025-03146-3
- [3] Huiqing Ge, Ye Jiang, Qijun Jin, Linjun Wan, Ximing Qian, and Zhongheng Zhang. 2018. Nomogram for the prediction of postoperative hypoxemia in patients with acute aortic dissection. *BMC Anesthesiology* 18, Article 146 (Oct. 2018), 9 pages. doi:doi.org/10.1186/s12871-018-0612-7
- [4] Ming Gong, Zining Wu, Shijun Xu, Lei Li, Xiaolong Wang, Xinliang Guan, and Hongjia Zhang. 2019. Increased risk for the development of postoperative severe hypoxemia in obese women with acute type a aortic dissection. *Journal of Cardiothoracic Surgery* 14, Article 81 (April 2019), 9 pages. doi:10.1186/s13019-019-0888-9
- [5] Debas Yaregal Melesse, Zewditu Abdissa Denu, Habtamu Getinet Kassahun, and Abatneh Feleke Agegnehu. 2020. The incidence of early post-operative hypoxemia and its contributing factors among patients underwent operation under anesthesia at University of Gondar comprehensive and specialized referral hospital, Gondar, North West Ethiopia, 2018. A prospective observational study. *International Journal of Surgery Open* 22 (2020), 38–46. doi:10.1016/j.ijso.2019.11.011
- [6] Moges Gelaw Taye, Amelework Molla, Diriba Teshome, Metages Hunie, Simegne Kibret, Yewlsew Fentie, Netsanet Temesgen, Melaku Tadege Engidaw, and Efreem Fenta. 2021. Predictors of hypoxemia after general anesthesia in the early postoperative period in a hospital in Ethiopia: an observational study. *Multidisciplinary Respiratory Medicine* 16 (Dec. 2021), 6 pages. doi:10.4081/mrm.2021.782

Updated 25 February 2026