

Investigating Postoperative Hypoxemia Using the Medical Informatics Operating Room Vitals and Events Repository (MOVER)

Authors¹

¹Dalla Lana School of Public Health, Biostatistics Division, University of Toronto, Toronto, Canada

Abstract

Introduction & Literature Review

- Postoperative hypoxemia is a frequent complication from anesthetic¹
- The MOVER EPIC dataset includes records from 64,354 surgeries between 2017 and 2022² and can be used to study this issue
- Prior research has assessed how postoperative hypoxemia is affected by different variables, but these models only contained up to four laboratory test results^{3,4,5}
- To fill this gap characterized by few laboratory test results, this work modelled postoperative hypoxemia probability using logistic regression with up to ten preoperative laboratory test results and three demographic variables as covariates
- The objective was to detect significant predictors of postoperative hypoxemia and determine the accuracy of predicting its probability with just these variables

Methodology (The Pipeline)




Figure 1. Flow chart of data wrangling architecture

Key Findings

Variable	Coefficient Estimate	P-value
Intercept	-6.576	0.237
Age	0.004	0.660
Sex: Female	-1.426	0.001
Body mass index	0.051	0.040
Carbon dioxide	0.015	0.716
log(Glucose)	0.786	0.114
Hematocrit	0.094	0.557
Hemoglobin	-0.521	0.264
log(Leukocytes)	0.257	0.485
Potassium	0.117	0.717
Sodium	0.014	0.650

Table 1. Full logistic regression model results for postoperative hypoxemia

- The reduced model, which included only sex and body mass index, achieved a lower test set misclassification rate (28.0%) than the full model (29.3%).

- In the full model, sex and body mass index were the only predictors significant at the 0.05 level (Table 1).
- Female sex is associated with lower odds of postoperative hypoxemia and higher body mass index associated with higher odds.

Conclusion & Impact

Key Takeaway: Patients who are **male** or **have high BMI** may benefit from closer postoperative monitoring for timely detection and intervention of hypoxemia

Key Risk Factors of Postoperative hypoxemia

- Higher BMI** elevates odds, aligning with established literature⁴
- Higher odds among **males**, contradicting previous findings⁴

Model Performance and Limitation

- The reduced model outperforms the full model, suggesting that additional variables introduced noise rather than clinical signal
- Only 98 out of 24,167 patients (0.4%) had hypoxemia. This extreme class imbalance limited model performance despite undersampling control cases to balance the data

Future Directions

- Employing systematic oversampling or re-weighting techniques to improve minority class classification
- Investigating alternative biomarkers and the interplay between hypoxemia and concurrent postoperative complications to enhance predictive accuracy

References

- ¹ Yosef Belay Bizuneh, Mulualem Endeshaw Zeleke, Ashebir Mamay Gebru, Destaw Fetene Teshome, Fantahun Ayenew Mekonnen, Dessie Abebaw Angaw, and Amare Belete Getahun. 2025. Global prevalence of postoperative hypoxemia among adult and pediatric surgical patients: a systematic review and meta-analysis. *BMC Anesthesiology* 25, Article 284 (June 2025), 10 pages. doi:10.1186/s12871-025-03146-3
- ² Muntaha Samad, Mirana Angel, Joseph Rinehart, Yuzo Kanomata, Pierre Baldi, and Maxime Cannesson. 2023. Medical Informatics Operating Room Vitals and Events Repository (MOVER): a public-access operating room database. *JAMIA Open* 6, 4, Article ooad084 (Oct. 2023), 8 pages. doi:10.1093/jamiaopen/ooad084
- ³ Huiqing Ge, Ye Jiang, Qijun Jin, Linjun Wan, Ximing Qian, and Zhongheng Zhang. 2018. Nomogram for the prediction of postoperative hypoxemia in patients with acute aortic dissection. *BMC Anesthesiology* 18, Article 146 (Oct. 2018), 9 pages. doi:10.1186/s12871-018-0612-7
- ⁴ Ming Gong, Zining Wu, Shijun Xu, Lei Li, Xiaolong Wang, Xinliang Guan, and Hongjia Zhang. 2019. Increased risk for the development of postoperative severe hypoxemia in obese women with acute type a aortic dissection. *Journal of Cardiothoracic Surgery* 14, Article 81 (April 2019), 9 pages. doi:10.1186/s13019-019-0888-9
- ⁵ Debas Yaregal Melesse, Zewditu Abdissa Denu, Habtamu Getinet Kassahun, and Abatneh Feleke Agegnehu. 2020. The incidence of early post-operative hypoxemia and its contributing factors among patients underwent operation under anesthesia at University of Gondar comprehensive and specialized referral hospital, Gondar, North West Ethiopia, 2018. A prospective observational study. *International Journal of Surgery Open* 22 (2020), 38–46. doi:10.1016/j.ijso.2019.11.011