ICU MEDICATIONS

MICU(1) 蕭旋玲藥師製表20240512更新

Drug	Class	Dose Range	Comment	Concentration	Speed Algorithm
Vasopressors, Inotropics, and Chronotropics					
Dopamine 200mg/5ml/amp	D	1 - 3 μg/kg/min	2 amp in 250cc NS/D5W run ***10*** cc/hr (BW= 60kg, 4.3 μg/kg/min)	~ 2.96 mg/ml	***X*** (流速)= **0.17** x BW (4.3 μg/kg/min)
	β_1 , D	3 - 10 μg/kg/min			
	α_1 ,β_1	10 - 20 μg/kg/min			
Dobutamine 250mg/20ml/vial	$\beta_1 > \beta_2 >> \alpha$	2 - 20 μg/kg/min	2 vial in 250cc NS/D5W run ***10*** cc/hr (BW= 60kg, 4.8 μg/kg/min)	~ 1.72 mg/ml	***X*** (流速)= **0.17** x BW (4.8 μg/kg/min)
Norepinephrine 4mg/4ml/amp	$\alpha_1 >> \beta_1$	1 - 40 μg/min, 0.05 - 2 μg/kg/min	2 amp in D5W(建議) 250cc run 10 cc/hr (5.2 μg/min)	~ 0.06 mg/ml	***X*** (流速)= **10** (5.2 μg/min)
Vasopressin 20U/ml/amp	V1	0.01- 0.03 U/min, Max: 0.03 U/min, (應於NE≧ 30μg/min才使用)	1vial in NS 100ml ivf **3** ml/hr (0.01 U/min) +/- 1.5ml/hr at 10 - 15 min interval, max: 10 ml/hr	~ 0.2 U/ml	***X*** (流速)= **3** (0.01 U/min)
Isoproterenol 0.2mg/ml/amp	α_1 , β_2	1- 10 μg/min	5 amp in NS/D5W 500cc run 30 cc/hr (1 μg/min)	~ 0.004 mg/ml	***X*** (流速)= **30** (1 μg/min)
Epinephrine 1mg/ml/amp	$\alpha_{1,2}$, $\beta_{1,2}$	2 - 20 μg/min	anaphylatic shock: 2 - 10 mcg/min (after 1mg/1ml IM) 5 amp in NS/D5W 500cc run 20 cc/hr (3.3 μg/min)	~ 0.04 mg/ml	***X*** (流速)= **20** (3.3 μg/min)
Milrinone 10mg/10ml/vial	PDE	50 μg/kg over 10 min then 0.375 - 0.75 μg/kg/min			
Vasodilators					
Nitroglycerin 0.2mg/ml, 250ml/bot, or 5mg/10ml/amp	NO	10 - 400 μg/min	IVF: (0.5mg/ml原液) 2 cc/hr (16.7μg/min) then q 10 min ↑1-2 cc/hr till symptom relief or hypotension	~ 0.5 mg/ml	***X*** (流速)= **2** (16.7 μg/min)
Nicardipine HCl 10mg/10ml/amp	CCB	5 - 15 mg/hr	5 amp in NS/D5W 250ml ivf 15 cc/hr (2.5 mg/hr) initial: 5 mg/hr ivf; titrate 2.5 mg/hr every 15 min until desired blood pressure control is achieved, max dose: 15 mg/hr	~ 0.29 mg/ml	***X*** (流速)= **15** (2.5 mg/hr)
Labetalol 25mg/5ml/amp	α_1, β_1, β_2 blocker	IV: 20mg over 2 min, then 20-80 mg q10 min or IVF: 10-120 mg/hr	8 amp (200mg) diluted with D5W to 250ml ivf 10 cc/hr (7 mg/hr)		***X*** (流速)= **10** (7 mg/hr)
Hydralazine 20mg/ml/amp	vasodilator	5-20 mg q 4 - 6 h	IV > 1 min, dilute with NS (建議)		

ICU MEDICATIONS

MICU(1) 蕭旋玲藥師製表20240512更新

Drug	Class	Dose	Comment	Concentration	Speed Algorithm
Anti-arrhythmics					
Amiodarone 150mg/3ml/amp	Class Ⅲ	LD: 150mg over 10 min, then 1mg/min* 6hrs, then 0.5mg/min* 18 hrs	LD: 1 amp in D5W(建議) 30ml ivf > 10min then 6 amp in D5W 500 cc 玻璃瓶or PP bag(建議) ivf 33 cc/hr * 6hrs then 17 cc/hr * 18hrs	~ 1.7 mg/ml	
Lidocaine 2%, 5ml/amp (100mg/pc)	Na channel (Class IB)	LD: 1-1.5mg/kg then MD: 1- 4mg/min	LD: 50 -75mg iv > 1min then MD: 2g in NS/D5W **400**cc run 15 cc/hr (1 mg/min)	~ 4 mg/ml	***X*** (流速)= **15** (1 mg/min)
Verapamil 5mg/2ml/amp	CCB	IV: 2.5 - 5mg, repeat 5 -10 mg in 15 -30 min prn then IVF: 5-20mg/hr	IV > 1 - 2 min		
Adenosine 6mg/2ml/vial	purinergic	6mg → 12mg → 12mg	IV > 1-2 sec, followed by 20ml NS, after 2 min, give the 2nd dose (12mg) if necessary, 12mg can repeat once		
Sedation, Paralysis, Pain-control and Miscellaneous					
Morphine 10mg/ml/amp	opioid	IV: 2 - 4 mg q1-2 hr IVF: 2 - 30 mg/hr	IV > 1 min, 2 amp in NS/D5W 250cc run 12 cc/hr (1 mg/hr) for tachypnea when weaning from MV		***X*** (流速)= **12** (1 mg/hr)
Fentanyl citrate 0.05mg/ml, **2**ml/amp (**0.1**mg/pc)	opioid	IV: 0.35 - 0.5 μg /kg q 0.5-1 hr IVF: **0.15 - 0.6** μg/kg/hr	2 amp in NS/D5W 250 ml ivf 12 ml/hr (BW= 60kg, 0.16 μg/kg/hr)	~ 0.002 mg/ml	***X*** (流速)= **0.2** x BW (0.16 μg/kg/hr)
Dexmedetomidine 100mcg/ml,2ml/vial	α2 agonist	LD: 0.5 μg/kg >10 mins (注意BP, HR) IVF: 0.2 - 0.7 μg/kg/hr	1 vial in NS 100ml ivf 6 ml/hr (BW= 60kg, 0.2 μg/kg/hr)	~ 2 μg/ml	***X*** (流速)= **0.1** x BW (0.2 μg/kg/hr)
Propofol 10mg/ml 200mg/20ml/amp,500mg/50ml/vial	anesthetic	IV: 0.03 -1 mg/kg > 5 mins IVF: 0.3 - 3 mg/kg/hr	LD: (原液) 3 - 6 cc then 3 - 6 cc/hr initially	~ 10 mg/ml	***X*** (流速)= **0.1** x BW (1 mg/kg/hr)
Midazolam 5mg/1ml/amp, 15mg/3ml/amp	BZD	LD: 0.01- 0.05 mg/kg > 3-5mins IVF: 0.02 - **0.1** mg/kg/hr (指引已下修上限)	IVF: 5 amp (15mg/amp) in 250cc NS/D5W run ***22*** cc/hr (BW= 60kg, **0.1** mg/kg/hr) Refractory seizure: **0.045 - 0.6** mg/kg/hr	~ 0.54 mg/ml	***X*** (流速)= **0.25** x BW (0.07 mg/kg/hr)
Diazepam 10mg/2ml/amp	BZD	LD: 5 - 10mg then MD: 0.03 - 0.1 mg/kg q 0.5 - 6 hr prn			
Lorazepam 2mg/ml/amp	BZD	LD: 0.02 - 0.04 mg/kg (≤ 2 mg) IVF: 0.01 - 0.1 mg/kg/hr (≤10 mg/hr)			
Cisatracurium 2mg/ml,5ml/amp	nACH	IV: 0.1 - 0.2 mg/kg IVF: 0.03 - 0.2 mg/kg/hr	IVF: 5 amp (10mg/amp) in 250cc NS/D5W run ***17*** cc/hr (BW= 60kg, 0.05 mg/kg/hr)	~ 0.33 mg/ml	***X*** (流速)= **0.28** x BW (0.05 mg/kg/hr)
Rocuronium 50mg/5mL/vial, 100mg/10ml/vial (Cisatracurium缺貨時, continuous ivf NMBA取代用藥)	nACH	IV: 0.5 - 1 mg/kg IVF: 0.24 - 0.96 mg/kg/hr	IVF: 5 amp (10mg/amp) in 250cc NS/D5W run ***15*** cc/hr (BW= 60kg, 0.23 mg/kg/hr)	~ 1.67 mg/ml	***X*** (流速)= **0.25** x BW (0.24 mg/kg/hr)
Aminophyllin 250mg/10ml/amp	PED	LD: 5.5mg/kg (> 30 mins) then IVF: 0.5mg/kg/hr	2 amp in NS/D5W 500cc run ***30*** cc/hr (BW= 60kg, 0.5 mg/kg/hr)	~ 1.85 mg/ml	***X*** (流速)= **0.5** x BW (0.5 mg/kg/hr)