

(a)
Healthcare utilisation
GP help-seeking
Annual healthcare cost
Number of illnesses
Psychological distress
Post traumatic stress disorder
Anxiety disorder
Major depressive disorder
Sleep disorder
Respiratory infections
Pneumonia
Upper respiratory tract infection
COVID-19 (hospitalized)
COVID-19 (susceptibility)
Loading on G (λ, SE)
Fixed covariance (r = 0)
Loading on FIs (λ, SE)
Structural path (β, SE)
Non-significant (P > .05)
0.813 (0.024)
0.739 (0.017)
0.691 (0.016)
0.899 (0.024)
0.844 (0.018)
0.887 (0.034)
0.580 (0.031)
0.638 (0.032)
0.315 (0.022)
0.143 (0.041)
0.279 (0.039)
G
General factor
(healthcare contact,
distress & illness)
β = 0.531 (0.038)
Influenza
r = 0.000 (fixed)
β = 0.431 (0.120)
+0.374 (0.109)
-0.051 (0.063)
+1.006 (0.181)
+0.786 (0.150)
FIs
Infection-specific
residual factor
(b)
Dual-pathway
Single-pathway
Opposite pathway
β (SE)*
Tobacco use disorder
+0.544 (0.099)*
Alcohol dependence
+0.388 (0.099)*
Opiate dependence
+0.252 (0.066)*
Attention-deficit/hyperactivity disorder
+0.128 (0.059)
Alzheimer's disease
+0.104 (0.064)
Tourette syndrome
-0.014 (0.085)
Suicide attempt
-0.020 (0.067)
Obsessive-compulsive disorder
-0.044 (0.099)
Bipolar disorder
-0.069 (0.048)
Schizophrenia
-0.073 (0.042)
Parkinson's disease
-0.079 (0.080)
Anorexia nervosa
-0.154 (0.060)*
Autism spectrum disorder
-0.252 (0.082)*
-0.5
0.0
0.5
Standardized coefficient on FIs (95% CI)