

PHYS11 CH:5.4

Static and Kinetic Friction

Mr. Gullo

November 2024

Table of Contents

Types of Friction

- **Friction:** Force that opposes motion
- Two main types:
 - **Static Friction:** Acts on objects at rest
 - **Kinetic Friction:** Acts on objects in motion
- Maximum static friction is usually greater than kinetic friction

CH5.4/Screenshot 2024-11-11 110912.png

Friction Formulas

Static Friction

$$f_s \leq \mu_s N$$

where:

- f_s is static friction force
- μ_s is coefficient of static friction
- N is normal force

Kinetic Friction

$$f_k = \mu_k N$$

where:

- f_k is kinetic friction force
- μ_k is coefficient of kinetic friction

Coefficients of Friction

System	Static Friction	Kinetic Friction
Rubber on dry concrete	1.0	0.7
Wood on wood	0.5	0.3
Steel on steel (dry)	0.6	0.3
Steel on steel (oiled)	0.05	0.03
Ice on ice	0.1	0.03

Table of Contents

Forces on an Inclined Plane

- Weight components on an incline:
 - Parallel to slope: $w_{\parallel} = mg \sin \theta$
 - Perpendicular to slope: $w_{\perp} = mg \cos \theta$
- Normal force (N) equals perpendicular component
- Friction acts parallel to surface, opposing motion

CH5.4/Screenshot 2024-11-11 110923.png

Problem Solving Steps

1. Draw a sketch of the problem
2. Identify known and unknown quantities
3. Draw free-body diagram with rotated coordinate system
4. Apply Newton's second law:
 - If no acceleration: $F_{net} = 0$
 - If accelerating: $F_{net} = ma$
5. Check answer for reasonableness

Table of Contents

Example: Skier on a Slope

Problem

A 62 kg skier slides down a snowy slope at 25° . Find μ_k if friction is 45.0 N.

CH5.4/Screenshot 2024-11-11 110959SansFBD.png

CH5.4/Screenshot 2024-11-11 110959.png

CH5.4/Screenshot 2024-11-11 111003.png

Solution

$$N = mg \cos \theta = (62 \text{ kg})(9.80 \text{ m/s}^2) \cos(25)$$

$$f_k = \mu_k N = 45.0 \text{ N}$$

$$\mu_k = \frac{f_k}{N} = \frac{45.0 \text{ N}}{551 \text{ N}} = 0.082$$

Example: Acceleration on an Incline

Problem - Part A

What is the skier's acceleration if friction is negligible?

Solution

$$a = g \sin \theta$$

$$a = (9.80 \text{ m/s}^2) \sin(25) = 4.14 \text{ m/s}^2$$

Example: Acceleration with Friction

Problem - Part B

What is the skier's acceleration with 45.0 N friction?

Solution

$$F_{net} = mg \sin \theta - f_k$$

$$a = g \sin \theta - \frac{f_k}{m}$$

$$a = 9.80 \sin(25) - \frac{45.0}{60.0} = 3.39 \text{ m/s}^2$$

Table of Contents

Review Questions

- ① What is friction?
 - An external force that opposes relative motion
- ② Compare static vs. kinetic friction:
 - Static: Acts on objects at rest
 - Kinetic: Acts on objects in motion
 - Static friction maximum $>$ Kinetic friction

- Static friction maximum $>$ Kinetic friction

CH5.4/Friction-Plot-980x724.png