

# PHYS12 CH18: The Invisible Forces Between Charges

## Coulomb's Law, Fields, Potential, and Capacitors

Mr. Gullo

December 2025

# Outline

# The Mystery of the Invisible

What if forces could act  
*without touching?*

# The Mystery of the Invisible

What if forces could act  
*without touching?*

From lightning bolts to the neurons in your brain...

# The Mystery of the Invisible

What if forces could act  
*without touching?*

From lightning bolts to the neurons in your brain...

Electric forces shape reality.

# Learning Objectives

By the end of this section, you will be able to:

- **18.2:** Describe Coulomb's law verbally and mathematically

# Learning Objectives

By the end of this section, you will be able to:

- **18.2:** Describe Coulomb's law verbally and mathematically
- **18.2:** Solve problems involving Coulomb's law

## 18.2 The Force Between Charges

### The Mental Model

Like gravity pulls masses together, electric force acts between charges. But there's a twist: charges can attract OR repel.



## 18.2 The Force Between Charges

### The Mental Model

Like gravity pulls masses together, electric force acts between charges. But there's a twist: charges can attract OR repel.

### Two types of charge:

- Positive (+) and Negative (-)

## 18.2 The Force Between Charges

### The Mental Model

Like gravity pulls masses together, electric force acts between charges. But there's a twist: charges can attract OR repel.

### Two types of charge:

- Positive (+) and Negative (-)
- Like charges repel

## 18.2 The Force Between Charges

### The Mental Model

Like gravity pulls masses together, electric force acts between charges. But there's a twist: charges can attract OR repel.

### Two types of charge:

- Positive (+) and Negative (-)
- Like charges repel
- Unlike charges attract

## 18.2 Coulomb's Discovery



## 18.2 Coulomb's Discovery



Charles-Augustin de Coulomb (1780s) used a torsion balance to measure forces between charged spheres.

## 18.2 The Source Code of Electric Force

### Universal Law: Coulomb's Law

$$F = \frac{kq_1q_2}{r^2}$$

Force between charges equals Coulomb's constant times the product of charges divided by distance squared.

## 18.2 The Source Code of Electric Force

### Universal Law: Coulomb's Law

$$F = \frac{kq_1q_2}{r^2}$$

Force between charges equals Coulomb's constant times the product of charges divided by distance squared.

Where:  $k = 8.99 \times 10^9 \text{ N}\cdot\text{m}^2/\text{C}^2$

## 18.2 Reading the Signs

### The Paradox

If both charges are positive OR both negative:  $F > 0$  (repulsive)

If charges have opposite signs:  $F < 0$  (attractive)



## 18.2 Reading the Signs


### The Paradox

If both charges are positive OR both negative:  $F > 0$  (repulsive)

If charges have opposite signs:  $F < 0$  (attractive)



(a)



(b)



(c)

## 18.2 Inverse-Square Law

**What happens when you change the distance?**

## 18.2 Inverse-Square Law

### What happens when you change the distance?

- Double the distance ( $r \rightarrow 2r$ ): Force becomes  $\frac{1}{4}$  as strong

## 18.2 Inverse-Square Law

### What happens when you change the distance?

- Double the distance ( $r \rightarrow 2r$ ): Force becomes  $\frac{1}{4}$  as strong
- Triple the distance ( $r \rightarrow 3r$ ): Force becomes  $\frac{1}{9}$  as strong

## 18.2 Inverse-Square Law

### What happens when you change the distance?

- Double the distance ( $r \rightarrow 2r$ ): Force becomes  $\frac{1}{4}$  as strong
- Triple the distance ( $r \rightarrow 3r$ ): Force becomes  $\frac{1}{9}$  as strong
- Halve the distance ( $r \rightarrow \frac{r}{2}$ ): Force becomes 4 times stronger

## 18.2 Inverse-Square Law

### What happens when you change the distance?

- Double the distance ( $r \rightarrow 2r$ ): Force becomes  $\frac{1}{4}$  as strong
- Triple the distance ( $r \rightarrow 3r$ ): Force becomes  $\frac{1}{9}$  as strong
- Halve the distance ( $r \rightarrow \frac{r}{2}$ ): Force becomes 4 times stronger

### The Mental Model

Bring charges twice as close: force quadruples. Move them twice as far: force drops to one-fourth.

## 18.2 Coulomb vs Gravity

### Similarities:

- Both are inverse-square laws

## 18.2 Coulomb vs Gravity

### Similarities:

- Both are inverse-square laws
- Both act at a distance



## 18.2 Coulomb vs Gravity

### Similarities:

- Both are inverse-square laws
- Both act at a distance

### Differences:

- $k$  is MUCH larger than  $G$  (electric forces are stronger)

## 18.2 Coulomb vs Gravity

### Similarities:

- Both are inverse-square laws
- Both act at a distance

### Differences:

- $k$  is MUCH larger than  $G$  (electric forces are stronger)
- Gravity only attracts; electric forces attract OR repel

## 18.2 Coulomb vs Gravity

### Similarities:

- Both are inverse-square laws
- Both act at a distance

### Differences:

- $k$  is MUCH larger than  $G$  (electric forces are stronger)
- Gravity only attracts; electric forces attract OR repel

### Civilian vs Reality

**Civilian:** "Gravity holds everything together."

**Physicist:** "Electric forces hold atoms and molecules together. Gravity holds planets and galaxies."

# Attempt: Decoding Electric Force

## The Challenge (3 min, silent)

Two charges  $q_1 = +3 \times 10^{-9} \text{ C}$  and  $q_2 = -4 \times 10^{-9} \text{ C}$  are separated by 3.0 cm.

### Given:

- $q_1 = +3 \times 10^{-9} \text{ C}$
- $q_2 = -4 \times 10^{-9} \text{ C}$
- $r = 3.0 \text{ cm} = 0.030 \text{ m}$

**Find:** Magnitude and direction of force

*Can you calculate the force? Work silently.*

# Compare: Force Calculation

## Turn and talk (2 min):

- 1 What did you substitute for  $q_1$  and  $q_2$ ?
- 2 Did you convert cm to m?
- 3 Is the force attractive or repulsive? How do you know?

# Compare: Force Calculation

## Turn and talk (2 min):

- 1 What did you substitute for  $q_1$  and  $q_2$ ?
- 2 Did you convert cm to m?
- 3 Is the force attractive or repulsive? How do you know?

**Name wheel:** One pair share your approach (not your answer).

# Reveal: The Electric Interaction

**Self-correct in a different color:**

**Step 1:**  $F = \frac{kq_1q_2}{r^2}$

# Reveal: The Electric Interaction

**Self-correct in a different color:**

**Step 1:**  $F = \frac{kq_1q_2}{r^2}$

**Step 2:**  $F = \frac{(8.99 \times 10^9)(3 \times 10^{-9})(-4 \times 10^{-9})}{(0.030)^2}$



# Reveal: The Electric Interaction

**Self-correct in a different color:**

**Step 1:**  $F = \frac{kq_1q_2}{r^2}$

**Step 2:**  $F = \frac{(8.99 \times 10^9)(3 \times 10^{-9})(-4 \times 10^{-9})}{(0.030)^2}$

**Step 3:**  $F = -1.2 \times 10^{-4} \text{ N}$

# Reveal: The Electric Interaction

**Self-correct in a different color:**

**Step 1:**  $F = \frac{kq_1q_2}{r^2}$

**Step 2:**  $F = \frac{(8.99 \times 10^9)(3 \times 10^{-9})(-4 \times 10^{-9})}{(0.030)^2}$

**Step 3:**  $F = -1.2 \times 10^{-4} \text{ N}$

$|F| = 1.2 \times 10^{-4} \text{ N, attractive}$

# Reveal: The Electric Interaction

**Self-correct in a different color:**

**Step 1:**  $F = \frac{kq_1q_2}{r^2}$

**Step 2:**  $F = \frac{(8.99 \times 10^9)(3 \times 10^{-9})(-4 \times 10^{-9})}{(0.030)^2}$

**Step 3:**  $F = -1.2 \times 10^{-4} \text{ N}$

$|F| = 1.2 \times 10^{-4} \text{ N, attractive}$

**Check:** Opposite charges attract. Negative force confirms this!

# Learning Objectives

By the end of this section, you will be able to:

- **18.3:** Calculate the strength of an electric field

# Learning Objectives

By the end of this section, you will be able to:

- **18.3:** Calculate the strength of an electric field
- **18.3:** Create and interpret drawings of electric fields

## 18.3 Force Fields in Physics

What if space itself could push on charges?

## 18.3 Force Fields in Physics

What if space itself could push on charges?

### The Mental Model

An electric field is an invisible map showing which way a positive charge would be pushed at every point in space.

## 18.3 Force Fields in Physics

What if space itself could push on charges?

### The Mental Model

An electric field is an invisible map showing which way a positive charge would be pushed at every point in space.

Not science fiction - this is how physicists think about forces at a distance.



## 18.3 The Source Code of Fields

### Universal Law: Electric Field

$$\vec{E} = \frac{\vec{F}}{q_{\text{test}}}$$

Electric field equals force per unit charge.

## 18.3 The Source Code of Fields

### Universal Law: Electric Field

$$\vec{E} = \frac{\vec{F}}{q_{\text{test}}}$$

Electric field equals force per unit charge.

For a point charge  $Q$ :

$$E = \frac{k|Q|}{r^2}$$

## 18.3 The Source Code of Fields

### Universal Law: Electric Field

$$\vec{E} = \frac{\vec{F}}{q_{\text{test}}}$$


Electric field equals force per unit charge.

For a point charge  $Q$ :


$$E = \frac{k|Q|}{r^2}$$

Units: N/C (newtons per coulomb)

## 18.3 Visualizing the Invisible



## 18.3 Visualizing the Invisible



### Field lines show:

- Direction of force on positive charge
- Strength (closer lines = stronger field)

## 18.3 Field Line Rules

- 1 Lines point **away** from positive charges

## 18.3 Field Line Rules

- 1 Lines point **away** from positive charges
- 2 Lines point **toward** negative charges

## 18.3 Field Line Rules

- 1 Lines point **away** from positive charges
- 2 Lines point **toward** negative charges
- 3 Lines NEVER cross each other



## 18.3 Field Line Rules

- 1 Lines point **away** from positive charges
- 2 Lines point **toward** negative charges
- 3 Lines NEVER cross each other
- 4 Denser lines = stronger field

## 18.3 Field Line Rules


- 1 Lines point **away** from positive charges
- 2 Lines point **toward** negative charges
- 3 Lines NEVER cross each other
- 4 Denser lines = stronger field

### The Paradox

**Misconception:** "Field lines are paths charges follow."


**Reality:** Field lines show force direction, but moving charges have inertia - they curve gradually.

## 18.3 Field Patterns




Positive and negative

## 18.3 Field Patterns



Positive and negative



Two negatives

Field lines connect opposite charges, repel from like charges.

# Attempt: Reading Field Maps

## The Challenge (2 min, silent)

Look at this field map. Three charges create these field lines.

### Questions:

- 1 Which charges are positive? Which are negative?
- 2 Which charge has the largest magnitude?
- 3 Where is the field strongest?

*Use field line density and direction to decode the charges.*

# Compare: Field Interpretation

## Turn and talk (2 min):

- 1 How did you identify positive vs negative charges?
- 2 How did you compare charge magnitudes?
- 3 Where is the field strongest, and how do you know?

# Compare: Field Interpretation

## Turn and talk (2 min):

- 1 How did you identify positive vs negative charges?
- 2 How did you compare charge magnitudes?
- 3 Where is the field strongest, and how do you know?

**Name wheel:** One pair share your reasoning.

# Reveal: Decoding the Field

**Self-correct in a different color:**

**Signs:**

- Lines OUT = positive charge
- Lines IN = negative charge



# Reveal: Decoding the Field

**Self-correct in a different color:**

**Signs:**

- Lines OUT = positive charge
- Lines IN = negative charge

**Magnitude:**

- More lines = larger charge
- Count field lines touching each charge

# Reveal: Decoding the Field

## Self-correct in a different color:

### Signs:

- Lines OUT = positive charge
- Lines IN = negative charge

### Magnitude:

- More lines = larger charge
- Count field lines touching each charge

### Field Strength:

- Closest lines = strongest field
- Usually near charges

# Learning Objectives

By the end of this section, you will be able to:

- **18.4:** Explain similarities and differences between electric and gravitational potential energy

# Learning Objectives

By the end of this section, you will be able to:

- **18.4:** Explain similarities and differences between electric and gravitational potential energy
- **18.4:** Calculate electric potential difference

## 18.4 The Universe's Pressure Gauge

### The Mental Model

Gravitational potential: height in a gravitational field

Electric potential: "height" in an electric field


## 18.4 The Universe's Pressure Gauge

### The Mental Model



Gravitational potential: height in a gravitational field

Electric potential: "height" in an electric field

Before  
interaction



After  
interaction




## 18.4 The Universe's Pressure Gauge

### The Mental Model

Gravitational potential: height in a gravitational field

Electric potential: "height" in an electric field



Both store energy that can be released to do work.

## 18.4 Potential Energy of Two Charges

### Universal Law: Electric Potential Energy

$$U_E = \frac{kq_1q_2}{r}$$

Energy stored in configuration of two charges.



## 18.4 Potential Energy of Two Charges

### Universal Law: Electric Potential Energy

$$U_E = \frac{kq_1q_2}{r}$$

Energy stored in configuration of two charges.

### Sign tells the story:

- $U_E > 0$ : like charges (they want to fly apart)

## 18.4 Potential Energy of Two Charges

### Universal Law: Electric Potential Energy

$$U_E = \frac{kq_1q_2}{r}$$

Energy stored in configuration of two charges.

### Sign tells the story:

- $U_E > 0$ : like charges (they want to fly apart)
- $U_E < 0$ : opposite charges (they want to come together)

## 18.4 Electric Potential (Voltage)

### Universal Law: Electric Potential

$$V = \frac{U_E}{q} = \frac{kq}{r}$$

Potential energy per unit charge. Units: volts (V)

## 18.4 Electric Potential (Voltage)

### Universal Law: Electric Potential

$$V = \frac{U_E}{q} = \frac{kq}{r}$$

Potential energy per unit charge. Units: volts (V)

### Civilian vs Reality

**Civilian:** "Voltage is electricity flowing."

**Physicist:** "Voltage is electric pressure - potential energy per charge."

## 18.4 Potential Difference

**What really matters: difference in potential**

## 18.4 Potential Difference

**What really matters: difference in potential**

In uniform field  $E$ :

$$\Delta V = -E(x_f - x_i)$$

## 18.4 Potential Difference

**What really matters: difference in potential**

In uniform field  $E$ :

$$\Delta V = -E(x_f - x_i)$$

**Rearranged:**

$$E = \frac{\Delta V}{d}$$

## 18.4 Potential Difference

### What really matters: difference in potential

In uniform field  $E$ :

$$\Delta V = -E(x_f - x_i)$$

Rearranged:

$$E = \frac{\Delta V}{d}$$

Electric field units: V/m (volts per meter)



## 18.4 The 9V Battery

### Real-World: Battery Voltage

A 9V battery creates 9V potential difference between terminals.

This means: moving 1 coulomb from - to + terminal requires 9 joules of work.

## 18.4 The 9V Battery

### Real-World: Battery Voltage

A 9V battery creates 9V potential difference between terminals.

This means: moving 1 coulomb from - to + terminal requires 9 joules of work.

Battery converts chemical energy to electric potential energy.

# Attempt: Calculating Voltage

## The Challenge (3 min, silent)

A point charge  $Q = +5 \times 10^{-9} \text{ C}$  creates an electric potential.

### Given:

- $Q = +5 \times 10^{-9} \text{ C}$
- $r = 0.10 \text{ m}$

**Find:** Electric potential at distance  $r = 0.10 \text{ m}$

*Can you calculate the voltage? Work silently.*

# Compare: Voltage Calculation

## Turn and talk (2 min):

- 1 What formula did you use?
- 2 What did you substitute for each variable?
- 3 What are the units of your answer?

# Compare: Voltage Calculation

## Turn and talk (2 min):

- 1 What formula did you use?
- 2 What did you substitute for each variable?
- 3 What are the units of your answer?

**Name wheel:** One pair share your approach.

# Reveal: The Electric Pressure

**Self-correct in a different color:**

**Formula:**  $V = \frac{kQ}{r}$

# Reveal: The Electric Pressure

**Self-correct in a different color:**

**Formula:**  $V = \frac{kQ}{r}$

**Substitute:**  $V = \frac{(8.99 \times 10^9)(5 \times 10^{-9})}{0.10}$

# Reveal: The Electric Pressure

**Self-correct in a different color:**

**Formula:**  $V = \frac{kQ}{r}$

**Substitute:**  $V = \frac{(8.99 \times 10^9)(5 \times 10^{-9})}{0.10}$

$V = 450 \text{ V}$



# Reveal: The Electric Pressure

**Self-correct in a different color:**

**Formula:**  $V = \frac{kQ}{r}$

**Substitute:**  $V = \frac{(8.99 \times 10^9)(5 \times 10^{-9})}{0.10}$

$$V = 450 \text{ V}$$

**Check:** 450 volts - much higher than a battery, but safe at this tiny charge!

# Learning Objectives

By the end of this section, you will be able to:

- **18.5:** Calculate energy stored in a capacitor and capacitance

# Learning Objectives

By the end of this section, you will be able to:



- **18.5:** Calculate energy stored in a capacitor and capacitance
- **18.5:** Explain properties of capacitors and dielectrics

## 18.5 Energy Storage Devices

What if you could bottle electric fields?



## 18.5 Energy Storage Devices

What if you could bottle electric fields?




## 18.5 Energy Storage Devices

What if you could bottle electric fields?




Capacitors store energy in electric fields between charged plates.

## 18.5 The Parallel-Plate Capacitor



# 18.5 The Parallel-Plate Capacitor



## Design:

- Two metal plates separated by small distance
- One plate charged +, other charged -
- Electric field between plates is uniform



## 18.5 The Source Code of Capacitance

### Universal Law: Capacitance

$$C = \frac{Q}{V}$$

Capacitance equals charge stored per volt applied. Units: farads (F)

## 18.5 The Source Code of Capacitance

### Universal Law: Capacitance

$$C = \frac{Q}{V}$$

Capacitance equals charge stored per volt applied. Units: farads (F)

For parallel plates:

$$C_0 = \epsilon_0 \frac{A}{d}$$

where  $\epsilon_0 = 8.85 \times 10^{-12}$  F/m

## 18.5 Energy Storage

### Universal Law: Energy in Capacitor

$$U_E = \frac{1}{2}CV^2$$

Energy stored equals half capacitance times voltage squared.

## 18.5 Energy Storage

### Universal Law: Energy in Capacitor

$$U_E = \frac{1}{2}CV^2$$

Energy stored equals half capacitance times voltage squared.

### The Mental Model

Like kinetic energy  $K = \frac{1}{2}mv^2$ , but for electric fields instead of motion.

# 18.5 Capacitance Factors

**What determines capacitance?**

# 18.5 Capacitance Factors

**What determines capacitance?**

$$C_0 = \epsilon_0 \frac{A}{d}$$

# 18.5 Capacitance Factors

**What determines capacitance?**

$$C_0 = \epsilon_0 \frac{A}{d}$$

- Increase plate area  $A$ : capacitance increases

# 18.5 Capacitance Factors

## What determines capacitance?

$$C_0 = \epsilon_0 \frac{A}{d}$$

- Increase plate area  $A$ : capacitance increases
- Increase separation  $d$ : capacitance decreases



# 18.5 Capacitance Factors

## What determines capacitance?

$$C_0 = \epsilon_0 \frac{A}{d}$$

- Increase plate area  $A$ : capacitance increases
- Increase separation  $d$ : capacitance decreases
- Geometry only - not charge or voltage!

# 18.5 Capacitance Factors

## What determines capacitance?

$$C_0 = \epsilon_0 \frac{A}{d}$$

- Increase plate area  $A$ : capacitance increases
- Increase separation  $d$ : capacitance decreases
- Geometry only - not charge or voltage!

## The Paradox

**Misconception:** "More charge means more capacitance."

**Reality:** Capacitance is constant for given geometry. More charge just means higher voltage.

# 18.5 Dielectrics

## Real-World Enhancement

Insert insulating material (dielectric) between plates:

- Capacitance increases
- Can store more energy
- Prevents electrical breakdown

# 18.5 Dielectrics

## Real-World Enhancement

Insert insulating material (dielectric) between plates:

- Capacitance increases
- Can store more energy
- Prevents electrical breakdown

## Common dielectrics:

- Paper, plastic, ceramic, air

# Attempt: Capacitor Design

## The Challenge (3 min, silent)

Design a parallel-plate capacitor with capacitance  $C = 1.0 \times 10^{-9} \text{ F}$ .

### Given:

- $C = 1.0 \times 10^{-9} \text{ F}$  (1.0 nF)
- Plate area  $A = 0.010 \text{ m}^2$
- $\epsilon_0 = 8.85 \times 10^{-12} \text{ F/m}$

**Find:** Required plate separation  $d$

*Can you find the spacing? Work silently.*

# Compare: Design Strategy

## Turn and talk (2 min):

- 1 What formula did you start with?
- 2 How did you rearrange to solve for  $d$ ?
- 3 What units did you get for  $d$ ?

# Compare: Design Strategy

## Turn and talk (2 min):

- 1 What formula did you start with?
- 2 How did you rearrange to solve for  $d$ ?
- 3 What units did you get for  $d$ ?

**Name wheel:** One pair share your approach.

# Reveal: The Design Solution

**Self-correct in a different color:**

**Formula:**  $C_0 = \varepsilon_0 \frac{A}{d}$



# Reveal: The Design Solution

**Self-correct in a different color:**

**Formula:**  $C_0 = \varepsilon_0 \frac{A}{d}$

**Rearrange:**  $d = \varepsilon_0 \frac{A}{C}$

# Reveal: The Design Solution

**Self-correct in a different color:**

**Formula:**  $C_0 = \epsilon_0 \frac{A}{d}$

**Rearrange:**  $d = \epsilon_0 \frac{A}{C}$

**Substitute:**  $d = (8.85 \times 10^{-12}) \frac{0.010}{1.0 \times 10^{-9}}$

# Reveal: The Design Solution

**Self-correct in a different color:**

**Formula:**  $C_0 = \epsilon_0 \frac{A}{d}$

**Rearrange:**  $d = \epsilon_0 \frac{A}{C}$

**Substitute:**  $d = (8.85 \times 10^{-12}) \frac{0.010}{1.0 \times 10^{-9}}$

$$d = 8.85 \times 10^{-5} \text{ m} = 0.089 \text{ mm}$$

# Reveal: The Design Solution

**Self-correct in a different color:**

**Formula:**  $C_0 = \epsilon_0 \frac{A}{d}$

**Rearrange:**  $d = \epsilon_0 \frac{A}{C}$

**Substitute:**  $d = (8.85 \times 10^{-12}) \frac{0.010}{1.0 \times 10^{-9}}$

$$d = 8.85 \times 10^{-5} \text{ m} = 0.089 \text{ mm}$$

**Check:** Less than a tenth of a millimeter - capacitors need very small spacing!

# The Four Revelations

## What You Now Know

- 1 Coulomb's Law:  $F = \frac{kq_1q_2}{r^2}$  - forces between charges

# The Four Revelations

## What You Now Know

- 1 Coulomb's Law:  $F = \frac{kq_1q_2}{r^2}$  - forces between charges
- 2 Electric Field:  $E = \frac{F}{q}$  - force per unit charge

# The Four Revelations

## What You Now Know

- 1 Coulomb's Law:  $F = \frac{kq_1q_2}{r^2}$  - forces between charges
- 2 Electric Field:  $E = \frac{F}{q}$  - force per unit charge
- 3 Electric Potential:  $V = \frac{kq}{r}$  - energy per unit charge

# The Four Revelations

## What You Now Know

- 1 Coulomb's Law:  $F = \frac{kq_1q_2}{r^2}$  - forces between charges
- 2 Electric Field:  $E = \frac{F}{q}$  - force per unit charge
- 3 Electric Potential:  $V = \frac{kq}{r}$  - energy per unit charge
- 4 Capacitance:  $C = \frac{Q}{V}$  - charge storage capacity



# Key Equations

$$F = \frac{kq_1q_2}{r^2} \quad (\text{Coulomb's law}) \quad (1)$$

$$E = \frac{k|Q|}{r^2} \quad (\text{Electric field from point charge}) \quad (2)$$

$$\vec{E} = \frac{\vec{F}}{q} \quad (\text{Field definition}) \quad (3)$$

$$U_E = \frac{kq_1q_2}{r} \quad (\text{Electric potential energy}) \quad (4)$$

$$V = \frac{kq}{r} \quad (\text{Electric potential from point charge}) \quad (5)$$

$$C = \frac{Q}{V} \quad (\text{Capacitance}) \quad (6)$$

$$C_0 = \epsilon_0 \frac{A}{d} \quad (\text{Parallel-plate capacitor}) \quad (7)$$

$$U_E = \frac{1}{2} CV^2 \quad (\text{Energy in capacitor}) \quad (8)$$

# Constants to Remember

| Constant                  | Value  |
|---------------------------|--|
| Coulomb's constant $k$    | $8.99 \times 10^9 \text{ N}\cdot\text{m}^2/\text{C}^2$ |
| Elementary charge $e$     | $1.602 \times 10^{-19} \text{ C}$                      |
| Permittivity $\epsilon_0$ | $8.85 \times 10^{-12} \text{ F/m}$                     |

These appear in every calculation!

# From Theory to Reality

## **You now understand:**

- Why static shocks happen (charge transfer)

# From Theory to Reality

## **You now understand:**

- Why static shocks happen (charge transfer)
- How touchscreens work (capacitance sensing)

# From Theory to Reality

## **You now understand:**

- Why static shocks happen (charge transfer)
- How touchscreens work (capacitance sensing)
- Why lightning forms (electric field breakdown)

# From Theory to Reality

## **You now understand:**

- Why static shocks happen (charge transfer)
- How touchscreens work (capacitance sensing)
- Why lightning forms (electric field breakdown)
- How defibrillators work (capacitor discharge)

# From Theory to Reality

## You now understand:

- Why static shocks happen (charge transfer)
- How touchscreens work (capacitance sensing)
- Why lightning forms (electric field breakdown)
- How defibrillators work (capacitor discharge)
- Why atoms bond (electric forces)

# From Theory to Reality

## You now understand:

- Why static shocks happen (charge transfer)
- How touchscreens work (capacitance sensing)
- Why lightning forms (electric field breakdown)
- How defibrillators work (capacitor discharge)
- Why atoms bond (electric forces)
- How computer memory works (capacitor charge storage)



Complete the assigned problems  
posted on the LMS

## **Temporary page!**

$\text{\LaTeX}$  was unable to guess the total number of pages correctly. There was some unprocessed data that should have been added to the document, so this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will disappear, because  $\text{\LaTeX}$  now knows how many pages to expect for the document.