

PHYS12 CH:5 Motion in Two Dimensions

Vectors and Projectiles

Mr. Gullo

December 2025

Outline

- 1 Introduction
- 2 Vector Addition and Subtraction: Graphical Methods
- 3 Vector Addition and Subtraction: Analytical Methods
- 4 Projectile Motion
- 5 Summary

The Mystery

How do you describe motion
when objects move in two directions at once?

The Mystery

How do you describe motion
when objects move in two directions at once?

A football spiraling downfield...

The Mystery

How do you describe motion
when objects move in two directions at once?

A football spiraling downfield...

A drone navigating through space...

The Mystery

How do you describe motion
when objects move in two directions at once?

A football spiraling downfield...

A drone navigating through space...

Motion happens in multiple dimensions simultaneously.

Learning Objectives

By the end of this section, you will be able to:

- **5.1:** Describe the graphical method of vector addition and subtraction

Learning Objectives

By the end of this section, you will be able to:

- **5.1:** Describe the graphical method of vector addition and subtraction
- **5.1:** Use the graphical method to solve physics problems

5.1 What Is a Vector?

The Dual Nature

A vector is a quantity with both magnitude AND direction.

5.1 What Is a Vector?

The Dual Nature

A vector is a quantity with both magnitude AND direction.

Vectors in physics:

- Displacement, velocity, acceleration

5.1 What Is a Vector?

The Dual Nature

A vector is a quantity with both magnitude AND direction.

Vectors in physics:

- Displacement, velocity, acceleration
- Force, momentum

5.1 What Is a Vector?

The Dual Nature

A vector is a quantity with both magnitude AND direction.

Vectors in physics:

- Displacement, velocity, acceleration
- Force, momentum
- Boldface notation: **F**, **v**, **a**

5.1 What Is a Vector?

The Dual Nature

A vector is a quantity with both magnitude AND direction.


Vectors in physics:

- Displacement, velocity, acceleration
- Force, momentum
- Boldface notation: **F**, **v**, **a**


The Mental Model

Velocity is like a speedometer with a compass attached. Speed tells how fast. Velocity adds where.

5.1 The Journey to the Destination




5.1 The Journey to the Destination



Path: 9 blocks east + 5 blocks north = 14 blocks walked

Displacement: 10.3 blocks at 29.1° north of east

5.1 The Journey to the Destination



Path: 9 blocks east + 5 blocks north = 14 blocks walked

Displacement: 10.3 blocks at 29.1° north of east

Civilian View vs. Reality

Civilian: "I walked 14 blocks."

Physicist: "Displacement was 10.3 blocks."

5.1 The Head-to-Tail Method

How to add vectors graphically:

- 1 Draw first vector to scale with protractor

5.1 The Head-to-Tail Method

How to add vectors graphically:

- 1 Draw first vector to scale with protractor
- 2 Place tail of second vector at head of first

5.1 The Head-to-Tail Method

How to add vectors graphically:

- 1 Draw first vector to scale with protractor
- 2 Place tail of second vector at head of first
- 3 Continue for all vectors

5.1 The Head-to-Tail Method

How to add vectors graphically:

- 1 Draw first vector to scale with protractor
- 2 Place tail of second vector at head of first
- 3 Continue for all vectors
- 4 Draw resultant from tail of first to head of last

5.1 The Head-to-Tail Method

How to add vectors graphically:

- 1 Draw first vector to scale with protractor
- 2 Place tail of second vector at head of first
- 3 Continue for all vectors
- 4 Draw resultant from tail of first to head of last
- 5 Measure magnitude with ruler, direction with protractor


5.1 The Head-to-Tail Method

How to add vectors graphically:


- 1 Draw first vector to scale with protractor
- 2 Place tail of second vector at head of first
- 3 Continue for all vectors
- 4 Draw resultant from tail of first to head of last
- 5 Measure magnitude with ruler, direction with protractor

Key insight: Order doesn't matter! $\mathbf{A} + \mathbf{B} = \mathbf{B} + \mathbf{A}$

5.1 Building the Resultant



5.1 Building the Resultant



The resultant vector connects start to finish.

5.1 Vector Subtraction


To subtract: Add the negative vector

$$\mathbf{A} - \mathbf{B} = \mathbf{A} + (-\mathbf{B})$$

5.1 Vector Subtraction

To subtract: Add the negative vector


$$\mathbf{A} - \mathbf{B} = \mathbf{A} + (-\mathbf{B})$$



5.1 Vector Subtraction


To subtract: Add the negative vector

$$\mathbf{A} - \mathbf{B} = \mathbf{A} + (-\mathbf{B})$$




The negative vector: Same magnitude, opposite direction

5.1 Forces on Ice




Free body diagram




(b)

5.1 Forces on Ice




Free body diagram




Two skaters push with 400 N each at right angles.

5.1 Forces on Ice




Free body diagram



Two skaters push with 400 N each at right angles.

Resultant force: $F_{tot} = \sqrt{(400)^2 + (400)^2} = 566 \text{ N}$

5.1 Forces on Ice



Two skaters push with 400 N each at right angles.

Resultant force: $F_{tot} = \sqrt{(400)^2 + (400)^2} = 566 \text{ N}$

Pythagorean theorem works when vectors are perpendicular!

Learning Objectives

By the end of this section, you will be able to:

- **5.2:** Define components of vectors

Learning Objectives

By the end of this section, you will be able to:


- **5.2:** Define components of vectors
- **5.2:** Describe the analytical method of vector addition

Learning Objectives

By the end of this section, you will be able to:

- **5.2:** Define components of vectors
- **5.2:** Describe the analytical method of vector addition
- **5.2:** Use the analytical method to solve problems

5.2 The Power of Trigonometry




$$\sin(\theta) = \frac{y}{b}$$

$$\cos(\theta) = \frac{x}{b}$$

$$\tan(\theta) = \frac{y}{x}$$

5.2 The Power of Trigonometry



$$\sin(\theta) = \frac{y}{b}$$


$$\cos(\theta) = \frac{x}{b}$$

$$\tan(\theta) = \frac{y}{x}$$

Refresher:

- $\sin \theta = \frac{\text{opposite}}{\text{hypotenuse}}$

5.2 The Power of Trigonometry



$$\sin(\theta) = \frac{y}{b}$$


$$\cos(\theta) = \frac{x}{b}$$

$$\tan(\theta) = \frac{y}{x}$$

Refresher:

- $\sin \theta = \frac{\text{opposite}}{\text{hypotenuse}}$
- $\cos \theta = \frac{\text{adjacent}}{\text{hypotenuse}}$

5.2 The Power of Trigonometry



$$\sin(\theta) = \frac{y}{b}$$

$$\cos(\theta) = \frac{x}{b}$$

$$\tan(\theta) = \frac{y}{x}$$

Refresher:

- $\sin \theta = \frac{\text{opposite}}{\text{hypotenuse}}$
- $\cos \theta = \frac{\text{adjacent}}{\text{hypotenuse}}$
- $\tan \theta = \frac{\text{opposite}}{\text{adjacent}}$

5.2 Breaking Vectors Into Components

The Source Code

$$A_x = A \cos \theta$$

$$A_y = A \sin \theta$$

Every 2D vector can be split into x and y components.


5.2 Breaking Vectors Into Components

The Source Code

$$A_x = A \cos \theta$$

$$A_y = A \sin \theta$$

Every 2D vector can be split into x and y components.



5.2 Example: City Walk Components

Displacement: $A = 10.3$ blocks at $\theta = 29.1^\circ$ north of east

5.2 Example: City Walk Components

Displacement: $A = 10.3$ blocks at $\theta = 29.1^\circ$ north of east

East component (x):

$$\begin{aligned} A_x &= A \cos \theta \\ &= (10.3)(\cos 29.1^\circ) = 9.0 \text{ blocks} \end{aligned}$$

5.2 Example: City Walk Components

Displacement: $A = 10.3$ blocks at $\theta = 29.1^\circ$ north of east

East component (x):

$$\begin{aligned} A_x &= A \cos \theta \\ &= (10.3)(\cos 29.1^\circ) = 9.0 \text{ blocks} \end{aligned}$$

North component (y):

$$\begin{aligned} A_y &= A \sin \theta \\ &= (10.3)(\sin 29.1^\circ) = 5.0 \text{ blocks} \end{aligned}$$

5.2 Example: City Walk Components

Displacement: $A = 10.3$ blocks at $\theta = 29.1^\circ$ north of east

East component (x):

$$\begin{aligned} A_x &= A \cos \theta \\ &= (10.3)(\cos 29.1^\circ) = 9.0 \text{ blocks} \end{aligned}$$

North component (y):

$$\begin{aligned} A_y &= A \sin \theta \\ &= (10.3)(\sin 29.1^\circ) = 5.0 \text{ blocks} \end{aligned}$$

Components match the original path: 9 east + 5 north!

5.2 Reverse: From Components to Vector

Reconstruction Formulas

$$A = \sqrt{A_x^2 + A_y^2}$$

$$\theta = \tan^{-1} \left(\frac{A_y}{A_x} \right)$$

Pythagorean theorem gives magnitude, inverse tangent gives direction.

5.2 Reverse: From Components to Vector

Reconstruction Formulas

$$A = \sqrt{A_x^2 + A_y^2}$$

$$\theta = \tan^{-1} \left(\frac{A_y}{A_x} \right)$$

Pythagorean theorem gives magnitude, inverse tangent gives direction.

The Paradox

Students confuse: $\mathbf{A}_x + \mathbf{A}_y = \mathbf{A}$ (vector addition)

with $A = \sqrt{A_x^2 + A_y^2}$ (magnitude calculation)

5.2 The Analytical Method

To add vectors **A and **B** analytically:**

- 1 Find components: A_x, A_y, B_x, B_y

5.2 The Analytical Method

To add vectors **A and **B** analytically:**

- 1 Find components: A_x, A_y, B_x, B_y
- 2 Add x-components: $R_x = A_x + B_x$

5.2 The Analytical Method

To add vectors **A and **B** analytically:**

- 1 Find components: A_x, A_y, B_x, B_y
- 2 Add x-components: $R_x = A_x + B_x$
- 3 Add y-components: $R_y = A_y + B_y$

5.2 The Analytical Method

To add vectors **A and **B** analytically:**

- 1 Find components: A_x, A_y, B_x, B_y
- 2 Add x-components: $R_x = A_x + B_x$
- 3 Add y-components: $R_y = A_y + B_y$
- 4 Find magnitude: $R = \sqrt{R_x^2 + R_y^2}$

5.2 The Analytical Method

To add vectors **A and **B** analytically:**

- 1 Find components: A_x, A_y, B_x, B_y
- 2 Add x-components: $R_x = A_x + B_x$
- 3 Add y-components: $R_y = A_y + B_y$
- 4 Find magnitude: $R = \sqrt{R_x^2 + R_y^2}$
- 5 Find direction: $\theta = \tan^{-1}(R_y/R_x)$

5.2 The Analytical Method

To add vectors **A and **B** analytically:**

- 1 Find components: A_x, A_y, B_x, B_y
- 2 Add x-components: $R_x = A_x + B_x$
- 3 Add y-components: $R_y = A_y + B_y$
- 4 Find magnitude: $R = \sqrt{R_x^2 + R_y^2}$
- 5 Find direction: $\theta = \tan^{-1}(R_y/R_x)$

More accurate than graphical method - not limited by drawing precision!

Attempt: Two-Leg Journey

The Challenge (3 min, silent)

A person walks 53.0 m at 20.0° north of east, then 34.0 m at 63.0° north of east.

Given:

- $A = 53.0 \text{ m}$, $\theta_A = 20.0^\circ$
- $B = 34.0 \text{ m}$, $\theta_B = 63.0^\circ$

Find: Total displacement magnitude and direction

Can you decode this journey? Work silently.

Compare: Vector Addition Strategy

Turn and talk (2 min):

- 1 What's the first step - find components or add vectors?
- 2 How many components do you need to calculate?
- 3 Do you add x and y together, or keep them separate?

Compare: Vector Addition Strategy

Turn and talk (2 min):

- 1 What's the first step - find components or add vectors?
- 2 How many components do you need to calculate?
- 3 Do you add x and y together, or keep them separate?

Name wheel: One pair share your approach (not your answer).

Reveal: The Navigation Solution

Self-correct in a different color:

Step 1: Find components of **A**

$$A_x = (53.0)(\cos 20.0^\circ) = 49.8 \text{ m}$$

$$A_y = (53.0)(\sin 20.0^\circ) = 18.1 \text{ m}$$

Reveal: The Navigation Solution

Self-correct in a different color:

Step 1: Find components of **A**

$$A_x = (53.0)(\cos 20.0^\circ) = 49.8 \text{ m}$$

$$A_y = (53.0)(\sin 20.0^\circ) = 18.1 \text{ m}$$

Step 2: Find components of **B**

$$B_x = (34.0)(\cos 63.0^\circ) = 15.4 \text{ m}$$

$$B_y = (34.0)(\sin 63.0^\circ) = 30.3 \text{ m}$$

Reveal: Combining Components

Step 3: Add components

$$R_x = A_x + B_x = 49.8 + 15.4 = 65.2 \text{ m}$$

$$R_y = A_y + B_y = 18.1 + 30.3 = 48.4 \text{ m}$$

Reveal: Combining Components

Step 3: Add components

$$R_x = A_x + B_x = 49.8 + 15.4 = 65.2 \text{ m}$$

$$R_y = A_y + B_y = 18.1 + 30.3 = 48.4 \text{ m}$$

Step 4: Find magnitude

$$R = \sqrt{(65.2)^2 + (48.4)^2} = \boxed{81.2 \text{ m}}$$

Reveal: Combining Components

Step 3: Add components

$$R_x = A_x + B_x = 49.8 + 15.4 = 65.2 \text{ m}$$

$$R_y = A_y + B_y = 18.1 + 30.3 = 48.4 \text{ m}$$

Step 4: Find magnitude

$$R = \sqrt{(65.2)^2 + (48.4)^2} = \boxed{81.2 \text{ m}}$$

Step 5: Find direction

$$\theta = \tan^{-1}(48.4/65.2) = \boxed{36.6^\circ \text{ north of east}}$$

Learning Objectives

By the end of this section, you will be able to:

- **5.3:** Describe the properties of projectile motion

Learning Objectives

By the end of this section, you will be able to:

- **5.3:** Describe the properties of projectile motion
- **5.3:** Apply kinematic equations and vectors to solve projectile problems

5.3 The Great Separation

Nature's Rule for Projectiles

Horizontal and vertical motions
are independent


They don't influence each other.

5.3 The Great Separation

Nature's Rule for Projectiles

Horizontal and vertical motions
are independent

They don't influence each other.



5.3 What Is a Projectile?

Definition: An object launched into the air that moves under gravity alone

5.3 What Is a Projectile?

Definition: An object launched into the air that moves under gravity alone

Key terms:

- **Projectile:** The moving object

5.3 What Is a Projectile?

Definition: An object launched into the air that moves under gravity alone

Key terms:

- **Projectile:** The moving object
- **Trajectory:** Its curved path

5.3 What Is a Projectile?

Definition: An object launched into the air that moves under gravity alone

Key terms:

- **Projectile:** The moving object
- **Trajectory:** Its curved path
- **Range:** Horizontal distance traveled

5.3 What Is a Projectile?

Definition: An object launched into the air that moves under gravity alone

Key terms:

- **Projectile:** The moving object
- **Trajectory:** Its curved path
- **Range:** Horizontal distance traveled
- **Maximum height:** Peak altitude

5.3 What Is a Projectile?

Definition: An object launched into the air that moves under gravity alone

Key terms:

- **Projectile:** The moving object
- **Trajectory:** Its curved path
- **Range:** Horizontal distance traveled
- **Maximum height:** Peak altitude

We ignore air resistance in introductory physics.

5.3 The Independence Principle

Horizontal motion:

- No acceleration: $a_x = 0$

5.3 The Independence Principle

Horizontal motion:

- No acceleration: $a_x = 0$
- Constant velocity: $v_x = v_{0x}$

5.3 The Independence Principle

Horizontal motion:

- No acceleration: $a_x = 0$
- Constant velocity: $v_x = v_{0x}$
- Position: $x = x_0 + v_x t$

5.3 The Independence Principle

Horizontal motion:

- No acceleration: $a_x = 0$
- Constant velocity: $v_x = v_{0x}$
- Position: $x = x_0 + v_x t$

Vertical motion:

- Constant acceleration: $a_y = -g = -9.8 \text{ m/s}^2$

5.3 The Independence Principle

Horizontal motion:

- No acceleration: $a_x = 0$
- Constant velocity: $v_x = v_{0x}$
- Position: $x = x_0 + v_x t$

Vertical motion:

- Constant acceleration: $a_y = -g = -9.8 \text{ m/s}^2$
- Velocity: $v_y = v_{0y} - gt$

5.3 The Independence Principle

Horizontal motion:

- No acceleration: $a_x = 0$
- Constant velocity: $v_x = v_{0x}$
- Position: $x = x_0 + v_x t$

Vertical motion:

- Constant acceleration: $a_y = -g = -9.8 \text{ m/s}^2$
- Velocity: $v_y = v_{0y} - gt$
- Position: $y = y_0 + v_{0y} t - \frac{1}{2}gt^2$

5.3 The Independence Principle

Horizontal motion:


- No acceleration: $a_x = 0$
- Constant velocity: $v_x = v_{0x}$
- Position: $x = x_0 + v_x t$

Vertical motion:


- Constant acceleration: $a_y = -g = -9.8 \text{ m/s}^2$
- Velocity: $v_y = v_{0y} - gt$
- Position: $y = y_0 + v_{0y}t - \frac{1}{2}gt^2$

Time t is the only variable connecting the two motions!

5.3 The Trajectory



5.3 The Trajectory



Ball kicked at angle θ follows parabolic path.

5.3 The Four-Step Method

To solve projectile problems:

- 1 Separate into x and y components

5.3 The Four-Step Method

To solve projectile problems:

- 1 Separate into x and y components
- 2 Treat as two independent 1D motions

5.3 The Four-Step Method

To solve projectile problems:

- 1 Separate into x and y components
- 2 Treat as two independent 1D motions
- 3 Solve for unknowns (use t to connect them)

5.3 The Four-Step Method

To solve projectile problems:

- 1 Separate into x and y components
- 2 Treat as two independent 1D motions
- 3 Solve for unknowns (use t to connect them)
- 4 Recombine to find total displacement and velocity

5.3 The Four-Step Method

To solve projectile problems:

- 1 Separate into x and y components
- 2 Treat as two independent 1D motions
- 3 Solve for unknowns (use t to connect them)
- 4 Recombine to find total displacement and velocity

Recombination formulas:

$$d = \sqrt{x^2 + y^2}, \quad \theta = \tan^{-1}(y/x)$$

$$v = \sqrt{v_x^2 + v_y^2}, \quad \theta_v = \tan^{-1}(v_y/v_x)$$

5.3 Maximum Height Formula

The Peak

$$h = \frac{v_{0y}^2}{2g}$$

Maximum height depends only on initial vertical velocity.

5.3 Maximum Height Formula

The Peak

$$h = \frac{v_{0y}^2}{2g}$$

Maximum height depends only on initial vertical velocity.

Key insight: At maximum height, $v_y = 0$

5.3 Maximum Height Formula

The Peak

$$h = \frac{v_{0y}^2}{2g}$$

Maximum height depends only on initial vertical velocity.

Key insight: At maximum height, $v_y = 0$

What Your Brain Gets Wrong

Intuition: "Faster launch means higher peak."

Reality: "Only the *vertical* component matters."

Attempt: Fireworks Launch

The Challenge (3 min, silent)

A fireworks shell is launched at 70.0 m/s at 75° above horizontal.

Given:

- $v_0 = 70.0 \text{ m/s}$
- $\theta = 75^\circ$

Find: Maximum height

Can you predict where it explodes? Work silently.

Compare: Projectile Strategy

Turn and talk (2 min):

- 1 What's the first step - find components or use a formula?
- 2 Which velocity matters - total or vertical?
- 3 Which kinematic equation did you choose?

Compare: Projectile Strategy

Turn and talk (2 min):

- 1 What's the first step - find components or use a formula?
- 2 Which velocity matters - total or vertical?
- 3 Which kinematic equation did you choose?

Name wheel: One pair share your approach (not your answer).

Reveal: Reaching the Sky

Self-correct in a different color:

Step 1: Find vertical velocity component

$$\begin{aligned}v_{0y} &= v_0 \sin \theta \\&= (70.0)(\sin 75^\circ) = 67.6 \text{ m/s}\end{aligned}$$

Reveal: Reaching the Sky

Self-correct in a different color:

Step 1: Find vertical velocity component

$$\begin{aligned}v_{0y} &= v_0 \sin \theta \\&= (70.0)(\sin 75^\circ) = 67.6 \text{ m/s}\end{aligned}$$

Step 2: Use maximum height formula

$$\begin{aligned}h &= \frac{v_{0y}^2}{2g} \\&= \frac{(67.6)^2}{2(9.8)} = \boxed{233 \text{ m}}\end{aligned}$$

Reveal: Reaching the Sky

Self-correct in a different color:

Step 1: Find vertical velocity component

$$\begin{aligned}v_{0y} &= v_0 \sin \theta \\&= (70.0)(\sin 75^\circ) = 67.6 \text{ m/s}\end{aligned}$$

Step 2: Use maximum height formula

$$\begin{aligned}h &= \frac{v_{0y}^2}{2g} \\&= \frac{(67.6)^2}{2(9.8)} = \boxed{233 \text{ m}}\end{aligned}$$

Check: About 765 feet - reasonable for large fireworks!

5.3 The Range Equation

The Distance Formula

$$R = \frac{v_0^2 \sin 2\theta}{g}$$

Range depends on initial speed and launch angle.

5.3 The Range Equation

The Distance Formula

$$R = \frac{v_0^2 \sin 2\theta}{g}$$

Range depends on initial speed and launch angle.

Key insights:

- Maximum range at $\theta = 45^\circ$

5.3 The Range Equation

The Distance Formula

$$R = \frac{v_0^2 \sin 2\theta}{g}$$

Range depends on initial speed and launch angle.

Key insights:

- Maximum range at $\theta = 45^\circ$
- Same range for complementary angles: 30° and 60°

5.3 The Range Equation

The Distance Formula

$$R = \frac{v_0^2 \sin 2\theta}{g}$$

Range depends on initial speed and launch angle.

Key insights:

- Maximum range at $\theta = 45^\circ$
- Same range for complementary angles: 30° and 60°
- Only valid when initial and final heights are equal

Attempt: Volcanic Rock

The Challenge (3 min, silent)

A rock is ejected from a volcano at 25.0 m/s at 35° above horizontal. It lands 20.0 m below its starting point.

Given:

- $v_0 = 25.0 \text{ m/s}$, $\theta = 35^\circ$
- $y = -20.0 \text{ m}$

Find: Time of flight

Can you predict the flight time? Work silently.

Compare: Quadratic Strategy

Turn and talk (2 min):

- 1 Which kinematic equation has both y and t ?
- 2 What values do you substitute for y_0 and y ?
- 3 How do you solve a quadratic equation?

Compare: Quadratic Strategy

Turn and talk (2 min):

- 1 Which kinematic equation has both y and t ?
- 2 What values do you substitute for y_0 and y ?
- 3 How do you solve a quadratic equation?

Name wheel: One pair share your approach (not your answer).

Reveal: Time of Flight

Self-correct in a different color:

Step 1: Find $v_{0y} = (25.0)(\sin 35^\circ) = 14.3 \text{ m/s}$

Reveal: Time of Flight

Self-correct in a different color:

Step 1: Find $v_{0y} = (25.0)(\sin 35^\circ) = 14.3 \text{ m/s}$

Step 2: Use kinematic equation with $y_0 = 0$, $y = -20.0 \text{ m}$

$$-20.0 = (14.3)t - (4.90)t^2$$

Reveal: Time of Flight

Self-correct in a different color:

Step 1: Find $v_{0y} = (25.0)(\sin 35^\circ) = 14.3 \text{ m/s}$

Step 2: Use kinematic equation with $y_0 = 0$, $y = -20.0 \text{ m}$

$$-20.0 = (14.3)t - (4.90)t^2$$

Step 3: Rearrange to standard form

$$(4.90)t^2 - (14.3)t - (20.0) = 0$$

Reveal: Time of Flight

Self-correct in a different color:

Step 1: Find $v_{0y} = (25.0)(\sin 35^\circ) = 14.3 \text{ m/s}$

Step 2: Use kinematic equation with $y_0 = 0$, $y = -20.0 \text{ m}$

$$-20.0 = (14.3)t - (4.90)t^2$$

Step 3: Rearrange to standard form

$$(4.90)t^2 - (14.3)t - (20.0) = 0$$

Step 4: Quadratic formula

$$t = \frac{14.3 \pm \sqrt{(14.3)^2 - 4(4.90)(-20.0)}}{2(4.90)}$$

Reveal: Solving the Quadratic

Step 5: Calculate discriminant

$$\sqrt{(14.3)^2 + 4(4.90)(20.0)} = \sqrt{596.5} = 24.4$$

Reveal: Solving the Quadratic

Step 5: Calculate discriminant

$$\sqrt{(14.3)^2 + 4(4.90)(20.0)} = \sqrt{596.5} = 24.4$$

Step 6: Two solutions

$$t = \frac{14.3 + 24.4}{9.8} = 3.96 \text{ s} \quad (\text{physical})$$

$$t = \frac{14.3 - 24.4}{9.8} = -1.03 \text{ s} \quad (\text{reject})$$

Reveal: Solving the Quadratic

Step 5: Calculate discriminant

$$\sqrt{(14.3)^2 + 4(4.90)(20.0)} = \sqrt{596.5} = 24.4$$

Step 6: Two solutions

$$t = \frac{14.3 + 24.4}{9.8} = 3.96 \text{ s} \quad (\text{physical})$$

$$t = \frac{14.3 - 24.4}{9.8} = -1.03 \text{ s} \quad (\text{reject})$$

$t = 3.96 \text{ s}$

Reveal: Solving the Quadratic

Step 5: Calculate discriminant

$$\sqrt{(14.3)^2 + 4(4.90)(20.0)} = \sqrt{596.5} = 24.4$$

Step 6: Two solutions

$$t = \frac{14.3 + 24.4}{9.8} = 3.96 \text{ s} \quad (\text{physical})$$

$$t = \frac{14.3 - 24.4}{9.8} = -1.03 \text{ s} \quad (\text{reject})$$

$t = 3.96 \text{ s}$

Negative time means before launch - physically impossible!

What You Now Know

The Revelations

- 1 Vectors have magnitude AND direction

What You Now Know

The Revelations

- 1 Vectors have magnitude AND direction
- 2 Graphical addition: head-to-tail method

What You Now Know

The Revelations

- ① Vectors have magnitude AND direction
- ② Graphical addition: head-to-tail method
- ③ Analytical method: break into components, recombine

What You Now Know

The Revelations

- 1 Vectors have magnitude AND direction
- 2 Graphical addition: head-to-tail method
- 3 Analytical method: break into components, recombine
- 4 Projectile motion: horizontal and vertical are independent

What You Now Know

The Revelations

- 1 Vectors have magnitude AND direction
- 2 Graphical addition: head-to-tail method
- 3 Analytical method: break into components, recombine
- 4 Projectile motion: horizontal and vertical are independent
- 5 Time connects the two dimensions

What You Now Know

The Revelations

- 1 Vectors have magnitude AND direction
- 2 Graphical addition: head-to-tail method
- 3 Analytical method: break into components, recombine
- 4 Projectile motion: horizontal and vertical are independent
- 5 Time connects the two dimensions
- 6 Maximum height depends only on v_{0y}

Key Equations

Vector Components:

$$A_x = A \cos \theta \quad (1)$$

$$A_y = A \sin \theta \quad (2)$$

Magnitude and Direction:

$$A = \sqrt{A_x^2 + A_y^2} \quad (3)$$

$$\theta = \tan^{-1}(A_y/A_x) \quad (4)$$

Projectile Motion:

$$\text{Horizontal: } x = x_0 + v_x t \quad (5)$$

$$\text{Vertical: } y = y_0 + v_{0y} t - \frac{1}{2} g t^2 \quad (6)$$

$$\text{Max height: } h = \frac{v_{0y}^2}{2g} \quad (7)$$

Complete the assigned problems
posted on the LMS