

PHYS11 CH:9 The Currency of the Universe

How Energy Powers Everything

Mr. Gullo

December 2025

Outline

What if nothing ever stopped?

What if motion could last forever?

What if nothing ever stopped?

What if motion could last forever?

Balls bounce lower. Roller coasters slow down. Pendulums stop swinging.

What if nothing ever stopped?

What if motion could last forever?

Balls bounce lower. Roller coasters slow down. Pendulums stop swinging.

Where does the motion go?

The Roller Coaster Experience



Figure: Roller coaster: energy transformation in action

The Roller Coaster Experience



Figure: Roller coaster: energy transformation in action

Lifted to the top. Released. Speed builds. Climbs again. Slows down.

The Roller Coaster Experience



Figure: Roller coaster: energy transformation in action

Lifted to the top. Released. Speed builds. Climbs again. Slows down.
Same energy. Different forms.

Learning Objectives

By the end of this section, you will be able to:

- **9.1:** Describe and apply the work-energy theorem

Learning Objectives

By the end of this section, you will be able to:

- **9.1:** Describe and apply the work-energy theorem
- **9.1:** Describe and calculate work and power

9.1 What Is Work?

Civilian View vs. Reality

Civilian: "Homework is work. Holding a heavy box is hard work."

Physicist: "Neither is work. No motion, no work."

9.1 What Is Work?

Civilian View vs. Reality

Civilian: "Homework is work. Holding a heavy box is hard work."

Physicist: "Neither is work. No motion, no work."

The Universal Law of Work

$$W = Fd$$

Work equals **force** times **distance** moved in the direction of the **force**.

9.1 Three Examples

Consider these scenarios:

9.1 Three Examples

Consider these scenarios:

- 1 Homework - objects not moving over distance

9.1 Three Examples

Consider these scenarios:

- 1 Homework - objects not moving over distance
- 2 Lifting a rock upward - force moves object upward

9.1 Three Examples

Consider these scenarios:

- 1 Homework - objects not moving over distance
- 2 Lifting a rock upward - force moves object upward
- 3 Carrying a rock horizontally at constant speed

9.1 Three Examples

Consider these scenarios:

- 1 Homework - objects not moving over distance
- 2 Lifting a rock upward - force moves object upward
- 3 Carrying a rock horizontally at constant speed

The Mental Model

Work requires TWO things: **force** AND motion in the direction of **force**.

9.1 Energy: The Ability to Do Work

Two forms of mechanical **energy**:

9.1 Energy: The Ability to Do Work

Two forms of mechanical **energy**:

Kinetic Energy (KE)

Energy of motion. A moving object has **kinetic energy**.

9.1 Energy: The Ability to Do Work

Two forms of mechanical **energy**:

Kinetic Energy (KE)

Energy of motion. A moving object has **kinetic energy**.

Potential Energy (PE)

Stored **energy**. Gravitational **PE** is **energy** an object has due to its position above Earth's surface.

9.1 Energy: The Ability to Do Work

Two forms of mechanical **energy**:

Kinetic Energy (KE)

Energy of motion. A moving object has **kinetic energy**.

Potential Energy (PE)

Stored **energy**. Gravitational **PE** is **energy** an object has due to its position above Earth's surface.

Both measured in joules (J), same unit as **work**.

9.1 Gravitational Potential Energy

Lift a rock off the ground. You exert **force** upward over **distance**.

9.1 Gravitational Potential Energy

Lift a rock off the ground. You exert **force** upward over **distance**.

The **force** equals the rock's **weight**: $F = w = mg$

9.1 Gravitational Potential Energy

Lift a rock off the ground. You exert **force** upward over **distance**.

The **force** equals the rock's **weight**: $F = w = mg$

Work done equals **force** times **distance**:

Gravitational **Potential Energy**

$$PE = mgh$$

Potential energy equals mass times gravity times **height**.

9.1 Kinetic Energy

Drop the rock. Gravity does **work** on it. Rock speeds up.

9.1 Kinetic Energy

Drop the rock. Gravity does **work** on it. Rock speeds up.

Kinetic Energy

$$KE = \frac{1}{2}mv^2$$

Kinetic energy equals one-half mass times **velocity** squared.

9.1 Kinetic Energy

Drop the rock. Gravity does **work** on it. Rock speeds up.

Kinetic Energy

$$KE = \frac{1}{2}mv^2$$

Kinetic energy equals one-half mass times **velocity** squared.

The Mental Model

Heavier objects and faster objects have more **KE**. **Velocity** matters more because it's squared.

9.1 The Work-Energy Theorem

Nature's Source Code

$$W = \Delta KE = \frac{1}{2}mv_2^2 - \frac{1}{2}mv_1^2$$

Net **work** equals change in **kinetic energy**.

9.1 The Work-Energy Theorem

Nature's Source Code

$$W = \Delta KE = \frac{1}{2}mv_2^2 - \frac{1}{2}mv_1^2$$

Net **work** equals change in **kinetic energy**.

When you do **work** on an object, you change its **kinetic energy**.

9.1 The Work-Energy Theorem

Nature's Source Code

$$W = \Delta KE = \frac{1}{2}mv_2^2 - \frac{1}{2}mv_1^2$$

Net **work** equals change in **kinetic energy**.

When you do **work** on an object, you change its **kinetic energy**.

Subscripts: ₁ is initial, ₂ is final.

9.1 James Joule



Figure: James Joule (1818-1889)

9.1 James Joule



Figure: James Joule (1818-1889)

The joule (J) is named after this physicist who proved the work-energy connection.

9.1 James Joule



Figure: James Joule (1818-1889)

The joule (J) is named after this physicist who proved the work-energy connection.

$$1.0 \text{ J} = 1.0 \text{ N} \cdot \text{m} = 1.0 \text{ kg} \cdot \text{m}^2/\text{s}^2$$

9.1 Power: Rate of Doing Work

Work tells you how much. **Power** tells you how fast.

9.1 Power: Rate of Doing Work

Work tells you how much. **Power** tells you how fast.

Power

$$P = \frac{W}{t}$$

Power equals **work** divided by **time**.

9.1 Power: Rate of Doing Work

Work tells you how much. **Power** tells you how fast.

Power

$$P = \frac{W}{t}$$

Power equals **work** divided by **time**.

Unit: Watt (W)

9.1 Power: Rate of Doing Work

Work tells you how much. **Power** tells you how fast.

Power

$$P = \frac{W}{t}$$

Power equals **work** divided by **time**.

Unit: Watt (W)

One watt equals one joule per second.

9.1 Work vs. Power




Figure: Two ways to move a TV to the fourth floor

9.1 Work vs. Power




Figure: Two ways to move a TV to the fourth floor

Same work. Different power.

Pulley (2 min) generates more power than stairs (5 min).

9.1 James Watt and the Steam Engine




Figure: James Watt (1736-1819) - The unit of power

9.1 James Watt and the Steam Engine



Figure: James Watt (1736-1819) - The unit of power

Watt improved the steam engine, converting reciprocal motion to circular motion.

9.1 James Watt and the Steam Engine



Figure: James Watt (1736-1819) - The unit of power

Watt improved the steam engine, converting reciprocal motion to circular motion.

This innovation powered the industrial revolution.

Attempt: The Skater's Push

The Challenge (3 min, silent)

An ice skater with mass 50 kg glides at 8 m/s. Her friend pushes, increasing speed to 12 m/s.

Given:

- $m = 50 \text{ kg}$
- $v_1 = 8 \text{ m/s}$
- $v_2 = 12 \text{ m/s}$

Find: How much work did the friend do on the skater?

Can you calculate the work? Try it silently.

Compare: The Skater's Push

Turn and talk (2 min):

- 1 What equation did you use?
- 2 Did you calculate KE initial and KE final?
- 3 What operation connects them to work?

Compare: The Skater's Push

Turn and talk (2 min):

- 1 What equation did you use?
- 2 Did you calculate KE initial and KE final?
- 3 What operation connects them to work?

Name wheel: One pair share your approach (not your answer).

Reveal: The Skater's Push

Self-correct in a different color:

Equation: $W = \Delta KE = \frac{1}{2}mv_2^2 - \frac{1}{2}mv_1^2$

Reveal: The Skater's Push

Self-correct in a different color:

Equation: $W = \Delta KE = \frac{1}{2}mv_2^2 - \frac{1}{2}mv_1^2$

Factor out: $W = \frac{1}{2}m(v_2^2 - v_1^2)$

Reveal: The Skater's Push

Self-correct in a different color:

Equation: $W = \Delta KE = \frac{1}{2}mv_2^2 - \frac{1}{2}mv_1^2$

Factor out: $W = \frac{1}{2}m(v_2^2 - v_1^2)$

Substitute: $W = \frac{1}{2}(50)(12^2 - 8^2)$

Reveal: The Skater's Push

Self-correct in a different color:

Equation: $W = \Delta KE = \frac{1}{2}mv_2^2 - \frac{1}{2}mv_1^2$

Factor out: $W = \frac{1}{2}m(v_2^2 - v_1^2)$

Substitute: $W = \frac{1}{2}(50)(12^2 - 8^2)$

$$W = 25(144 - 64) = 25(80)$$

Reveal: The Skater's Push

Self-correct in a different color:

Equation: $W = \Delta KE = \frac{1}{2}mv_2^2 - \frac{1}{2}mv_1^2$

Factor out: $W = \frac{1}{2}m(v_2^2 - v_1^2)$

Substitute: $W = \frac{1}{2}(50)(12^2 - 8^2)$

$$W = 25(144 - 64) = 25(80)$$

$W = 2000 \text{ J}$

Reveal: The Skater's Push

Self-correct in a different color:

Equation: $W = \Delta KE = \frac{1}{2}mv_2^2 - \frac{1}{2}mv_1^2$

Factor out: $W = \frac{1}{2}m(v_2^2 - v_1^2)$

Substitute: $W = \frac{1}{2}(50)(12^2 - 8^2)$

$$W = 25(144 - 64) = 25(80)$$

$W = 2000 \text{ J}$

Check: Energy increased because friend did work. Reasonable!

Learning Objectives

By the end of this section, you will be able to:

- **9.2:** Explain the law of conservation of energy in terms of kinetic and potential energy

Learning Objectives

By the end of this section, you will be able to:

- **9.2:** Explain the law of conservation of energy in terms of kinetic and potential energy
- **9.2:** Perform calculations related to kinetic and potential energy and apply conservation of energy

9.2 The Universe's Accounting System

Energy is never created or destroyed.

It only transforms.

9.2 The Universe's Accounting System

Energy is never created or destroyed.

It only transforms.

The Law of Conservation of Energy

In a closed system, total energy remains constant.

9.2 The Universe's Accounting System

Energy is never created or destroyed.

It only transforms.

The Law of Conservation of Energy

In a closed system, total energy remains constant.

The Illusion

Civilian: "The ball lost energy when it stopped bouncing."

Physicist: "Energy transformed to heat from friction and sound."

9.2 Energy Transformations

Lift a rock to the top. You do **work**. Rock gains **PE**.

9.2 Energy Transformations

Lift a rock to the top. You do **work**. Rock gains **PE**.
Release the rock. **PE** converts to **KE** as it falls.

9.2 Energy Transformations

Lift a rock to the top. You do **work**. Rock gains **PE**.
Release the rock. **PE** converts to **KE** as it falls.
Hit the ground. **KE** converts to heat and sound.

9.2 Energy Transformations

Lift a rock to the top. You do **work**. Rock gains **PE**.

Release the rock. **PE** converts to **KE** as it falls.

Hit the ground. **KE** converts to heat and sound.

The Mental Model

Energy changes form constantly. The total amount stays the same.

9.2 The Roller Coaster




Figure: Energy transformations on a roller coaster

9.2 The Roller Coaster




Figure: Energy transformations on a roller coaster

Top: High PE, low KE (slow)

9.2 The Roller Coaster




Figure: Energy transformations on a roller coaster

Top: High PE, low KE (slow)

Bottom: Low PE, high KE (fast)

9.2 The Roller Coaster




Figure: Energy transformations on a roller coaster

Top: High PE, low KE (slow)

Bottom: Low PE, high KE (fast)

Next hill: KE converts back to PE

9.2 Conservation Equation

Assume no **energy** lost to friction.

9.2 Conservation Equation

Assume no **energy** lost to friction.

Conservation of Mechanical **Energy**

$$KE_1 + PE_1 = KE_2 + PE_2$$

Initial **kinetic** plus initial **potential** equals final **kinetic** plus final **potential**.

9.2 Conservation Equation

Assume no **energy** lost to friction.

Conservation of Mechanical **Energy**

$$KE_1 + PE_1 = KE_2 + PE_2$$

Initial **kinetic** plus initial **potential** equals final **kinetic** plus final **potential**.

Either side equals total mechanical **energy**.

9.2 Conservation Equation

Assume no **energy** lost to friction.

Conservation of Mechanical **Energy**

$$KE_1 + PE_1 = KE_2 + PE_2$$

Initial **kinetic** plus initial **potential** equals final **kinetic** plus final **potential**.

Either side equals total mechanical **energy**.

Closed system: No **energy** lost to surroundings.

9.2 Equations Summary

$$KE = \frac{1}{2}mv^2$$

$$PE = mgh$$

$$KE_1 + PE_1 = KE_2 + PE_2$$

9.2 Equations Summary

$$KE = \frac{1}{2}mv^2$$

$$PE = mgh$$

$$KE_1 + PE_1 = KE_2 + PE_2$$

Combining them:

$$\frac{1}{2}mv_1^2 + mgh_1 = \frac{1}{2}mv_2^2 + mgh_2$$

9.2 Equations Summary

$$KE = \frac{1}{2}mv^2$$

$$PE = mgh$$

$$KE_1 + PE_1 = KE_2 + PE_2$$

Combining them:

$$\frac{1}{2}mv_1^2 + mgh_1 = \frac{1}{2}mv_2^2 + mgh_2$$

Pro tip: Mass often cancels out!

Attempt: The Falling Rock

The Challenge (3 min, silent)

A 10 kg rock falls from a 20 m cliff. When it has fallen 10 m, what are its KE and PE?

Given:

- $m = 10 \text{ kg}$
- $h_1 = 20 \text{ m}$ (initial height)
- $h_2 = 10 \text{ m}$ (after falling 10 m)
- $v_1 = 0$ (dropped from rest)
- $g = 9.8 \text{ m/s}^2$

Find: KE_2 and PE_2

Can you use conservation of energy? Work silently.

Compare: The Falling Rock

Turn and talk (2 min):

- 1 What is the initial KE? Why?
- 2 How did you calculate PE at 10 m height?
- 3 How did you find KE at 10 m height?

Compare: The Falling Rock

Turn and talk (2 min):

- 1 What is the initial KE? Why?
- 2 How did you calculate PE at 10 m height?
- 3 How did you find KE at 10 m height?

Name wheel: One pair share your approach (not your answer).

Reveal: The Falling Rock

Self-correct in a different color:

Initial energy: $KE_1 = 0$ (at rest), $PE_1 = mgh_1 = 10 \times 9.8 \times 20 = 1960 \text{ J}$

Reveal: The Falling Rock

Self-correct in a different color:

Initial **energy**: $KE_1 = 0$ (at rest), $PE_1 = mgh_1 = 10 \times 9.8 \times 20 = 1960 \text{ J}$

At 10 m **height**: $PE_2 = mgh_2 = 10 \times 9.8 \times 10 = 980 \text{ J}$

Reveal: The Falling Rock

Self-correct in a different color:

Initial **energy**: $KE_1 = 0$ (at rest), $PE_1 = mgh_1 = 10 \times 9.8 \times 20 = 1960 \text{ J}$

At 10 m **height**: $PE_2 = mgh_2 = 10 \times 9.8 \times 10 = 980 \text{ J}$

Conservation: $KE_1 + PE_1 = KE_2 + PE_2$

Reveal: The Falling Rock

Self-correct in a different color:

Initial **energy**: $KE_1 = 0$ (at rest), $PE_1 = mgh_1 = 10 \times 9.8 \times 20 = 1960 \text{ J}$

At 10 m **height**: $PE_2 = mgh_2 = 10 \times 9.8 \times 10 = 980 \text{ J}$

Conservation: $KE_1 + PE_1 = KE_2 + PE_2$

$$0 + 1960 = KE_2 + 980$$

Reveal: The Falling Rock

Self-correct in a different color:

Initial **energy**: $KE_1 = 0$ (at rest), $PE_1 = mgh_1 = 10 \times 9.8 \times 20 = 1960 \text{ J}$

At 10 m **height**: $PE_2 = mgh_2 = 10 \times 9.8 \times 10 = 980 \text{ J}$

Conservation: $KE_1 + PE_1 = KE_2 + PE_2$

$$0 + 1960 = KE_2 + 980$$

$KE_2 = 980 \text{ J}$

Reveal: The Falling Rock

Self-correct in a different color:

Initial **energy**: $KE_1 = 0$ (at rest), $PE_1 = mgh_1 = 10 \times 9.8 \times 20 = 1960 \text{ J}$

At 10 m **height**: $PE_2 = mgh_2 = 10 \times 9.8 \times 10 = 980 \text{ J}$

Conservation: $KE_1 + PE_1 = KE_2 + PE_2$

$$0 + 1960 = KE_2 + 980$$

$KE_2 = 980 \text{ J}$

Check: Lost 980 J of **PE**, gained 980 J of **KE**. **Energy** conserved!

9.2 Why Conservation Works

Total mechanical **energy** at any point equals **energy** at start.

9.2 Why Conservation Works

Total mechanical **energy** at any point equals **energy** at start.

At the top of roller coaster: mostly **PE**, little **KE**

9.2 Why Conservation Works

Total mechanical **energy** at any point equals **energy** at start.

At the top of roller coaster: mostly **PE**, little **KE**

At the bottom: mostly **KE**, little **PE**

9.2 Why Conservation Works

Total mechanical **energy** at any point equals **energy** at start.

At the top of roller coaster: mostly **PE**, little **KE**

At the bottom: mostly **KE**, little **PE**

At any point in between: **PE** + **KE** = constant

9.2 Why Conservation Works

Total mechanical **energy** at any point equals **energy** at start.

At the top of roller coaster: mostly **PE**, little **KE**

At the bottom: mostly **KE**, little **PE**

At any point in between: **PE** + **KE** = constant

The Paradox

Your brain says: "The roller coaster is slowing down, losing **energy**."

Reality: "**KE** converting to **PE**. Total **energy** unchanged."

9.2 Real Systems and Friction

Ideal system: No friction, **energy** perfectly conserved

9.2 Real Systems and Friction

Ideal system: No friction, **energy** perfectly conserved

Real system: Friction converts mechanical **energy** to heat

9.2 Real Systems and Friction

Ideal system: No friction, **energy** perfectly conserved

Real system: Friction converts mechanical **energy** to heat

The Mental Model

Friction is not a violation of conservation. Heat is still **energy**, just less useful.

9.2 Real Systems and Friction

Ideal system: No friction, **energy** perfectly conserved

Real system: Friction converts mechanical **energy** to heat

The Mental Model

Friction is not a violation of conservation. Heat is still **energy**, just less useful.

Engineers design to minimize friction losses.

9.2 The Closed System

Closed system: No **energy** enters or leaves

9.2 The Closed System

Closed system: No **energy** enters or leaves
In reality, perfectly closed systems don't exist.

9.2 The Closed System

Closed system: No **energy** enters or leaves

In reality, perfectly closed systems don't exist.

But many systems are approximately closed for short times.

9.2 The Closed System

Closed system: No **energy** enters or leaves

In reality, perfectly closed systems don't exist.

But many systems are approximately closed for short times.

The Approximation

Falling objects, roller coasters, pendulums: closed system is good approximation if friction is small.

What You Now Know

The Revelations

① **Work** = **force** times **distance** (only if motion happens)

What You Now Know

The Revelations

- 1 **Work** = **force** times **distance** (only if motion happens)
- 2 **Work** changes **energy** (work-energy theorem)

What You Now Know

The Revelations

- 1 **Work** = **force** times **distance** (only if motion happens)
- 2 **Work** changes **energy** (work-energy theorem)
- 3 **Energy** has two forms: **kinetic** (motion) and **potential** (stored)

What You Now Know

The Revelations

- 1 **Work** = **force** times **distance** (only if motion happens)
- 2 **Work** changes **energy** (work-energy theorem)
- 3 **Energy** has two forms: **kinetic** (motion) and **potential** (stored)
- 4 **Power** = rate of doing **work**

What You Now Know

The Revelations

- 1 **Work** = **force** times **distance** (only if motion happens)
- 2 **Work** changes **energy** (work-energy theorem)
- 3 **Energy** has two forms: **kinetic** (motion) and **potential** (stored)
- 4 **Power** = rate of doing **work**
- 5 **Energy** is conserved: $KE + PE = \text{constant}$

What You Now Know

The Revelations

- 1 **Work** = **force** times **distance** (only if motion happens)
- 2 **Work** changes **energy** (work-energy theorem)
- 3 **Energy** has two forms: **kinetic** (motion) and **potential** (stored)
- 4 **Power** = rate of doing **work**
- 5 **Energy** is conserved: $KE + PE = \text{constant}$
- 6 **Energy** transforms but never vanishes

Key Equations

$$W = Fd$$

$$KE = \frac{1}{2}mv^2$$

$$PE = mgh$$

$$W = \Delta KE = \frac{1}{2}mv_2^2 - \frac{1}{2}mv_1^2$$

$$P = \frac{W}{t}$$

$$KE_1 + PE_1 = KE_2 + PE_2$$

Complete the assigned problems
posted on the LMS

Temporary page!

\LaTeX was unable to guess the total number of pages correctly. There was some unprocessed data that should have been added to the document, so this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will disappear, because \LaTeX now knows how many pages to expect for the document.