

# PHYS11 CH:18 The Force That Moves Everything

## Electric Charge and Conservation

Mr. Gullo

December 2025

# Outline

- 1 Introduction
- 2 Electrical Charges, Conservation of Charge, and Transfer of Charge
- 3 Worked Example
- 4 Practice Application
- 5 Summary

# Why Does Hair Stand on End?



# Why Does Hair Stand on End?



*What invisible force makes strands of hair repel each other?*

What if everything you touch  
*is held together by invisible forces?*

# The Invisible World

What if everything you touch  
*is held together by invisible forces?*

The atoms in your fingertips never actually touch the atoms in this desk...

# The Invisible World

What if everything you touch  
*is held together by invisible forces?*

The atoms in your fingertips never actually touch the atoms in this desk...

Electric forces keep them apart.

# Learning Objectives

By the end of this lesson, you will be able to:

- **18.1:** Describe positive and negative electric charges



# Learning Objectives

By the end of this lesson, you will be able to:

- **18.1:** Describe positive and negative electric charges
- **18.1:** Use conservation of charge to calculate charge transfers

# Learning Objectives

By the end of this lesson, you will be able to:

- **18.1:** Describe positive and negative electric charges
- **18.1:** Use conservation of charge to calculate charge transfers
- **18.1:** Characterize conductors vs insulators

# Learning Objectives

By the end of this lesson, you will be able to:

- **18.1:** Describe positive and negative electric charges
- **18.1:** Use conservation of charge to calculate charge transfers
- **18.1:** Characterize conductors vs insulators
- **18.1:** Describe electric polarization and charging by induction

# 18.1 Two Types of Charge

## Nature's Binary Code

Electric charge is a property of matter that causes objects to attract or repel each other.

# 18.1 Two Types of Charge

## Nature's Binary Code

Electric charge is a property of matter that causes objects to attract or repel each other.

### The discovery:

- Glass rubbed with silk: glass becomes positive, silk becomes negative

# 18.1 Two Types of Charge

## Nature's Binary Code

Electric charge is a property of matter that causes objects to attract or repel each other.

### The discovery:

- Glass rubbed with silk: glass becomes positive, silk becomes negative
- Like charges **repel** each other

# 18.1 Two Types of Charge

## Nature's Binary Code

Electric charge is a property of matter that causes objects to attract or repel each other.

### The discovery:

- Glass rubbed with silk: glass becomes positive, silk becomes negative
- Like charges **repel** each other
- Unlike charges **attract** each other

# 18.1 Two Types of Charge

## Nature's Binary Code

Electric charge is a property of matter that causes objects to attract or repel each other.

### The discovery:

- Glass rubbed with silk: glass becomes positive, silk becomes negative
- Like charges **repel** each other
- Unlike charges **attract** each other

## The Mental Model


Charge is nature's binary system: positive or negative. No neutral charge exists.



# 18.1 Experimental Evidence



(a)




(b)



(c)

# 18.1 Experimental Evidence



**Pattern:** Glass rods repel each other. Silk cloths repel each other. Glass attracts silk.

# 18.1 The Discovery of the Electron

**The mystery (1897):** What carries negative charge?

# 18.1 The Discovery of the Electron

**The mystery (1897):** What carries negative charge?

**The experiment:** J.J. Thomson studied cathode rays

# 18.1 The Discovery of the Electron

**The mystery (1897):** What carries negative charge?

**The experiment:** J.J. Thomson studied cathode rays

**The revelation:** Cathode rays are particles carrying negative charge

# 18.1 The Discovery of the Electron

**The mystery (1897):** What carries negative charge?



**The experiment:** J.J. Thomson studied cathode rays

**The revelation:** Cathode rays are particles carrying negative charge



## Universal Law: The Electron

The electron carries the fundamental unit of *negative* electric charge.

# 18.1 Inside the Atom



# 18.1 Inside the Atom



Rutherford's model: electrons orbit a tiny, dense nucleus of protons.



# 18.1 The Fundamental Unit of Charge

## Nature's Quantum

$$e = 1.602 \times 10^{-19} \text{ C}$$

The fundamental unit of electric charge (magnitude).

# 18.1 The Fundamental Unit of Charge

## Nature's Quantum

$$e = 1.602 \times 10^{-19} \text{ C}$$

The fundamental unit of electric charge (magnitude).

### The carriers:

- Proton charge:  $+e = +1.602 \times 10^{-19} \text{ C}$

# 18.1 The Fundamental Unit of Charge

## Nature's Quantum

$$e = 1.602 \times 10^{-19} \text{ C}$$

The fundamental unit of electric charge (magnitude).

### The carriers:

- Proton charge:  $+e = +1.602 \times 10^{-19} \text{ C}$
- Electron charge:  $-e = -1.602 \times 10^{-19} \text{ C}$

# 18.1 The Fundamental Unit of Charge

## Nature's Quantum

$$e = 1.602 \times 10^{-19} \text{ C}$$

The fundamental unit of electric charge (magnitude).

### The carriers:


- Proton charge:  $+e = +1.602 \times 10^{-19} \text{ C}$
- Electron charge:  $-e = -1.602 \times 10^{-19} \text{ C}$

## The Paradox


**Civilian:** "Why is charge so tiny?"

**Physicist:** "It takes  $6.25 \times 10^{18}$  protons to make just 1 coulomb!"

# 18.1 Measuring the Electron Charge



# 18.1 Measuring the Electron Charge



## Millikan Oil-Drop Experiment (1909):

- Spray oil droplets between charged plates
- Balance electric force against gravity
- Measure charge on individual drops

# 18.1 Charge Quantization

**Discovery:** Charge always comes in multiples of  $e$

# 18.1 Charge Quantization

**Discovery:** Charge always comes in multiples of  $e$

Universal Law: Charge Quantization

$$Q = ne$$

where  $n$  is an integer ( $\pm 1, \pm 2, \pm 3, \dots$ )



# 18.1 Charge Quantization

**Discovery:** Charge always comes in multiples of  $e$

Universal Law: Charge Quantization

$$Q = ne$$

where  $n$  is an integer ( $\pm 1, \pm 2, \pm 3, \dots$ )

**Meaning:** You can have 5 electrons or 5 million, but never 5.5 electrons.

# 18.1 Conservation of Charge

## Nature's Accounting System

$$q_{\text{initial}} = q_{\text{final}}$$

Electric charge cannot be created or destroyed.

# 18.1 Conservation of Charge

## Nature's Accounting System

$$q_{\text{initial}} = q_{\text{final}}$$

Electric charge cannot be created or destroyed.

### What this means:

- Total charge before interaction = total charge after

# 18.1 Conservation of Charge

## Nature's Accounting System

$$q_{\text{initial}} = q_{\text{final}}$$

Electric charge cannot be created or destroyed.

### What this means:

- Total charge before interaction = total charge after
- Charge can move, but net charge stays constant

# 18.1 Conservation of Charge

## Nature's Accounting System

$$q_{\text{initial}} = q_{\text{final}}$$

Electric charge cannot be created or destroyed.

### What this means:

- Total charge before interaction = total charge after
- Charge can move, but net charge stays constant
- Most fundamental conservation law in physics

# 18.1 Conductors vs Insulators

## Conductors

Materials that allow charge to move freely

### Examples:

- Metals (copper, silver, aluminum)
- Electrons loosely bound

# 18.1 Conductors vs Insulators

## Conductors

Materials that allow charge to move freely

### Examples:

- Metals (copper, silver, aluminum)
- Electrons loosely bound

## Insulators

Materials that prevent charge from moving

### Examples:

- Rubber, plastic, glass, wood
- Electrons tightly bound

# 18.1 Conductors vs Insulators

## Conductors

Materials that allow charge to move freely

### Examples:

- Metals (copper, silver, aluminum)
- Electrons loosely bound

## Insulators

Materials that prevent charge from moving

### Examples:


- Rubber, plastic, glass, wood
- Electrons tightly bound

## The Mental Model


Conductor = highway for electrons. Insulator = roadblock.



## 18.1 The Conductivity Spectrum





# 18.1 The Conductivity Spectrum




**Semiconductors:** Between conductors and insulators (silicon, germanium)

# 18.1 Charge Distribution



# 18.1 Charge Distribution



**Conductor:** Charge spreads to outer surface (repulsion wins).

**Insulator:** Charge stays in place (can't move).

# 18.1 Transferring Charge: Contact

**Charging by contact:** Surfaces touch and share electrons

# 18.1 Transferring Charge: Contact

**Charging by contact:** Surfaces touch and share electrons

**Example:** Walking across carpet

- Shoes rub carpet

# 18.1 Transferring Charge: Contact

**Charging by contact:** Surfaces touch and share electrons

**Example:** Walking across carpet

- Shoes rub carpet
- Electrons transfer from carpet to shoes

# 18.1 Transferring Charge: Contact

**Charging by contact:** Surfaces touch and share electrons

**Example:** Walking across carpet

- Shoes rub carpet
- Electrons transfer from carpet to shoes
- You accumulate excess negative charge



# 18.1 Transferring Charge: Contact

**Charging by contact:** Surfaces touch and share electrons

**Example:** Walking across carpet

- Shoes rub carpet
- Electrons transfer from carpet to shoes
- You accumulate excess negative charge
- Touch doorknob → ZAP!

# 18.1 Transferring Charge: Contact

**Charging by contact:** Surfaces touch and share electrons



**Example:** Walking across carpet

- Shoes rub carpet
- Electrons transfer from carpet to shoes
- You accumulate excess negative charge
- Touch doorknob → ZAP!


## Real-World: Static Shock

Rubbing increases contact between materials, transferring more electrons.

# 18.1 Transferring Charge: Conduction




## 18.1 Transferring Charge: Conduction



**Charging by conduction:** Touching charged object to neutral object

# 18.1 Transferring Charge: Conduction



**Charging by conduction:** Touching charged object to neutral object  
Charges redistribute to equalize - like water finding same level.

# 18.1 Polarization

**What is polarization?** Separation of positive and negative charges within an object

# 18.1 Polarization

**What is polarization?** Separation of positive and negative charges within an object

**Example:** Child on slide (Figure 18.1)

- Excess positive charge on body

# 18.1 Polarization

**What is polarization?** Separation of positive and negative charges within an object

**Example:** Child on slide (Figure 18.1)

- Excess positive charge on body
- Positive charges repel each other



# 18.1 Polarization

**What is polarization?** Separation of positive and negative charges within an object

**Example:** Child on slide (Figure 18.1)

- Excess positive charge on body
- Positive charges repel each other
- Spread to extremities (hair)

# 18.1 Polarization

**What is polarization?** Separation of positive and negative charges within an object

**Example:** Child on slide (Figure 18.1)

- Excess positive charge on body
- Positive charges repel each other
- Spread to extremities (hair)
- Hair strands repel → stand on end!

# 18.1 Polarization

**What is polarization?** Separation of positive and negative charges within an object


**Example:** Child on slide (Figure 18.1)

- Excess positive charge on body
- Positive charges repel each other
- Spread to extremities (hair)
- Hair strands repel → stand on end!


## The Mental Model

Like dominos - push on one end, disturbance travels to other end.

# 18.1 Charging by Induction




# 18.1 Charging by Induction




**Induction:** Creating charge separation by approaching a charged object (no touching!)

# 18.1 Van de Graaff Generator



# 18.1 Van de Graaff Generator



## How it works:

- Belt transfers electrons to metal globe
- Electrons spread over outer surface (repulsion)
- Can accumulate millions of volts!

# 18.1 Hair-Raising Physics





# 18.1 Hair-Raising Physics



## Why does hair stand up?

- Each hair strand gets excess charge (same sign)
- Like charges repel
- Strands push away from each other as far as possible

# Attempt: Conservation Challenge

## The Challenge (3 min, silent)

Two metal spheres initially have charges of  $+4\text{ C}$  and  $+8\text{ C}$ . After touching each other, one sphere has  $+10\text{ C}$ .

### Given:

- Blue sphere initial:  $q_1 = +4\text{ C}$
- Red sphere initial:  $q_2 = +8\text{ C}$
- Blue sphere final:  $q'_1 = +10\text{ C}$

**Find:** Final charge on red sphere  $q'_2$

*Can you use conservation of charge? Work silently.*

# Compare: Conservation Strategy

**Turn and talk (2 min):**

- 1 What law did you use?
- 2 What is the total initial charge?
- 3 How did you find the final charge on the red sphere?

# Compare: Conservation Strategy

## Turn and talk (2 min):

- 1 What law did you use?
- 2 What is the total initial charge?
- 3 How did you find the final charge on the red sphere?

**Name wheel:** One pair share your approach (not your answer).

# Reveal: Conservation in Action

**Self-correct in a different color:**

**Law:**  $q_{\text{initial}} = q_{\text{final}}$

# Reveal: Conservation in Action

**Self-correct in a different color:**

**Law:**  $q_{\text{initial}} = q_{\text{final}}$

**Step 1:** Find total initial charge

$$q_{\text{initial}} = +4 \text{ C} + 8 \text{ C} = +12 \text{ C}$$

# Reveal: Conservation in Action

**Self-correct in a different color:**

**Law:**  $q_{\text{initial}} = q_{\text{final}}$

**Step 1:** Find total initial charge

$$q_{\text{initial}} = +4 \text{ C} + 8 \text{ C} = +12 \text{ C}$$

**Step 2:** Apply conservation law

$$q_{\text{final}} = +10 \text{ C} + q'_2$$

# Reveal: Conservation in Action

**Self-correct in a different color:**

**Law:**  $q_{\text{initial}} = q_{\text{final}}$

**Step 1:** Find total initial charge

$$q_{\text{initial}} = +4 \text{ C} + 8 \text{ C} = +12 \text{ C}$$

**Step 2:** Apply conservation law

$$q_{\text{final}} = +10 \text{ C} + q'_2$$

**Step 3:** Solve for  $q'_2$

$$+12 \text{ C} = +10 \text{ C} + q'_2$$

$$q'_2 = +2 \text{ C}$$



# Reveal: Conservation in Action

**Self-correct in a different color:**

**Law:**  $q_{\text{initial}} = q_{\text{final}}$

**Step 1:** Find total initial charge

$$q_{\text{initial}} = +4 \text{ C} + 8 \text{ C} = +12 \text{ C}$$

**Step 2:** Apply conservation law

$$q_{\text{final}} = +10 \text{ C} + q'_2$$

**Step 3:** Solve for  $q'_2$

$$+12 \text{ C} = +10 \text{ C} + q'_2$$

$$q'_2 = +2 \text{ C}$$

**Check:**  $12 = 10 + 2$  ✓ Charge is conserved!

# Attempt: Counting Electrons

## The Challenge (3 min, silent)

An ink droplet in a printer has net charge  $q = -1.0 \times 10^{-10} \text{ C}$  after passing through an electron beam.

### Given:

- Droplet charge:  $q = -1.0 \times 10^{-10} \text{ C}$
- Electron charge:  $e = -1.602 \times 10^{-19} \text{ C}$

**Find:** Number of electrons captured by droplet

*How many electrons does it take? Work silently.*

# Compare: Quantization Strategy

## Turn and talk (2 min):

- 1 What equation relates total charge to number of electrons?
- 2 Did you multiply or divide?
- 3 What happens with the negative signs?

# Compare: Quantization Strategy

## Turn and talk (2 min):

- 1 What equation relates total charge to number of electrons?
- 2 Did you multiply or divide?
- 3 What happens with the negative signs?

**Name wheel:** One pair share your approach (not your answer).

# Reveal: Quantization at Work

**Self-correct in a different color:**

**Equation:**  $Q = ne$ , so  $n = \frac{Q}{e}$

# Reveal: Quantization at Work

**Self-correct in a different color:**

**Equation:**  $Q = ne$ , so  $n = \frac{Q}{e}$

**Substitute:**

$$n = \frac{q}{q_{e^-}} = \frac{-1.0 \times 10^{-10} \text{ C}}{-1.602 \times 10^{-19} \text{ C}}$$

# Reveal: Quantization at Work

**Self-correct in a different color:**

**Equation:**  $Q = ne$ , so  $n = \frac{Q}{e}$

**Substitute:**

$$n = \frac{q}{q_{e^-}} = \frac{-1.0 \times 10^{-10} \text{ C}}{-1.602 \times 10^{-19} \text{ C}}$$

**Simplify:**

$$n = \frac{1.0 \times 10^{-10}}{1.602 \times 10^{-19}} = 6.24 \times 10^8$$

# Reveal: Quantization at Work

**Self-correct in a different color:**

**Equation:**  $Q = ne$ , so  $n = \frac{Q}{e}$

**Substitute:**

$$n = \frac{q}{q_{e^-}} = \frac{-1.0 \times 10^{-10} \text{ C}}{-1.602 \times 10^{-19} \text{ C}}$$

**Simplify:**

$$n = \frac{1.0 \times 10^{-10}}{1.602 \times 10^{-19}} = 6.24 \times 10^8$$

$n = 6 \times 10^8 \text{ electrons}$



# Reveal: Quantization at Work

**Self-correct in a different color:**

**Equation:**  $Q = ne$ , so  $n = \frac{Q}{e}$

**Substitute:**

$$n = \frac{q}{q_{e^-}} = \frac{-1.0 \times 10^{-10} \text{ C}}{-1.602 \times 10^{-19} \text{ C}}$$

**Simplify:**

$$n = \frac{1.0 \times 10^{-10}}{1.602 \times 10^{-19}} = 6.24 \times 10^8$$

$n = 6 \times 10^8 \text{ electrons}$

**Check:** About 600 million electrons - seems large but atoms have  $10^{16}$  atoms!

# What You Now Know

## The Revelations

- 1 Two types of charge: positive (protons) and negative (electrons)

# What You Now Know

## The Revelations

- 1 Two types of charge: positive (protons) and negative (electrons)
- 2 Fundamental charge:  $e = 1.602 \times 10^{-19} \text{ C}$

# What You Now Know

## The Revelations

- 1 Two types of charge: positive (protons) and negative (electrons)
- 2 Fundamental charge:  $e = 1.602 \times 10^{-19} \text{ C}$
- 3 Charge quantization:  $Q = ne$  (only integer multiples)

# What You Now Know

## The Revelations

- 1 Two types of charge: positive (protons) and negative (electrons)
- 2 Fundamental charge:  $e = 1.602 \times 10^{-19} \text{ C}$
- 3 Charge quantization:  $Q = ne$  (only integer multiples)
- 4 Conservation:  $q_{\text{initial}} = q_{\text{final}}$  (never violated)

# What You Now Know

## The Revelations

- 1 Two types of charge: positive (protons) and negative (electrons)
- 2 Fundamental charge:  $e = 1.602 \times 10^{-19} \text{ C}$
- 3 Charge quantization:  $Q = ne$  (only integer multiples)
- 4 Conservation:  $q_{\text{initial}} = q_{\text{final}}$  (never violated)
- 5 Conductors let charge move; insulators don't

# What You Now Know

## The Revelations

- 1 Two types of charge: positive (protons) and negative (electrons)
- 2 Fundamental charge:  $e = 1.602 \times 10^{-19} \text{ C}$
- 3 Charge quantization:  $Q = ne$  (only integer multiples)
- 4 Conservation:  $q_{\text{initial}} = q_{\text{final}}$  (never violated)
- 5 Conductors let charge move; insulators don't
- 6 Transfer methods: contact, conduction, induction

# Key Equations

$$e = 1.602 \times 10^{-19} \text{ C} \quad (\text{fundamental charge}) \quad (1)$$

$$Q = ne \quad (\text{charge quantization}) \quad (2)$$

$$q_{\text{initial}} = q_{\text{final}} \quad (\text{conservation of charge}) \quad (3)$$



Complete the assigned problems  
posted on the LMS