

PHYS11 CH:1-3

Foundations of Physics, Motion, and Kinematics

Mr. Gullo

July 23, 2025

Learning Objectives

- Define physics and its role as a fundamental science.

Learning Objectives

- Define physics and its role as a fundamental science.
- Differentiate between physical quantities, units, accuracy, and precision.

Learning Objectives

- Define physics and its role as a fundamental science.
- Differentiate between physical quantities, units, accuracy, and precision.
- Distinguish between scalar quantities (distance, speed) and vector quantities (displacement, velocity).

Learning Objectives

- Define physics and its role as a fundamental science.
- Differentiate between physical quantities, units, accuracy, and precision.
- Distinguish between scalar quantities (distance, speed) and vector quantities (displacement, velocity).
- Define acceleration as the rate of change of velocity.

Learning Objectives

- Define physics and its role as a fundamental science.
- Differentiate between physical quantities, units, accuracy, and precision.
- Distinguish between scalar quantities (distance, speed) and vector quantities (displacement, velocity).
- Define acceleration as the rate of change of velocity.
- Analyze and interpret motion using position-time and velocity-time graphs.

Learning Objectives

- Define physics and its role as a fundamental science.
- Differentiate between physical quantities, units, accuracy, and precision.
- Distinguish between scalar quantities (distance, speed) and vector quantities (displacement, velocity).
- Define acceleration as the rate of change of velocity.
- Analyze and interpret motion using position-time and velocity-time graphs.
- Understand the meaning of slope and area in motion graphs.

Learning Objectives

- Define physics and its role as a fundamental science.
- Differentiate between physical quantities, units, accuracy, and precision.
- Distinguish between scalar quantities (distance, speed) and vector quantities (displacement, velocity).
- Define acceleration as the rate of change of velocity.
- Analyze and interpret motion using position-time and velocity-time graphs.
- Understand the meaning of slope and area in motion graphs.
- Identify and apply the kinematic equations for motion with constant acceleration.

What is Physics?

- Physics is the most fundamental science.

What is Physics?

- Physics is the most fundamental science.
- Physics studies four main things:

What is Physics?

- Physics is the most fundamental science.
- Physics studies four main things:
 - **Energy** - the ability to do work
 - **Matter** - anything that has mass and takes up space
 - **Space** - the dimensions around us
 - **Time** - how events happen in sequence

What is Physics?

- Physics is the most fundamental science.
- Physics studies four main things:
 - **Energy** - the ability to do work
 - **Matter** - anything that has mass and takes up space
 - **Space** - the dimensions around us
 - **Time** - how events happen in sequence
- Physics provides the foundation for other sciences like chemistry, biology, and geology.

What is Physics?

- Physics is the most fundamental science.
- Physics studies four main things:
 - **Energy** - the ability to do work
 - **Matter** - anything that has mass and takes up space
 - **Space** - the dimensions around us
 - **Time** - how events happen in sequence
- Physics provides the foundation for other sciences like chemistry, biology, and geology.
- **Modern Physics** includes two important theories:

What is Physics?

- Physics is the most fundamental science.
- Physics studies four main things:
 - **Energy** - the ability to do work
 - **Matter** - anything that has mass and takes up space
 - **Space** - the dimensions around us
 - **Time** - how events happen in sequence
- Physics provides the foundation for other sciences like chemistry, biology, and geology.
- **Modern Physics** includes two important theories:
 - **Relativity**: Shows how time, space, and gravity work differently for different observers.

What is Physics?

- Physics is the most fundamental science.
- Physics studies four main things:
 - **Energy** - the ability to do work
 - **Matter** - anything that has mass and takes up space
 - **Space** - the dimensions around us
 - **Time** - how events happen in sequence
- Physics provides the foundation for other sciences like chemistry, biology, and geology.
- **Modern Physics** includes two important theories:
 - **Relativity**: Shows how time, space, and gravity work differently for different observers.
 - **Quantum Mechanics**: Explains how very small particles behave.

Physical Quantities & Units

- A **physical quantity** is a property of an object that can be measured (e.g., length, mass, time).

Physical Quantities & Units

- A **physical quantity** is a property of an object that can be measured (e.g., length, mass, time).
- We use the **SI system** (metric system) for consistency.

Physical Quantities & Units

- A **physical quantity** is a property of an object that can be measured (e.g., length, mass, time).
- We use the **SI system** (metric system) for consistency.
- The four fundamental units we will use are:

Physical Quantities & Units

- A **physical quantity** is a property of an object that can be measured (e.g., length, mass, time).
- We use the **SI system** (metric system) for consistency.
- The four fundamental units we will use are:
 - Length: **meter (m)**

Physical Quantities & Units

- A **physical quantity** is a property of an object that can be measured (e.g., length, mass, time).
- We use the **SI system** (metric system) for consistency.
- The four fundamental units we will use are:
 - Length: **meter (m)**
 - Mass: **kilogram (kg)**

Physical Quantities & Units

- A **physical quantity** is a property of an object that can be measured (e.g., length, mass, time).
- We use the **SI system** (metric system) for consistency.
- The four fundamental units we will use are:
 - Length: **meter (m)**
 - Mass: **kilogram (kg)**
 - Time: **second (s)**

Physical Quantities & Units

- A **physical quantity** is a property of an object that can be measured (e.g., length, mass, time).
- We use the **SI system** (metric system) for consistency.
- The four fundamental units we will use are:
 - Length: **meter (m)**
 - Mass: **kilogram (kg)**
 - Time: **second (s)**
 - Electric Current: **ampere (A)**

Physical Quantities & Units

- A **physical quantity** is a property of an object that can be measured (e.g., length, mass, time).
- We use the **SI system** (metric system) for consistency.
- The four fundamental units we will use are:
 - Length: **meter (m)**
 - Mass: **kilogram (kg)**
 - Time: **second (s)**
 - Electric Current: **ampere (A)**
- Unit conversions are essential for solving problems correctly.

Accuracy vs. Precision

Accuracy How close a measurement is to the correct or accepted value.

Precision How close repeated measurements are to each other, regardless of their accuracy.

An instrument can be precise without being accurate, or accurate without being precise. Our goal is to be both accurate and precise.

Concept Visualization: Accuracy & Precision

To understand the difference between accuracy and precision, we can use the analogy of a dartboard.

- **Accuracy** is hitting the bullseye.
- **Precision** is having all your darts land close together.

On the next slide, we will see four different scenarios.

Visualizing Accuracy & Precision




Figure: Four targets showing different combinations of accuracy and precision

Distance vs. Displacement

- **Distance** is a **scalar** quantity. It is the total length of the path traveled.

Distance vs. Displacement

- **Distance** is a **scalar** quantity. It is the total length of the path traveled.
 - Example: If you walk 5 meters east and 5 meters west, your distance traveled is 10 m.

Distance vs. Displacement

- **Distance** is a **scalar** quantity. It is the total length of the path traveled.
 - Example: If you walk 5 meters east and 5 meters west, your distance traveled is 10 m.
- **Displacement** ($\Delta \vec{d}$) is a **vector** quantity. It is the change in position from the start point to the end point.

Distance vs. Displacement

- **Distance** is a **scalar** quantity. It is the total length of the path traveled.
 - Example: If you walk 5 meters east and 5 meters west, your distance traveled is 10 m.
- **Displacement** ($\Delta\vec{d}$) is a **vector** quantity. It is the change in position from the start point to the end point.
 - $\Delta\vec{d} = \vec{d}_{final} - \vec{d}_{initial}$

Distance vs. Displacement

- **Distance** is a **scalar** quantity. It is the total length of the path traveled.
 - Example: If you walk 5 meters east and 5 meters west, your distance traveled is 10 m.
- **Displacement** ($\Delta \vec{d}$) is a **vector** quantity. It is the change in position from the start point to the end point.
 - $\Delta \vec{d} = \vec{d}_{final} - \vec{d}_{initial}$
 - In the example above, your displacement is 0 m because you ended where you started.

Concept Visualization: Distance vs. Displacement

Imagine a person walking from their house to a store. The path they take might involve several turns.

- The **distance** is the full length of the winding path they walked.
- The **displacement** is the straight-line distance and direction from the house to the store.

The next slide shows a map illustrating this.

Visualizing Distance vs. Displacement




Figure: Map showing winding path (distance) vs. straight-line path (displacement)

Speed vs. Velocity

- **Average Speed** is a **scalar** quantity that describes the rate of covering distance.

Speed vs. Velocity

- **Average Speed** is a **scalar** quantity that describes the rate of covering distance.
 - $v_{avg} = \frac{\text{distance}}{\text{time}}$

Speed vs. Velocity

- **Average Speed** is a **scalar** quantity that describes the rate of covering distance.
 - $v_{avg} = \frac{\text{distance}}{\text{time}}$
- **Velocity** is a **vector** quantity that describes the rate of displacement. It includes speed **and direction**.

Speed vs. Velocity

- **Average Speed** is a **scalar** quantity that describes the rate of covering distance.
 - $v_{avg} = \frac{\text{distance}}{\text{time}}$
- **Velocity** is a **vector** quantity that describes the rate of displacement. It includes speed **and direction**.
 - $\vec{v}_{avg} = \frac{\text{displacement}}{\text{time}} = \frac{\Delta \vec{d}}{\Delta t}$

Speed vs. Velocity

- **Average Speed** is a **scalar** quantity that describes the rate of covering distance.
 - $v_{avg} = \frac{\text{distance}}{\text{time}}$
- **Velocity** is a **vector** quantity that describes the rate of displacement. It includes speed **and direction**.
 - $\vec{v}_{avg} = \frac{\text{displacement}}{\text{time}} = \frac{\Delta \vec{d}}{\Delta t}$
- **Instantaneous velocity** is the velocity at a specific moment in time.

Position vs. Time Graphs

Graphs are a powerful way to visualize and analyze motion.

Position vs. Time Graphs

Graphs are a powerful way to visualize and analyze motion.

- On a position-time (p-t) graph, the **y-axis** is position and the **x-axis** is time.

Position vs. Time Graphs

Graphs are a powerful way to visualize and analyze motion.

- On a position-time (p-t) graph, the **y-axis** is position and the **x-axis** is time.
- The **slope** of the line on a p-t graph represents **velocity**.

Position vs. Time Graphs

Graphs are a powerful way to visualize and analyze motion.

- On a position-time (p-t) graph, the **y-axis** is position and the **x-axis** is time.
- The **slope** of the line on a p-t graph represents **velocity**.
 - $\text{Slope} = \frac{\text{rise}}{\text{run}} = \frac{\Delta \text{position}}{\Delta \text{time}} = \text{velocity}$

Position vs. Time Graphs

Graphs are a powerful way to visualize and analyze motion.

- On a position-time (p-t) graph, the **y-axis** is position and the **x-axis** is time.
- The **slope** of the line on a p-t graph represents **velocity**.
 - $\text{Slope} = \frac{\text{rise}}{\text{run}} = \frac{\Delta \text{position}}{\Delta \text{time}} = \text{velocity}$
- A **straight line** means constant velocity.

Position vs. Time Graphs

Graphs are a powerful way to visualize and analyze motion.

- On a position-time (p-t) graph, the **y-axis** is position and the **x-axis** is time.
- The **slope** of the line on a p-t graph represents **velocity**.
 - $\text{Slope} = \frac{\text{rise}}{\text{run}} = \frac{\Delta \text{position}}{\Delta \text{time}} = \text{velocity}$
- A **straight line** means constant velocity.
- A **curved line** means changing velocity (acceleration).

Position vs. Time Graphs


Graphs are a powerful way to visualize and analyze motion.

- On a position-time (p-t) graph, the **y-axis** is position and the **x-axis** is time.
- The **slope** of the line on a p-t graph represents **velocity**.
 - $\text{Slope} = \frac{\text{rise}}{\text{run}} = \frac{\Delta \text{position}}{\Delta \text{time}} = \text{velocity}$
- A **straight line** means constant velocity.
- A **curved line** means changing velocity (acceleration).
- **Key Idea:** The steeper the slope, the faster the object is moving.

Visualizing Constant Velocity (p-t Graph)

Let's visualize an object moving at a constant, positive velocity. The p-t graph will be a straight line with a positive slope. The value of that slope is the object's velocity.

Position vs. Time Graph (Constant Velocity)



Velocity vs. Time Graphs

We can also graph velocity against time.

Velocity vs. Time Graphs

We can also graph velocity against time.

- On a velocity-time (v - t) graph, the **y-axis** is velocity and the **x-axis** is time.

Velocity vs. Time Graphs

We can also graph velocity against time.

- On a velocity-time (v-t) graph, the **y-axis** is velocity and the **x-axis** is time.
- The **slope** of the line on a v-t graph represents **acceleration**.

Velocity vs. Time Graphs

We can also graph velocity against time.

- On a velocity-time (v-t) graph, the **y-axis** is velocity and the **x-axis** is time.
- The **slope** of the line on a v-t graph represents **acceleration**.
 - $\text{Slope} = \frac{\Delta \text{velocity}}{\Delta \text{time}} = \text{acceleration}$

Velocity vs. Time Graphs

We can also graph velocity against time.

- On a velocity-time (v-t) graph, the **y-axis** is velocity and the **x-axis** is time.
- The **slope** of the line on a v-t graph represents **acceleration**.
 - $\text{Slope} = \frac{\Delta \text{velocity}}{\Delta \text{time}} = \text{acceleration}$
- The **area under the curve** of a v-t graph represents **displacement**.

Velocity vs. Time Graphs

We can also graph velocity against time.

- On a velocity-time (v-t) graph, the **y-axis** is velocity and the **x-axis** is time.
- The **slope** of the line on a v-t graph represents **acceleration**.
 - $\text{Slope} = \frac{\Delta \text{velocity}}{\Delta \text{time}} = \text{acceleration}$
- The **area under the curve** of a v-t graph represents **displacement**.
 - $\text{Area} = \text{velocity} \times \text{time} = \text{displacement}$

Velocity vs. Time Graphs

We can also graph velocity against time.


- On a velocity-time (v-t) graph, the **y-axis** is velocity and the **x-axis** is time.
- The **slope** of the line on a v-t graph represents **acceleration**.
 - $\text{Slope} = \frac{\Delta \text{velocity}}{\Delta \text{time}} = \text{acceleration}$
- The **area under the curve** of a v-t graph represents **displacement**.
 - $\text{Area} = \text{velocity} \times \text{time} = \text{displacement}$
- **Key Ideas:** Slope gives acceleration, area gives displacement.

Visualizing Constant Acceleration (v-t Graph)

Let's visualize an object moving with constant, positive acceleration. The v-t graph will be a straight line with a positive slope.

- The **slope** of this line is the acceleration.
- The **area** under this line is the total displacement.

Velocity vs. Time Graph (Constant Acceleration)



Connecting v-t and a-t Graphs

When an object has constant acceleration, we can show this on two different graphs:

Connecting v-t and a-t Graphs

When an object has constant acceleration, we can show this on two different graphs:

Velocity-Time Graph:

- Shows a straight line
- Slope = acceleration
- Example: slope = 2 m/s^2

Acceleration-Time Graph:

- Shows a horizontal line
- Height = acceleration value
- Example: $a = 2 \text{ m/s}^2$ (constant)

Connecting v-t and a-t Graphs

When an object has constant acceleration, we can show this on two different graphs:

Velocity-Time Graph:

- Shows a straight line
- Slope = acceleration
- Example: slope = 2 m/s^2

Acceleration-Time Graph:

- Shows a horizontal line
- Height = acceleration value
- Example: $a = 2 \text{ m/s}^2$ (constant)

Key Connection

For constant acceleration: v-t graph is a straight line, a-t graph is a horizontal line.

Acceleration

- **Acceleration** is the rate at which an object's velocity changes.

Acceleration

- **Acceleration** is the rate at which an object's velocity changes.
- Acceleration is a **vector** quantity. This means it has both size (magnitude) and direction.

Acceleration

- **Acceleration** is the rate at which an object's velocity changes.
- Acceleration is a **vector** quantity. This means it has both size (magnitude) and direction.
- We measure acceleration in meters per second squared (m/s^2).

Acceleration

- **Acceleration** is the rate at which an object's velocity changes.
- Acceleration is a **vector** quantity. This means it has both size (magnitude) and direction.
- We measure acceleration in meters per second squared (m/s^2).
- The formula for average acceleration is:

$$\vec{a} = \frac{\Delta \vec{v}}{\Delta t} = \frac{\vec{v}_f - \vec{v}_0}{t_f - t_0}$$

Acceleration

- **Acceleration** is the rate at which an object's velocity changes.
- Acceleration is a **vector** quantity. This means it has both size (magnitude) and direction.
- We measure acceleration in meters per second squared (m/s^2).
- The formula for average acceleration is:

$$\vec{a} = \frac{\Delta \vec{v}}{\Delta t} = \frac{\vec{v}_f - \vec{v}_0}{t_f - t_0}$$

- This means: acceleration = change in velocity \div time taken

Three Types of Acceleration

- An object is accelerating when it does any of these three things:

Three Types of Acceleration

- An object is accelerating when it does any of these three things:
- **1. Speeding up** - velocity increases in the same direction

Three Types of Acceleration

- An object is accelerating when it does any of these three things:
- **1. Speeding up** - velocity increases in the same direction
- **2. Slowing down** - velocity decreases

Three Types of Acceleration

- An object is accelerating when it does any of these three things:
- **1. Speeding up** - velocity increases in the same direction
- **2. Slowing down** - velocity decreases
 - This is sometimes called **deceleration**
 - But it's still acceleration - just in the opposite direction of motion
 - Example: A car going 20 m/s that slows to 15 m/s has acceleration = $-5 \text{ m/s} \div \text{time}$

Three Types of Acceleration

- An object is accelerating when it does any of these three things:
- **1. Speeding up** - velocity increases in the same direction
- **2. Slowing down** - velocity decreases
 - This is sometimes called **deceleration**
 - But it's still acceleration - just in the opposite direction of motion
 - Example: A car going 20 m/s that slows to 15 m/s has acceleration = $-5 \text{ m/s} \div \text{time}$
- **3. Changing direction** - even if speed stays the same

Three Types of Acceleration

- An object is accelerating when it does any of these three things:
- **1. Speeding up** - velocity increases in the same direction
- **2. Slowing down** - velocity decreases
 - This is sometimes called **deceleration**
 - But it's still acceleration - just in the opposite direction of motion
 - Example: A car going 20 m/s that slows to 15 m/s has acceleration = $-5 \text{ m/s} \div \text{time}$
- **3. Changing direction** - even if speed stays the same
- **Key Point:** Any change in velocity (speed OR direction) means acceleration is happening.

Essential Equations: The Kinematics

These equations describe motion with **constant acceleration**.

Essential Equations: The Kinematics

These equations describe motion with **constant acceleration**.

$$① \quad \vec{v} = \vec{v}_0 + \vec{a}t$$

Essential Equations: The Kinematics

These equations describe motion with **constant acceleration**.

$$\textcircled{1} \quad \vec{v} = \vec{v}_0 + \vec{a}t$$

$$\textcircled{2} \quad \Delta \vec{d} = \vec{v}_0 t + \frac{1}{2} \vec{a} t^2$$

Essential Equations: The Kinematics

These equations describe motion with **constant acceleration**.

$$\textcircled{1} \quad \vec{v} = \vec{v}_0 + \vec{a}t$$

$$\textcircled{2} \quad \Delta \vec{d} = \vec{v}_0 t + \frac{1}{2} \vec{a} t^2$$

$$\textcircled{3} \quad \vec{v}^2 = \vec{v}_0^2 + 2\vec{a}\Delta \vec{d}$$

Essential Equations: The Kinematics

These equations describe motion with **constant acceleration**.

$$\textcircled{1} \quad \vec{v} = \vec{v}_0 + \vec{a}t$$

$$\textcircled{2} \quad \Delta \vec{d} = \vec{v}_0 t + \frac{1}{2} \vec{a} t^2$$

$$\textcircled{3} \quad \vec{v}^2 = \vec{v}_0^2 + 2\vec{a}\Delta \vec{d}$$

$$\textcircled{4} \quad \Delta \vec{d} = \frac{\vec{v}_0 + \vec{v}}{2} t$$

Essential Equations: The Kinematics

These equations describe motion with **constant acceleration**.

Variables:

$\Delta \vec{d}$: displacement (m)

t : time (s)

\vec{a} : acceleration (m/s^2)

\vec{v}_0 : initial velocity (m/s)

\vec{v} : final velocity (m/s)

① $\vec{v} = \vec{v}_0 + \vec{a}t$

② $\Delta \vec{d} = \vec{v}_0 t + \frac{1}{2} \vec{a} t^2$

③ $\vec{v}^2 = \vec{v}_0^2 + 2\vec{a}\Delta \vec{d}$

④ $\Delta \vec{d} = \frac{\vec{v}_0 + \vec{v}}{2} t$

Essential Equations: The Kinematics

These equations describe motion with **constant acceleration**.

Variables:

$\Delta \vec{d}$: displacement (m)

t : time (s)

\vec{a} : acceleration (m/s^2)

\vec{v}_0 : initial velocity (m/s)

\vec{v} : final velocity (m/s)

① $\vec{v} = \vec{v}_0 + \vec{a}t$

② $\Delta \vec{d} = \vec{v}_0 t + \frac{1}{2} \vec{a} t^2$

③ $\vec{v}^2 = \vec{v}_0^2 + 2\vec{a}\Delta \vec{d}$

④ $\Delta \vec{d} = \frac{\vec{v}_0 + \vec{v}}{2} t$

Note: We often write these without vector arrows in 1D problems, using $+/-$ signs for direction.

The GUESS Method

To solve physics problems consistently, we will use the **GUESS** method.

The GUESS Method

To solve physics problems consistently, we will use the **GUESS** method.

G - **Givens** List all known quantities from the problem, with variable symbols and units.

The GUESS Method

To solve physics problems consistently, we will use the **GUESS** method.

G - Givens List all known quantities from the problem, with variable symbols and units.

U - Unknown Identify the quantity you need to find.

The GUESS Method

To solve physics problems consistently, we will use the **GUESS** method.

G - Givens List all known quantities from the problem, with variable symbols and units.

U - Unknown Identify the quantity you need to find.

E - Equation Select the physics equation that relates your givens and unknown.

The GUESS Method

To solve physics problems consistently, we will use the **GUESS** method.

- G - Givens** List all known quantities from the problem, with variable symbols and units.
- U - Unknown** Identify the quantity you need to find.
- E - Equation** Select the physics equation that relates your givens and unknown.
- S - Substitute** Plug the known values (with units) into the equation.

The GUESS Method

To solve physics problems consistently, we will use the **GUESS** method.

- G - Givens** List all known quantities from the problem, with variable symbols and units.
- U - Unknown** Identify the quantity you need to find.
- E - Equation** Select the physics equation that relates your givens and unknown.
- S - Substitute** Plug the known values (with units) into the equation.
- S - Solve** Calculate the result, ensuring correct units and significant figures.

The GUESS Method

To solve physics problems consistently, we will use the **GUESS** method.

- G - Givens** List all known quantities from the problem, with variable symbols and units.
- U - Unknown** Identify the quantity you need to find.
- E - Equation** Select the physics equation that relates your givens and unknown.
- S - Substitute** Plug the known values (with units) into the equation.
- S - Solve** Calculate the result, ensuring correct units and significant figures.

This structured approach helps prevent mistakes and makes your work easy to follow.

I Do: Calculating Average Acceleration

Problem (12, p. 6)

The driver of a sports car traveling at 10.0 m/s steps down hard on the accelerator for 5.0 s and the velocity increases to 30.0 m/s . What was the average acceleration of the car during the 5.0-s time interval?

I Do: Calculating Average Acceleration - Solution Part 1

Solution using GUESS Method

- G - Givens:**
- Initial velocity (v_i) = 10.0 m/s
 - Final velocity (v_f) = 30.0 m/s
 - Time interval (t) = 5.0 s

U - Unknown: • Average acceleration (a) = ?

E - Equation:

$$a = \frac{\Delta v}{t} = \frac{v_f - v_i}{t}$$

I Do: Calculating Average Acceleration - Solution Part 2

Solution using GUESS Method (continued)

S - Substitute:

$$a = \frac{30.0 \text{ m/s} - 10.0 \text{ m/s}}{5.0 \text{ s}}$$

S - Solve:

$$a = \frac{20.0 \text{ m/s}}{5.0 \text{ s}}$$

$$a = 4.0 \text{ m/s}^2$$

We Do: Finding Initial Speed

Problem (8, p. 4)

A motorcycle moving at a constant velocity suddenly accelerates at a rate of 4.0 m/s^2 to a speed of 35 m/s in 5.0 s . What was the initial speed of the motorcycle?

We Do: Finding Initial Speed - Setup

Let's use the GUESS Method

- G - Givens:**
- Acceleration (a) = 4.0 m/s^2
 - Final velocity (v_f) = 35 m/s
 - Time interval (t) = 5.0 s

- U - Unknown:**
- Initial velocity (v_i) = ?

We Do: Finding Initial Speed - Equation

Let's use the GUESS Method

E - Equation:

$$v_f = v_i + at$$

We need to rearrange this equation to solve for v_i .

We Do: Finding Initial Speed - Equation

Let's use the GUESS Method

E - Equation:

$$v_f = v_i + at$$

We need to rearrange this equation to solve for v_i .

- Subtract at from both sides:
- $v_f - at = v_i$

We Do: Finding Initial Speed - Equation

Let's use the GUESS Method

E - Equation:

$$v_f = v_i + at$$

We need to rearrange this equation to solve for v_i .

- Subtract at from both sides:
- $v_f - at = v_i$

$$v_i = v_f - at$$

We Do: Finding Initial Speed - Solution

Let's use the GUESS Method

S - Substitute: Now we substitute our known values into the equation:

We Do: Finding Initial Speed - Solution

Let's use the GUESS Method

S - Substitute: Now we substitute our known values into the equation:

- $v_f = 35 \text{ m/s}$, $a = 4.0 \text{ m/s}^2$, $t = 5.0 \text{ s}$
$$v_i = 35 \text{ m/s} - (4.0 \text{ m/s}^2)(5.0 \text{ s})$$

S - Solve:

$$v_i = 35 \text{ m/s} - 20 \text{ m/s}$$

$$v_i = 15 \text{ m/s}$$

You Do: Practice Problem

Problem (24, p. 12)

How long does it take to accelerate from 8.0 m/s to 20.0 m/s at a rate of acceleration of 3.0 m/s^2 ?

Your Turn

Use the GUESS method to solve the problem.

- Identify your **Givens**.
- State the **Unknown**.
- Select the correct **Equation** and rearrange it.
- **Substitute** the values and **Solve**.

Lesson Summary

- Physics provides the fundamental rules for describing the universe.

Lesson Summary

- Physics provides the fundamental rules for describing the universe.
- Motion is described using **scalars** (distance, speed) and **vectors** (displacement, velocity, acceleration).

Lesson Summary

- Physics provides the fundamental rules for describing the universe.
- Motion is described using **scalars** (distance, speed) and **vectors** (displacement, velocity, acceleration).
- **Graphs** are a key tool for analyzing motion:

Lesson Summary

- Physics provides the fundamental rules for describing the universe.
- Motion is described using **scalars** (distance, speed) and **vectors** (displacement, velocity, acceleration).
- **Graphs** are a key tool for analyzing motion:
 - The slope of a position-time graph is **velocity**.

Lesson Summary

- Physics provides the fundamental rules for describing the universe.
- Motion is described using **scalars** (distance, speed) and **vectors** (displacement, velocity, acceleration).
- **Graphs** are a key tool for analyzing motion:
 - The slope of a position-time graph is **velocity**.
 - The slope of a velocity-time graph is **acceleration**.

Lesson Summary

- Physics provides the fundamental rules for describing the universe.
- Motion is described using **scalars** (distance, speed) and **vectors** (displacement, velocity, acceleration).
- **Graphs** are a key tool for analyzing motion:
 - The slope of a position-time graph is **velocity**.
 - The slope of a velocity-time graph is **acceleration**.
 - The area under a velocity-time graph is **displacement**.

Lesson Summary

- Physics provides the fundamental rules for describing the universe.
- Motion is described using **scalars** (distance, speed) and **vectors** (displacement, velocity, acceleration).
- **Graphs** are a key tool for analyzing motion:
 - The slope of a position-time graph is **velocity**.
 - The slope of a velocity-time graph is **acceleration**.
 - The area under a velocity-time graph is **displacement**.
- For motion with **constant acceleration**, we use the kinematic equations to relate displacement, velocity, acceleration, and time.