

PHYS12 CH:10 When the Universe Gets Weird

Einstein's Revolution

Mr. Gullo

December 2025

Outline

- 1 Introduction
- 2 Postulates of Special Relativity
- 3 Consequences of Special Relativity
- 4 Summary

Have you ever dreamed of traveling
to other star systems?

Have you ever dreamed of traveling
to other star systems?

Just fly fast enough, right?

Have you ever dreamed of traveling
to other star systems?

Just fly fast enough, right?

Wrong. Physics has other plans.

Distant Worlds




Figure: The Orion Nebula - home to distant star systems




Figure: The Orion Nebula - home to distant star systems

The Barrier

Special relativity explains why we can't reach these stars with current technology.

Before Einstein

Newton and Galileo were right... mostly.

Before Einstein

Newton and Galileo were right... mostly.

Classical relativity worked for centuries:

- Motion is relative to your frame of reference

Newton and Galileo were right... mostly.

Classical relativity worked for centuries:

- Motion is relative to your frame of reference
- Velocities add together

Newton and Galileo were right... mostly.

Classical relativity worked for centuries:

- Motion is relative to your frame of reference
- Velocities add together
- Time flows the same for everyone

Newton and Galileo were right... mostly.

Classical relativity worked for centuries:

- Motion is relative to your frame of reference
- Velocities add together
- Time flows the same for everyone

But at extreme speeds, everything breaks down.

Learning Objectives

By the end of this section, you will be able to:

- **10.1:** Describe the experiments that led Einstein to special relativity

Learning Objectives

By the end of this section, you will be able to:

- **10.1:** Describe the experiments that led Einstein to special relativity
- **10.1:** Understand the two postulates on which the theory is based

Learning Objectives

By the end of this section, you will be able to:

- **10.1:** Describe the experiments that led Einstein to special relativity
- **10.1:** Understand the two postulates on which the theory is based
- **10.1:** Explain why simultaneity depends on frame of reference

10.1 The Phantom Medium

19th century belief: Light must travel through a medium

10.1 The Phantom Medium

19th century belief: Light must travel through a medium

Like sound travels through air, light travels through... *ether*?

10.1 The Phantom Medium

19th century belief: Light must travel through a medium

Like sound travels through air, light travels through... *ether*?

The Mental Model

Scientists believed space was filled with an invisible fluid called the ether.

10.1 The Phantom Medium

19th century belief: Light must travel through a medium

Like sound travels through air, light travels through... *ether*?

The Mental Model

Scientists believed space was filled with an invisible fluid called the ether.

Problem: The ether doesn't exist.

10.1 The Most Famous Failed Experiment




Figure: Michelson-Morley interferometer (1887)

10.1 The Most Famous Failed Experiment




Figure: Michelson-Morley interferometer (1887)

Goal: Measure Earth's speed through the ether

10.1 The Most Famous Failed Experiment




Figure: Michelson-Morley interferometer (1887)

Goal: Measure Earth's speed through the ether

Result: No ether detected. Light speed is constant.

10.1 What They Expected

The swimmer analogy:

Two swimmers leave a moving platform:

- One swims with and against the current

10.1 What They Expected

The swimmer analogy:

Two swimmers leave a moving platform:

- One swims with and against the current
- One swims perpendicular to the current

10.1 What They Expected

The swimmer analogy:

Two swimmers leave a moving platform:

- One swims with and against the current
- One swims perpendicular to the current
- Even at same speed, they arrive at different times

10.1 What They Expected

The swimmer analogy:

Two swimmers leave a moving platform:

- One swims with and against the current
- One swims perpendicular to the current
- Even at same speed, they arrive at different times

But light beams don't behave like swimmers.

10.1 Enter Einstein




Figure: Albert Einstein (1879-1955)

10.1 Enter Einstein




Figure: Albert Einstein (1879-1955)

1905: Einstein proposes special relativity

10.1 Enter Einstein




Figure: Albert Einstein (1879-1955)

1905: Einstein proposes special relativity
Based on two simple postulates...

10.1 The Two Postulates

Postulate 1: Universal Laws

The laws of physics are the same in all inertial reference frames.

10.1 The Two Postulates

Postulate 1: Universal Laws

The laws of physics are the same in all inertial reference frames.

Postulate 2: The Cosmic Speed Limit

$$c = 3.00 \times 10^8 \text{ m/s}$$

The speed of light is the same in all inertial frames and is NOT affected by the speed of its source.

10.1 The Speed of Light

$$c = 299,792,458 \text{ m/s}$$

10.1 The Speed of Light

$$c = 299,792,458 \text{ m/s}$$

For most purposes: $c \approx 3.00 \times 10^8 \text{ m/s}$

10.1 The Speed of Light

$$c = 299,792,458 \text{ m/s}$$

For most purposes: $c \approx 3.00 \times 10^8 \text{ m/s}$

Civilian View vs. Reality

Civilian: "Light from a speeding car goes faster."

Physicist: "Light always travels at c , regardless of source speed."

10.1 Inertial Reference Frame

Definition

A reference frame where objects follow Newton's First Law: Objects at rest stay at rest, objects in motion stay in motion at constant velocity, unless acted upon by external force.

10.1 Inertial Reference Frame

Definition

A reference frame where objects follow Newton's First Law: Objects at rest stay at rest, objects in motion stay in motion at constant velocity, unless acted upon by external force.

Examples:

- Inside a car moving at constant velocity

10.1 Inertial Reference Frame

Definition

A reference frame where objects follow Newton's First Law: Objects at rest stay at rest, objects in motion stay in motion at constant velocity, unless acted upon by external force.

Examples:

- Inside a car moving at constant velocity
- Inside a stationary house

10.1 Inertial Reference Frame

Definition

A reference frame where objects follow Newton's First Law: Objects at rest stay at rest, objects in motion stay in motion at constant velocity, unless acted upon by external force.

Examples:

- Inside a car moving at constant velocity
- Inside a stationary house
- Inside a spacecraft coasting through space

10.1 The Paradox of Velocities

Newtonian mechanics: Velocities add

10.1 The Paradox of Velocities

Newtonian mechanics: Velocities add

The Mental Model

You run at 3 m/s and throw a ball forward at 10 m/s.

Ball speed: $3 + 10 = 13$ m/s

10.1 The Paradox of Velocities

Newtonian mechanics: Velocities add

The Mental Model

You run at 3 m/s and throw a ball forward at 10 m/s.

Ball speed: $3 + 10 = 13$ m/s

But what about light?

10.1 The Paradox of Velocities

Newtonian mechanics: Velocities add

The Mental Model

You run at 3 m/s and throw a ball forward at 10 m/s.

Ball speed: $3 + 10 = 13$ m/s

But what about light?

The Illusion

Airliner traveling at 200 m/s emits light forward.

Your brain says: Light speed = $c + 200$ m/s

Reality: Light speed = c (always)

10.1 Simultaneity Is Relative




Figure: Two flash lamps on a moving train

10.1 Simultaneity Is Relative




Figure: Two flash lamps on a moving train

Observer A (on train): Flashes simultaneous

10.1 Simultaneity Is Relative




Figure: Two flash lamps on a moving train

Observer A (on train): Flashes simultaneous

Observer B (on platform): Flashes NOT simultaneous

10.1 Why Simultaneity Breaks

Key insight: $t = \frac{d}{v}$

10.1 Why Simultaneity Breaks

Key insight: $t = \frac{d}{v}$

If light speed c is constant for all observers...

10.1 Why Simultaneity Breaks

Key insight: $t = \frac{d}{v}$

If light speed c is constant for all observers...

...and distance d appears different to different observers...

10.1 Why Simultaneity Breaks

Key insight: $t = \frac{d}{v}$

If light speed c is constant for all observers...

...and distance d appears different to different observers...

...then time t must also be different!

10.1 Why Simultaneity Breaks

Key insight: $t = \frac{d}{v}$

If light speed c is constant for all observers...

...and distance d appears different to different observers...

...then time t must also be different!

The Universal Law

Two events are simultaneous only if an observer measures them as occurring at the same time. Two events are NOT necessarily simultaneous to all observers.

Attempt: Light Travel Time

The Challenge (3 min, silent)

The sun is 1.50×10^8 km from Earth. How long does it take light to travel from the sun to Earth?

Given:

- Distance $d = 1.50 \times 10^8$ km
- Speed of light $c = 3.00 \times 10^8$ m/s

Find: Time in seconds and minutes

Work silently. Convert units carefully.

Compare: Light Travel Time

Turn and talk (2 min):

- 1 What equation did you use?
- 2 How did you handle the units (km vs m)?
- 3 What did you get for your answer?

Compare: Light Travel Time

Turn and talk (2 min):

- 1 What equation did you use?
- 2 How did you handle the units (km vs m)?
- 3 What did you get for your answer?

Name wheel: One pair share your approach (not your answer).

Reveal: Light From the Sun

Self-correct in a different color:

Equation: $v = \frac{d}{t}$ so $t = \frac{d}{v}$

Reveal: Light From the Sun

Self-correct in a different color:

Equation: $v = \frac{d}{t}$ so $t = \frac{d}{v}$

Convert units: $1.50 \times 10^8 \text{ km} \times \frac{10^3 \text{ m}}{1 \text{ km}} = 1.50 \times 10^{11} \text{ m}$

Reveal: Light From the Sun

Self-correct in a different color:

Equation: $v = \frac{d}{t}$ so $t = \frac{d}{v}$

Convert units: $1.50 \times 10^8 \text{ km} \times \frac{10^3 \text{ m}}{1 \text{ km}} = 1.50 \times 10^{11} \text{ m}$

Substitute: $t = \frac{1.50 \times 10^{11} \text{ m}}{3.00 \times 10^8 \text{ m/s}}$

Reveal: Light From the Sun

Self-correct in a different color:

Equation: $v = \frac{d}{t}$ so $t = \frac{d}{v}$

Convert units: $1.50 \times 10^8 \text{ km} \times \frac{10^3 \text{ m}}{1 \text{ km}} = 1.50 \times 10^{11} \text{ m}$

Substitute: $t = \frac{1.50 \times 10^{11} \text{ m}}{3.00 \times 10^8 \text{ m/s}}$

$$t = 500 \text{ s} = 8 \text{ min } 20 \text{ s}$$

Reveal: Light From the Sun

Self-correct in a different color:

Equation: $v = \frac{d}{t}$ so $t = \frac{d}{v}$

Convert units: $1.50 \times 10^8 \text{ km} \times \frac{10^3 \text{ m}}{1 \text{ km}} = 1.50 \times 10^{11} \text{ m}$

Substitute: $t = \frac{1.50 \times 10^{11} \text{ m}}{3.00 \times 10^8 \text{ m/s}}$

$$t = 500 \text{ s} = 8 \text{ min } 20 \text{ s}$$

Check: Sunlight takes 8 minutes to reach Earth. When you see a sunspot, it happened 8 minutes ago!

Learning Objectives

By the end of this section, you will be able to:

- **10.2:** Describe time dilation, length contraction, and relativistic momentum

Learning Objectives

By the end of this section, you will be able to:

- **10.2:** Describe time dilation, length contraction, and relativistic momentum
- **10.2:** Explain mass-energy equivalence

Learning Objectives

By the end of this section, you will be able to:

- **10.2:** Describe time dilation, length contraction, and relativistic momentum
- **10.2:** Explain mass-energy equivalence
- **10.2:** Perform calculations involving relativistic effects

10.2 The Relativistic Factor

The Universal Factor

$$\gamma = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}}$$

γ (gamma) is the relativistic factor that appears in ALL relativistic effects.

10.2 The Relativistic Factor

The Universal Factor

$$\gamma = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}}$$

γ (gamma) is the relativistic factor that appears in ALL relativistic effects.

When $v \ll c$: $\gamma \approx 1$ (classical physics works)

10.2 The Relativistic Factor

The Universal Factor

$$\gamma = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}}$$

γ (gamma) is the relativistic factor that appears in ALL relativistic effects.

When $v \ll c$: $\gamma \approx 1$ (classical physics works)

When $v \approx c$: $\gamma \gg 1$ (relativistic effects dominate)

10.2 Testing the Relativistic Factor

Try some values for v :

10.2 Testing the Relativistic Factor

Try some values for v :

- Airliner: $v = 225 \text{ m/s} \rightarrow \gamma = 1.0000000003$

10.2 Testing the Relativistic Factor

Try some values for v :

- Airliner: $v = 225 \text{ m/s} \rightarrow \gamma = 1.0000000003$
- Earth's orbit: $v = 2.98 \times 10^4 \text{ m/s} \rightarrow \gamma = 1.0000000005$

10.2 Testing the Relativistic Factor

Try some values for v :

- Airliner: $v = 225 \text{ m/s} \rightarrow \gamma = 1.0000000003$
- Earth's orbit: $v = 2.98 \times 10^4 \text{ m/s} \rightarrow \gamma = 1.0000000005$
- Particle accelerator: $v = 2.99 \times 10^8 \text{ m/s} \rightarrow \gamma = 7.1$

10.2 Testing the Relativistic Factor

Try some values for v :

- Airliner: $v = 225 \text{ m/s} \rightarrow \gamma = 1.0000000003$
- Earth's orbit: $v = 2.98 \times 10^4 \text{ m/s} \rightarrow \gamma = 1.0000000005$
- Particle accelerator: $v = 2.99 \times 10^8 \text{ m/s} \rightarrow \gamma = 7.1$

Relativistic effects only matter near light speed!

10.2 Time Dilation

The Law of Time

$$\Delta t = \gamma \Delta t_0$$

Time passes MORE SLOWLY for an observer moving relative to you.

10.2 Time Dilation

The Law of Time

$$\Delta t = \gamma \Delta t_0$$

Time passes MORE SLOWLY for an observer moving relative to you.

- Δt_0 = proper time (measured by moving observer)

10.2 Time Dilation

The Law of Time

$$\Delta t = \gamma \Delta t_0$$

Time passes MORE SLOWLY for an observer moving relative to you.

- Δt_0 = proper time (measured by moving observer)
- Δt = dilated time (measured by stationary observer)

10.2 Time Dilation

The Law of Time

$$\Delta t = \gamma \Delta t_0$$

Time passes MORE SLOWLY for an observer moving relative to you.

- Δt_0 = proper time (measured by moving observer)
- Δt = dilated time (measured by stationary observer)
- $\Delta t > \Delta t_0$ always

10.2 The Astronaut's Clock




Figure: Light crossing a moving spacecraft

10.2 The Astronaut's Clock




Figure: Light crossing a moving spacecraft

Astronaut measures: Time Δt_0 (shorter path)

10.2 The Astronaut's Clock




Figure: Light crossing a moving spacecraft

Astronaut measures: Time Δt_0 (shorter path)

Earth observer measures: Time Δt (longer path)

10.2 The Astronaut's Clock




Figure: Light crossing a moving spacecraft

Astronaut measures: Time Δt_0 (shorter path)

Earth observer measures: Time Δt (longer path)

Same light, different distances, different times!

10.2 The Twin Paradox

Thought experiment:

Twin A travels to a distant star at near light speed

10.2 The Twin Paradox

Thought experiment:

Twin A travels to a distant star at near light speed

Twin B stays on Earth

10.2 The Twin Paradox

Thought experiment:

Twin A travels to a distant star at near light speed

Twin B stays on Earth

When Twin A returns...

10.2 The Twin Paradox

Thought experiment:

Twin A travels to a distant star at near light speed

Twin B stays on Earth

When Twin A returns...

Twin B has aged much more than Twin A!

10.2 The Twin Paradox

Thought experiment:

Twin A travels to a distant star at near light speed

Twin B stays on Earth

When Twin A returns...

Twin B has aged much more than Twin A!

Real-World Confirmation

Atomic clocks on GPS satellites run slower than Earth clocks. GPS must correct for time dilation to give accurate positioning.

10.2 Length Contraction

The Law of Length

$$L = \frac{L_0}{\gamma} = L_0 \sqrt{1 - \frac{v^2}{c^2}}$$

Objects appear SHORTER when moving relative to you.

10.2 Length Contraction

The Law of Length

$$L = \frac{L_0}{\gamma} = L_0 \sqrt{1 - \frac{v^2}{c^2}}$$

Objects appear SHORTER when moving relative to you.

- L_0 = proper length (measured at rest)

10.2 Length Contraction

The Law of Length

$$L = \frac{L_0}{\gamma} = L_0 \sqrt{1 - \frac{v^2}{c^2}}$$

Objects appear SHORTER when moving relative to you.

- L_0 = proper length (measured at rest)
- L = contracted length (measured by moving observer)

10.2 Length Contraction

The Law of Length

$$L = \frac{L_0}{\gamma} = L_0 \sqrt{1 - \frac{v^2}{c^2}}$$

Objects appear SHORTER when moving relative to you.

- L_0 = proper length (measured at rest)
- L = contracted length (measured by moving observer)
- $L < L_0$ always

10.2 The Road Ahead




Figure: The road ahead

10.2 The Road Ahead




Figure: The road ahead

At everyday speeds: You both measure the same distance

10.2 The Road Ahead




Figure: The road ahead

At everyday speeds: You both measure the same distance

At relativistic speeds: You measure different distances!

10.2 The Road Ahead




Figure: The road ahead

At everyday speeds: You both measure the same distance

At relativistic speeds: You measure different distances!

Because $v = \frac{d}{t}$ and you agree on v but not on t , you must also disagree on d !

Attempt: The Alien Spaceship

The Challenge (3 min, silent)

An alien spaceship is 50 m long and travels at 95% of the speed of light. What is the ship's length as measured from Earth?

Given:

- Proper length $L_0 = 50$ m
- Velocity $v = 0.95c$

Find: Contracted length L

Use the length contraction formula. Work silently.

Compare: Spaceship Length

Turn and talk (2 min):

- 1 What formula did you use?
- 2 Did you calculate γ first or use the combined formula?
- 3 How did you handle $v = 0.95c$?

Compare: Spaceship Length

Turn and talk (2 min):

- 1 What formula did you use?
- 2 Did you calculate γ first or use the combined formula?
- 3 How did you handle $v = 0.95c$?

Name wheel: One pair share your approach (not your answer).

Reveal: The Contracted Spaceship

Self-correct in a different color:

Equation: $L = L_0 \sqrt{1 - \frac{v^2}{c^2}}$

Reveal: The Contracted Spaceship

Self-correct in a different color:

Equation: $L = L_0 \sqrt{1 - \frac{v^2}{c^2}}$

Substitute: $L = 50 \text{ m} \sqrt{1 - \frac{(0.95c)^2}{c^2}}$

Reveal: The Contracted Spaceship

Self-correct in a different color:

Equation: $L = L_0 \sqrt{1 - \frac{v^2}{c^2}}$

Substitute: $L = 50 \text{ m} \sqrt{1 - \frac{(0.95c)^2}{c^2}}$

Simplify: $L = 50 \text{ m} \sqrt{1 - (0.95)^2}$

Reveal: The Contracted Spaceship

Self-correct in a different color:

Equation: $L = L_0 \sqrt{1 - \frac{v^2}{c^2}}$

Substitute: $L = 50 \text{ m} \sqrt{1 - \frac{(0.95c)^2}{c^2}}$

Simplify: $L = 50 \text{ m} \sqrt{1 - (0.95)^2}$

$$L = 50 \text{ m} \sqrt{1 - 0.9025} = 50 \text{ m} \sqrt{0.0975}$$

Reveal: The Contracted Spaceship

Self-correct in a different color:

Equation: $L = L_0 \sqrt{1 - \frac{v^2}{c^2}}$

Substitute: $L = 50 \text{ m} \sqrt{1 - \frac{(0.95c)^2}{c^2}}$

Simplify: $L = 50 \text{ m} \sqrt{1 - (0.95)^2}$

$$L = 50 \text{ m} \sqrt{1 - 0.9025} = 50 \text{ m} \sqrt{0.0975}$$

$$L = 16 \text{ m}$$

Reveal: The Contracted Spaceship

Self-correct in a different color:

Equation: $L = L_0 \sqrt{1 - \frac{v^2}{c^2}}$

Substitute: $L = 50 \text{ m} \sqrt{1 - \frac{(0.95c)^2}{c^2}}$

Simplify: $L = 50 \text{ m} \sqrt{1 - (0.95)^2}$

$$L = 50 \text{ m} \sqrt{1 - 0.9025} = 50 \text{ m} \sqrt{0.0975}$$

$$L = 16 \text{ m}$$

Check: Ship contracted from 50 m to 16 m - only 32% of original length!

10.2 Relativistic Momentum

The Law of Momentum

$$p = \gamma mu$$

Momentum increases without limit as velocity approaches c .

10.2 Relativistic Momentum

The Law of Momentum

$$p = \gamma mu$$

Momentum increases without limit as velocity approaches c .

- m = rest mass

10.2 Relativistic Momentum

The Law of Momentum

$$p = \gamma mu$$

Momentum increases without limit as velocity approaches c .

- m = rest mass
- u = velocity of object

10.2 Relativistic Momentum

The Law of Momentum

$$p = \gamma mu$$

Momentum increases without limit as velocity approaches c .

- m = rest mass
- u = velocity of object
- As $u \rightarrow c$, $\gamma \rightarrow \infty$, so $p \rightarrow \infty$

10.2 The Momentum Barrier




Figure: Relativistic momentum approaches infinity

10.2 The Momentum Barrier




Figure: Relativistic momentum approaches infinity

As $v \rightarrow c$, momentum $p \rightarrow \infty$

10.2 The Momentum Barrier




Figure: Relativistic momentum approaches infinity

As $v \rightarrow c$, momentum $p \rightarrow \infty$

This is why you can't reach the speed of light!

10.2 Mass-Energy Equivalence

The Source Code of Energy

$$E = mc^2$$

Mass and energy are interchangeable. Matter IS energy.

10.2 Mass-Energy Equivalence

The Source Code of Energy

$$E = mc^2$$

Mass and energy are interchangeable. Matter IS energy.

- E = rest energy (joules)

10.2 Mass-Energy Equivalence

The Source Code of Energy

$$E = mc^2$$

Mass and energy are interchangeable. Matter IS energy.

- E = rest energy (joules)
- m = rest mass (kg)

10.2 Mass-Energy Equivalence

The Source Code of Energy

$$E = mc^2$$

Mass and energy are interchangeable. Matter IS energy.

- E = rest energy (joules)
- m = rest mass (kg)
- c = speed of light (3.00×10^8 m/s)

10.2 The Power of c^2

How much energy in 1 gram of matter?

10.2 The Power of c^2

How much energy in 1 gram of matter?

$$E = mc^2 = (0.001 \text{ kg})(3.00 \times 10^8 \text{ m/s})^2$$

10.2 The Power of c^2

How much energy in 1 gram of matter?

$$E = mc^2 = (0.001 \text{ kg})(3.00 \times 10^8 \text{ m/s})^2$$

$$E = 9.0 \times 10^{13} \text{ J}$$

10.2 The Power of c^2

How much energy in 1 gram of matter?

$$E = mc^2 = (0.001 \text{ kg})(3.00 \times 10^8 \text{ m/s})^2$$

$$E = 9.0 \times 10^{13} \text{ J}$$

That's enough to power 750,000 homes for one hour!

10.2 The Power of c^2

How much energy in 1 gram of matter?

$$E = mc^2 = (0.001 \text{ kg})(3.00 \times 10^8 \text{ m/s})^2$$

$$E = 9.0 \times 10^{13} \text{ J}$$


That's enough to power 750,000 homes for one hour!

Comparison

Burning 1 gram of coal: 24 J

Converting 1 gram of mass to energy: $9.0 \times 10^{13} \text{ J}$

10.2 Where Mass Becomes Energy




(a)



(b)

The Sun

10.2 Where Mass Becomes Energy



(a)




(b)

The Sun



Nuclear power plant

10.2 Where Mass Becomes Energy



(a)



(b)

The Sun



Nuclear power plant

Both convert mass into energy through nuclear reactions.

10.2 Nuclear Binding Energy

Example: Helium nucleus

10.2 Nuclear Binding Energy

Example: Helium nucleus

Made of: 2 protons + 2 neutrons = 4.0330 u

10.2 Nuclear Binding Energy

Example: Helium nucleus

Made of: 2 protons + 2 neutrons = 4.0330 u

Actual mass: 4.0003 u

10.2 Nuclear Binding Energy

Example: Helium nucleus

Made of: 2 protons + 2 neutrons = 4.0330 u

Actual mass: 4.0003 u

Mass defect: 0.0327 u

10.2 Nuclear Binding Energy

Example: Helium nucleus

Made of: 2 protons + 2 neutrons = 4.0330 u

Actual mass: 4.0003 u

Mass defect: 0.0327 u

This "missing" mass became binding energy when the nucleus formed:

$$E = (5.04 \times 10^{-30} \text{ kg})(3.00 \times 10^8 \text{ m/s})^2 = 4.54 \times 10^{-12} \text{ J}$$

Attempt: Positron-Electron Annihilation

The Challenge (3 min, silent)

When a positron and electron collide, they annihilate and convert completely to energy. How much energy is released?

Given:

- Both particles have rest mass $m = 9.11 \times 10^{-31}$ kg
- Total mass: $2 \times 9.11 \times 10^{-31}$ kg

Find: Energy E in joules

Use $E = mc^2$. Work silently.

Compare: Annihilation Energy

Turn and talk (2 min):

- 1 Did you account for both particles?
- 2 What value did you use for c ?
- 3 What units did you get for your answer?

Compare: Annihilation Energy

Turn and talk (2 min):

- 1 Did you account for both particles?
- 2 What value did you use for c ?
- 3 What units did you get for your answer?

Name wheel: One pair share your approach (not your answer).

Reveal: Total Annihilation

Self-correct in a different color:

Equation: $E = mc^2$

Reveal: Total Annihilation

Self-correct in a different color:

Equation: $E = mc^2$

Total mass: $m = 2(9.11 \times 10^{-31} \text{ kg}) = 1.822 \times 10^{-30} \text{ kg}$

Reveal: Total Annihilation

Self-correct in a different color:

Equation: $E = mc^2$

Total mass: $m = 2(9.11 \times 10^{-31} \text{ kg}) = 1.822 \times 10^{-30} \text{ kg}$

Substitute: $E = (1.822 \times 10^{-30} \text{ kg})(3.00 \times 10^8 \text{ m/s})^2$

Reveal: Total Annihilation

Self-correct in a different color:

Equation: $E = mc^2$

Total mass: $m = 2(9.11 \times 10^{-31} \text{ kg}) = 1.822 \times 10^{-30} \text{ kg}$

Substitute: $E = (1.822 \times 10^{-30} \text{ kg})(3.00 \times 10^8 \text{ m/s})^2$

$$E = (1.822 \times 10^{-30})(9.00 \times 10^{16})$$

Reveal: Total Annihilation

Self-correct in a different color:

Equation: $E = mc^2$

Total mass: $m = 2(9.11 \times 10^{-31} \text{ kg}) = 1.822 \times 10^{-30} \text{ kg}$

Substitute: $E = (1.822 \times 10^{-30} \text{ kg})(3.00 \times 10^8 \text{ m/s})^2$

$$E = (1.822 \times 10^{-30})(9.00 \times 10^{16})$$

$$E = 1.64 \times 10^{-13} \text{ J}$$

Reveal: Total Annihilation

Self-correct in a different color:

Equation: $E = mc^2$

Total mass: $m = 2(9.11 \times 10^{-31} \text{ kg}) = 1.822 \times 10^{-30} \text{ kg}$

Substitute: $E = (1.822 \times 10^{-30} \text{ kg})(3.00 \times 10^8 \text{ m/s})^2$

$$E = (1.822 \times 10^{-30})(9.00 \times 10^{16})$$

$E = 1.64 \times 10^{-13} \text{ J}$

Check: Tiny particles, but enormous energy density. This becomes gamma rays!

10.2 The RHIC Collider



Figure: Brookhaven National Laboratory RHIC

10.2 The RHIC Collider



Figure: Brookhaven National Laboratory RHIC

Speed: 99.7% of light speed $\rightarrow \gamma = 12.9$

10.2 The RHIC Collider



Figure: Brookhaven National Laboratory RHIC

Speed: 99.7% of light speed $\rightarrow \gamma = 12.9$

Result: Time dilates by factor of 13, length contracts by factor of 13

10.2 The RHIC Collider



Figure: Brookhaven National Laboratory RHIC

Speed: 99.7% of light speed $\rightarrow \gamma = 12.9$

Result: Time dilates by factor of 13, length contracts by factor of 13

Goal: Recreate conditions from the Big Bang!

10.2 Summary of Relativistic Effects

The Three Laws

Time Dilation: $\Delta t = \gamma \Delta t_0$

Length Contraction: $L = \frac{L_0}{\gamma}$

Mass-Energy: $E = mc^2$

10.2 Summary of Relativistic Effects

The Three Laws

Time Dilation: $\Delta t = \gamma \Delta t_0$

Length Contraction: $L = \frac{L_0}{\gamma}$

Mass-Energy: $E = mc^2$

Where: $\gamma = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}}$

10.2 Summary of Relativistic Effects

The Three Laws

Time Dilation: $\Delta t = \gamma \Delta t_0$

Length Contraction: $L = \frac{L_0}{\gamma}$

Mass-Energy: $E = mc^2$

Where: $\gamma = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}}$

All controlled by the relativistic factor γ !

What You Now Know

The Revelations

- 1 The ether doesn't exist - light speed is constant

What You Now Know

The Revelations

- 1 The ether doesn't exist - light speed is constant
- 2 Two postulates: physics is universal, c is constant

What You Now Know

The Revelations

- 1 The ether doesn't exist - light speed is constant
- 2 Two postulates: physics is universal, c is constant
- 3 Simultaneity is relative - depends on frame of reference

What You Now Know

The Revelations

- 1 The ether doesn't exist - light speed is constant
- 2 Two postulates: physics is universal, c is constant
- 3 Simultaneity is relative - depends on frame of reference
- 4 Time dilates: moving clocks run slow

What You Now Know

The Revelations

- 1 The ether doesn't exist - light speed is constant
- 2 Two postulates: physics is universal, c is constant
- 3 Simultaneity is relative - depends on frame of reference
- 4 Time dilates: moving clocks run slow
- 5 Length contracts: moving objects shrink

What You Now Know

The Revelations

- 1 The ether doesn't exist - light speed is constant
- 2 Two postulates: physics is universal, c is constant
- 3 Simultaneity is relative - depends on frame of reference
- 4 Time dilates: moving clocks run slow
- 5 Length contracts: moving objects shrink
- 6 $E = mc^2$: mass and energy are equivalent

Key Equations

$$\gamma = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}} \quad (1)$$

$$\Delta t = \gamma \Delta t_0 \quad (2)$$

$$L = \frac{L_0}{\gamma} = L_0 \sqrt{1 - \frac{v^2}{c^2}} \quad (3)$$

$$p = \gamma m u \quad (4)$$

$$E = mc^2 \quad (5)$$

$$c = 3.00 \times 10^8 \text{ m/s} \quad (6)$$

Why We Can't Reach the Stars

The barrier:

As $v \rightarrow c$:

• $\gamma \rightarrow \infty$

Why We Can't Reach the Stars

The barrier:

As $v \rightarrow c$:

- $\gamma \rightarrow \infty$
- Momentum $\rightarrow \infty$

Why We Can't Reach the Stars

The barrier:

As $v \rightarrow c$:

- $\gamma \rightarrow \infty$
- Momentum $\rightarrow \infty$
- Energy required $\rightarrow \infty$

Why We Can't Reach the Stars

The barrier:

As $v \rightarrow c$:

- $\gamma \rightarrow \infty$
- Momentum $\rightarrow \infty$
- Energy required $\rightarrow \infty$

You can never reach light speed. The universe has a speed limit.

Why We Can't Reach the Stars

The barrier:

As $v \rightarrow c$:

- $\gamma \rightarrow \infty$
- Momentum $\rightarrow \infty$
- Energy required $\rightarrow \infty$

You can never reach light speed. The universe has a speed limit.

The Mental Model

The faster you go, the more energy you need. At light speed, you'd need infinite energy. Impossible.

Complete the assigned problems
posted on the LMS