PHYS11 CH123:

Test Prep

Mr. Gullo

October 2024

Mr. Gullo Unit 1 Review Oct 2024

Mr. Gullo Unit 1 Review Oct 2024 2/1

Introduction

- This presentation covers key concepts and problem-solving techniques for your upcoming physics test
- We'll review problems from three chapters:
 - Chapter 1: Unit Conversion and Estimation
 - Chapter 2: Frames of Reference and Displacement
 - Chapter 3: Kinematics and Motion
- Each problem will be presented with a step-by-step solution
- Focus on understanding the concepts and problem-solving approaches

Mr. Gullo Unit 1 Review Oct 2024 3/1

Mr. Gullo Unit 1 Review Oct 2024 4/1

Chapter 1: Problem 30

A generation is about one-third of a lifetime. Approximately how many generations have passed since the year 0 AD?

Mr. Gullo Unit 1 Review Oct 2024 5/1

Chapter 1: Problem 30 - Solution

Solution:

$$\mathsf{history} \cdot \frac{10^{11} \; \mathsf{s}}{\mathsf{history}} \cdot \frac{1 \; \mathsf{generation}}{1/3 \; \mathsf{lifetime}} \cdot \frac{0.5 \; \mathsf{lifetime}}{10^9 \; \mathsf{s}} = \underline{150 \; \mathsf{generations}}$$

Mr. Gullo Unit 1 Review Oct 2024

Chapter 1: Problem 30 - Explanation

Explanation:

- Start with the total time since 0 AD (history)
- 2 Convert this time to seconds: $\frac{10^{11} \text{ s}}{\text{history}}$
- $\begin{tabular}{ll} \hline \textbf{3} & Convert seconds to lifetimes: $\frac{0.5 \text{ lifetime}}{10^9 \text{ s}}$ \\ \hline \end{tabular}$
- **3** Convert lifetimes to generations: $\frac{1 \text{ generation}}{1/3 \text{ lifetime}}$
- Multiply all these factors together
- This simplifies to 150 generations

Key concept: Unit conversion and fraction multiplication

Mr. Gullo Unit 1 Review Oct 2024 7/1

Mr. Gullo Unit 1 Review Oct 2024 8/1

Chapter 2: Problem 28

Suppose a train is moving along a track. Is there a single, correct reference frame from which to describe the train's motion?

A. Yes, there is a single, correct frame of reference because motion is a relative term.

B. Yes, there is a single, correct frame of reference which is in terms of Earth's position.

C. No, there is not a single, correct frame of reference because motion is a relative term.

D. No, there is not a single, correct frame of reference because motion is independent of frame of reference.

Mr. Gullo Unit 1 Review Oct 2024 9/1

Chapter 2: Problem 28 - Solution

Solution: The correct answer is **C**. No, there is not a single correct frame of reference because motion is a relative term.

Screenshot 2024-10-11 141024.png



 Mr. Gullo
 Unit 1 Review
 Oct 2024
 10 / 1

Chapter 2: Problem 28 - Solution

Solution: The correct answer is **C**. No, there is not a single correct frame of reference because motion is a relative term. Explanation:

- Option C is correct: Motion depends on what you compare it to (relative motion).
- Option D is incorrect: It wrongly states that motion doesn't change with perspective.
- Examples:
 - On the train: Train seems still, landscape moves.
 - On the platform: Train moves, platform seems still.
 - In a car alongside: Train might seem to move slowly.
- This illustrates the principle of relativity in physics: motion description depends on the chosen reference frame.



Mr. Gullo Unit 1 Review Oct 2024 11/1

Chapter 2: Problem 28 - Explanation

Explanation:

- Motion is relative: The movement of an object can be described differently depending on the observer's frame of reference
- Multiple valid frames: For a moving train, we could describe its motion relative to the ground, a passenger inside the train, or even another moving object
- Each frame is correct: The motion described from each of these perspectives would be correct for that particular frame of reference
- No absolute frame: There isn't a single, universally "correct" frame of reference that takes precedence over others

Key concept: Relativity of motion and frames of reference

Oct 2024

Chapter 2: Problem 51

Calculate that object's net displacement over the time shown.

2024_09_22_d75bb9ada91612339d1ag-12.jpg

A. 540 m B. 2,520 m C. 2,790 m D. 5,040 m

Mr. Gullo Unit 1 Review Oct 2024 13/1

Chapter 2: Problem 51 - Solution

Solution: The correct answer is **C**. The net displacement is 2,790 m. Calculation:

- Area of rectangle: $18 \text{ m/s} \cdot 30 \text{ s} = 540 \text{ m}$
- Area of triangle: $\frac{1}{2} \cdot 150 \text{ m/s} \cdot 30 \text{ s} = 2,250 \text{ m}$
- Total area: 540 m + 2,250 m = 2,790 m

◆ロト ◆個ト ◆差ト ◆差ト を めへぐ

Mr. Gullo Unit 1 Review Oct 2024 14/1

Chapter 2: Problem 51 - Explanation

Explanation:

- Identify the shape: The area under the line consists of a rectangle and a triangle
- 2 Calculate the area of the rectangle: $18 \text{ m/s} \cdot 30 \text{ s} = 540 \text{ m}$
- **3** Calculate the area of the triangle: $\frac{1}{2} \cdot 150 \text{ m/s} \cdot 30 \text{ s} = 2,250 \text{ m}$
- **9** Sum the areas: 540 m + 2,250 m = 2,790 m

Key concept: In a velocity-time graph, the area under the curve represents displacement



Mr. Gullo Unit 1 Review Oct 2024 15/1

Mr. Gullo Unit 1 Review Oct 2024 16/1

Chapter 3: Problem 32

You throw a ball straight up with an initial velocity of 15.0 m/s. It passes a tree branch on the way up at a height of 7.00 m. How much additional time will pass before the ball passes the tree branch on the way back down? A. 0.574 s B. 0.956 s C. 1.53 s D. 1.91 s

Mr. Gullo Unit 1 Review Oct 2024 17/1

Chapter 3: Problem 32 - Solution Part 1

Solution: The correct answer is **D**. 1.91 s Step 1: Find the velocity at the branch

$$v^{2} = v_{0}^{2} + 2a(x - x_{0})$$

$$v = \sqrt{v_{0}^{2} + 2a(x - x_{0})}$$

$$v = \sqrt{(15.0 \text{ m/s})^{2} + 2(-9.8 \text{ m/s}^{2})(7 \text{ m} - 0 \text{ m})}$$

$$v = 9.37 \text{ m/s}$$

18/1

Mr. Gullo Unit 1 Review Oct 2024

Chapter 3: Problem 32 - Solution Part 2

Step 2: Find the time from the branch to the top

$$v = v_0 + at$$

$$t_1 = \frac{v - v_0}{a} = \frac{0 \text{ m/s} - 9.37 \text{ m/s}}{-9.8 \text{ m/s}^2} = 0.956 \text{ s}$$

Step 3: Double the time for total up and down

$$t = 2t_1 = 2(0.956 \text{ s}) = 1.91 \text{ s}$$

Mr. Gullo Unit 1 Review Oct 2024

Chapter 3: Problem 32 - Explanation

Explanation:

- Use $v^2 = v_0^2 + 2a(x x_0)$ to find velocity at the branch
- ② Use $v = v_0 + at$ to find time from branch to top
- Ouble this time for total time up and down

Key concepts:

- Kinematic equations for constant acceleration
- Symmetry of motion under constant acceleration
- Negative acceleration for upward motion

◆ロト ◆個ト ◆差ト ◆差ト 差 める(*)

Mr. Gullo Unit 1 Review Oct 2024 20 / 1

 Mr. Gullo
 Unit 1 Review
 Oct 2024
 21/1

Conclusion

- We've covered key problems from three important chapters:
 - Unit conversion and estimation
 - Frames of reference and displacement
 - Kinematics and motion
- Focus on understanding the concepts behind each problem
- Practice applying these problem-solving techniques
- Remember to identify given information, choose appropriate equations, and check units
- Good luck on your upcoming test!

Mr. Gullo Unit 1 Review Oct 2024 22 / 1