

# PHYS11 CH:11 The Hidden Energy

## Temperature, Heat, and Phase Change

Mr. Gullo

December 2025

# Outline

- 1 Introduction
- 2 Temperature and Thermal Energy
- 3 Heat, Specific Heat, and Heat Transfer
- 4 Phase Change and Latent Heat
- 5 Summary

# The Mystery of the Welder



# The Mystery of the Welder



How does thermal energy travel from the arc to your skin meters away?

# Learning Objectives

By the end of this section, you will be able to:

- **11.1:** Explain that temperature is a measure of internal kinetic energy

# Learning Objectives

By the end of this section, you will be able to:

- **11.1:** Explain that temperature is a measure of internal kinetic energy
- **11.1:** Interconvert temperatures between Celsius, Kelvin, and Fahrenheit scales

# 11.1 What Is Temperature?

## The Illusion

**You think:** Temperature measures heat

**Reality:** Temperature measures average kinetic energy of molecules

# 11.1 What Is Temperature?

## The Illusion

**You think:** Temperature measures heat

**Reality:** Temperature measures average kinetic energy of molecules

## The Source Code

Temperature = what we measure on a thermometer

Heat = transfer of energy due to temperature difference



# 11.1 What Is Temperature?

## The Illusion

**You think:** Temperature measures heat

**Reality:** Temperature measures average kinetic energy of molecules

## The Source Code

Temperature = what we measure on a thermometer

Heat = transfer of energy due to temperature difference

Heat and temperature are NOT the same thing.

# 11.1 Thermal Energy: The Invisible Motion

## The Mental Model

Atoms and molecules are constantly bouncing around in random directions.  
Faster motion = higher temperature.

# 11.1 Thermal Energy: The Invisible Motion

## The Mental Model

Atoms and molecules are constantly bouncing around in random directions.  
Faster motion = higher temperature.

**Thermal energy** = average kinetic energy of particles in a substance

# 11.1 Thermal Energy: The Invisible Motion

## The Mental Model

Atoms and molecules are constantly bouncing around in random directions.  
Faster motion = higher temperature.

**Thermal energy** = average kinetic energy of particles in a substance

$$KE = \frac{1}{2}mv^2$$

# 11.1 Thermal Energy: The Invisible Motion

## The Mental Model

Atoms and molecules are constantly bouncing around in random directions.  
Faster motion = higher temperature.

**Thermal energy** = average kinetic energy of particles in a substance

$$KE = \frac{1}{2}mv^2$$

Higher speed  $\rightarrow$  greater kinetic energy  $\rightarrow$  higher temperature

# 11.1 Temperature Scales




Figure: Fahrenheit, Celsius, and Kelvin scales compared

# 11.1 Temperature Scales




Figure: Fahrenheit, Celsius, and Kelvin scales compared

## Three scales, same physics:

- Celsius: water freezes at 0, boils at 100
- Fahrenheit: water freezes at 32, boils at 212
- Kelvin: absolute zero at 0, water freezes at 273.15

# 11.1 Absolute Zero: The Coldest Possible

## The Ultimate Limit

**Absolute zero** =  $0\text{ K} = -273.15\text{ C}$

The temperature at which all molecular motion ceases.



# 11.1 Absolute Zero: The Coldest Possible

## The Ultimate Limit

**Absolute zero** =  $0\text{ K} = -273.15\text{ C}$

The temperature at which all molecular motion ceases.

## Can we reach it?

- Theoretically possible

# 11.1 Absolute Zero: The Coldest Possible

## The Ultimate Limit

**Absolute zero** =  $0\text{ K} = -273.15\text{ C}$

The temperature at which all molecular motion ceases.

## Can we reach it?

- Theoretically possible
- Never achieved in practice

# 11.1 Absolute Zero: The Coldest Possible

## The Ultimate Limit

**Absolute zero** = 0 K = -273.15 C

The temperature at which all molecular motion ceases.

## Can we reach it?

- Theoretically possible
- Never achieved in practice
- Coldest lab temperature:  $1.0 \times 10^{-10}$  K (Helsinki)

# 11.1 Absolute Zero: The Coldest Possible

## The Ultimate Limit

**Absolute zero** = 0 K = -273.15 C

The temperature at which all molecular motion ceases.

## Can we reach it?

- Theoretically possible
- Never achieved in practice
- Coldest lab temperature:  $1.0 \times 10^{-10}$  K (Helsinki)

At absolute zero, particles have minimum possible energy (quantum zero-point energy).

# 11.1 Converting Temperature Scales

## Key Conversion Equations

$$T_{\circ F} = \frac{9}{5} T_{\circ C} + 32$$

$$T_{\circ C} = \frac{5}{9} (T_{\circ F} - 32)$$

$$T_K = T_{\circ C} + 273.15$$

$$T_{\circ C} = T_K - 273.15$$

# 11.1 Converting Temperature Scales

## Key Conversion Equations

$$T_{\circ F} = \frac{9}{5} T_{\circ C} + 32$$

$$T_{\circ C} = \frac{5}{9} (T_{\circ F} - 32)$$

$$T_K = T_{\circ C} + 273.15$$

$$T_{\circ C} = T_K - 273.15$$

## Quick Check

Room temperature: 25 C = ? F = ? K

# Attempt: Temperature Conversion

## The Challenge (3 min, silent)

Your body temperature is normally 98.6 F.

### Given:

- $T_{oF} = 98.6 \text{ F}$

**Find:** Body temperature in Celsius and Kelvin

*Can you decode your body's thermal state? Work silently.*

# Compare: Temperature Conversion

**Turn and talk (2 min):**

- 1 Which equation did you use to convert F to C?
- 2 Did you subtract 32 before or after multiplying by  $5/9$ ?
- 3 How did you convert C to K?



# Compare: Temperature Conversion

## Turn and talk (2 min):

- 1 Which equation did you use to convert F to C?
- 2 Did you subtract 32 before or after multiplying by  $5/9$ ?
- 3 How did you convert C to K?

**Name wheel:** One pair share your approach (not your answer).

# Reveal: Decoding Body Temperature

**Self-correct in a different color:**

**Step 1:** Convert F to C using  $T_{\circ C} = \frac{5}{9}(T_{\circ F} - 32)$

# Reveal: Decoding Body Temperature

**Self-correct in a different color:**

**Step 1:** Convert F to C using  $T_{\circ C} = \frac{5}{9}(T_{\circ F} - 32)$

**Substitute:**  $T_{\circ C} = \frac{5}{9}(98.6 - 32) = \frac{5}{9}(66.6)$

# Reveal: Decoding Body Temperature

**Self-correct in a different color:**

**Step 1:** Convert F to C using  $T_{\circ C} = \frac{5}{9}(T_{\circ F} - 32)$

**Substitute:**  $T_{\circ C} = \frac{5}{9}(98.6 - 32) = \frac{5}{9}(66.6)$

$$T_{\circ C} = \boxed{37.0 \text{ C}}$$

# Reveal: Decoding Body Temperature

**Self-correct in a different color:**

**Step 1:** Convert F to C using  $T_{\circ C} = \frac{5}{9}(T_{\circ F} - 32)$

**Substitute:**  $T_{\circ C} = \frac{5}{9}(98.6 - 32) = \frac{5}{9}(66.6)$

$$T_{\circ C} = \boxed{37.0 \text{ C}}$$

**Step 2:** Convert C to K using  $T_K = T_{\circ C} + 273.15$

# Reveal: Decoding Body Temperature

**Self-correct in a different color:**

**Step 1:** Convert F to C using  $T_{\circ C} = \frac{5}{9}(T_{\circ F} - 32)$

**Substitute:**  $T_{\circ C} = \frac{5}{9}(98.6 - 32) = \frac{5}{9}(66.6)$

$$T_{\circ C} = \boxed{37.0 \text{ C}}$$

**Step 2:** Convert C to K using  $T_K = T_{\circ C} + 273.15$

$$T_K = 37.0 + 273.15 = \boxed{310.2 \text{ K}}$$

# Reveal: Decoding Body Temperature

**Self-correct in a different color:**

**Step 1:** Convert F to C using  $T_{\circ C} = \frac{5}{9}(T_{\circ F} - 32)$

**Substitute:**  $T_{\circ C} = \frac{5}{9}(98.6 - 32) = \frac{5}{9}(66.6)$

$$T_{\circ C} = \boxed{37.0 \text{ C}}$$

**Step 2:** Convert C to K using  $T_K = T_{\circ C} + 273.15$

$$T_K = 37.0 + 273.15 = \boxed{310.2 \text{ K}}$$

**Check:** 37 C is normal body temp. Makes sense!

# Learning Objectives

By the end of this section, you will be able to:

- **11.2:** Explain heat, heat capacity, and specific heat



# Learning Objectives

By the end of this section, you will be able to:

- **11.2:** Explain heat, heat capacity, and specific heat
- **11.2:** Distinguish between conduction, convection, and radiation

# Learning Objectives

By the end of this section, you will be able to:

- **11.2:** Explain heat, heat capacity, and specific heat
- **11.2:** Distinguish between conduction, convection, and radiation
- **11.2:** Solve problems involving specific heat and heat transfer

## 11.2 Heat: Energy on the Move

### The Universal Law

**Heat** = transfer of thermal energy from hot to cold

## 11.2 Heat: Energy on the Move

### The Universal Law

**Heat** = transfer of thermal energy from hot to cold

### Key insights:

- Heat is NOT a substance - it's energy in transit

## 11.2 Heat: Energy on the Move

### The Universal Law

**Heat** = transfer of thermal energy from hot to cold

#### Key insights:

- Heat is NOT a substance - it's energy in transit
- Heat always flows from high to low temperature

# 11.2 Heat: Energy on the Move

## The Universal Law

**Heat** = transfer of thermal energy from hot to cold

### Key insights:

- Heat is NOT a substance - it's energy in transit
- Heat always flows from high to low temperature
- Heat stops flowing when temperatures equalize

# 11.2 Heat: Energy on the Move

## The Universal Law

**Heat** = transfer of thermal energy from hot to cold

### Key insights:

- Heat is NOT a substance - it's energy in transit
- Heat always flows from high to low temperature
- Heat stops flowing when temperatures equalize
- Heat is measured in joules (J), like all energy

## 11.2 The Heat Equation

### Nature's Rule for Temperature Change

$$Q = mc\Delta T$$

Heat transferred = mass  $\times$  specific heat  $\times$  temperature change



## 11.2 The Heat Equation

### Nature's Rule for Temperature Change

$$Q = mc\Delta T$$

Heat transferred = mass  $\times$  specific heat  $\times$  temperature change

#### What each variable means:

- $Q$  = heat transferred (J)
- $m$  = mass (kg)
- $c$  = specific heat (J/kg·C)
- $\Delta T$  = temperature change (C or K)

## 11.2 Specific Heat: Material Stubbornness

### The Mental Model

Specific heat = how much energy needed to raise 1 kg of material by 1 C

## 11.2 Specific Heat: Material Stubbornness

### The Mental Model

Specific heat = how much energy needed to raise 1 kg of material by 1 C

#### Common values:

- Water: 4186 J/kg·C (very high!)

## 11.2 Specific Heat: Material Stubbornness

### The Mental Model

Specific heat = how much energy needed to raise 1 kg of material by 1 C

#### Common values:

- Water: 4186 J/kg·C (very high!)
- Aluminum: 900 J/kg·C

## 11.2 Specific Heat: Material Stubbornness

### The Mental Model

Specific heat = how much energy needed to raise 1 kg of material by 1 C

#### Common values:

- Water: 4186 J/kg·C (very high!)
- Aluminum: 900 J/kg·C
- Iron: 450 J/kg·C

## 11.2 Specific Heat: Material Stubbornness

### The Mental Model

Specific heat = how much energy needed to raise 1 kg of material by 1 C

#### Common values:

- Water: 4186 J/kg·C (very high!)
- Aluminum: 900 J/kg·C
- Iron: 450 J/kg·C
- Copper: 387 J/kg·C

## 11.2 Specific Heat: Material Stubbornness

### The Mental Model

Specific heat = how much energy needed to raise 1 kg of material by 1 C

#### Common values:

- Water: 4186 J/kg·C (very high!)
- Aluminum: 900 J/kg·C
- Iron: 450 J/kg·C
- Copper: 387 J/kg·C

Water requires 5× more energy than iron to heat the same amount!

## 11.2 Why Metal Feels Colder

### The Paradox

Wood and metal at room temperature.

**Why does metal feel colder when you touch it?**



## 11.2 Why Metal Feels Colder

### The Paradox

Wood and metal at room temperature.

**Why does metal feel colder when you touch it?**

**The answer:**

- Both are same temperature

## 11.2 Why Metal Feels Colder

### The Paradox

Wood and metal at room temperature.

**Why does metal feel colder when you touch it?**

**The answer:**

- Both are same temperature
- Metal conducts heat faster than wood

## 11.2 Why Metal Feels Colder

### The Paradox

Wood and metal at room temperature.

**Why does metal feel colder when you touch it?**

**The answer:**

- Both are same temperature
- Metal conducts heat faster than wood
- Metal pulls heat from your hand rapidly

## 11.2 Why Metal Feels Colder

### The Paradox

Wood and metal at room temperature.

**Why does metal feel colder when you touch it?**

**The answer:**

- Both are same temperature
- Metal conducts heat faster than wood
- Metal pulls heat from your hand rapidly
- You sense rate of heat loss, not temperature!

## 11.2 Three Modes of Heat Transfer




Figure: Fireplace: all three modes at once

## 11.2 Three Modes of Heat Transfer




Figure: Fireplace: all three modes at once

- **Conduction:** through physical contact
- **Convection:** by fluid movement
- **Radiation:** by electromagnetic waves

## 11.2 Conduction: Touch Transfer

### The Mental Model

Molecules vibrating faster bump into slower neighbors, sharing energy.  
Like dominoes falling in sequence.

## 11.2 Conduction: Touch Transfer

### The Mental Model

Molecules vibrating faster bump into slower neighbors, sharing energy. Like dominoes falling in sequence.




Figure: Energy transfers through collisions



## 11.2 Conduction: Touch Transfer

### The Mental Model

Molecules vibrating faster bump into slower neighbors, sharing energy. Like dominoes falling in sequence.





Figure: Energy transfers through collisions

**Good conductors:** metals (copper, aluminum, gold)

**Poor conductors:** wood, plastic, rubber (insulators)

## 11.2 Convection: Fluid Flow



**Figure:** Heated air rises, cool air sinks

## 11.2 Convection: Fluid Flow




Figure: Heated air rises, cool air sinks






Figure: Water circulation in pot

## 11.2 Convection: Fluid Flow



**Figure:** Heated air rises, cool air sinks



**Figure:** Water circulation in pot

**Key idea:** Hot fluid expands, becomes less dense, rises

## 11.2 Radiation: No Medium Needed

### The Mystery

How does Sun's energy reach Earth through vacuum of space?

## 11.2 Radiation: No Medium Needed

### The Mystery

How does Sun's energy reach Earth through vacuum of space?

### The Source Code

**Radiation** = energy transfer by electromagnetic waves

No physical medium required!

## 11.2 Radiation: No Medium Needed

### The Mystery

How does Sun's energy reach Earth through vacuum of space?

### The Source Code

**Radiation** = energy transfer by electromagnetic waves  
No physical medium required!

### Examples:

- Feel warmth from fire without touching
- Microwave oven heating food
- Infrared heat lamps
- Sun warming Earth

## 11.2 Color and Radiation

### Color affects absorption and emission:

- **Black:** best absorber AND radiator



## 11.2 Color and Radiation

### Color affects absorption and emission:

- **Black:** best absorber AND radiator
- **White:** worst absorber AND radiator

## 11.2 Color and Radiation

### Color affects absorption and emission:

- **Black:** best absorber AND radiator
- **White:** worst absorber AND radiator
- **Shiny:** reflects radiation like mirror

## 11.2 Color and Radiation

### Color affects absorption and emission:

- **Black:** best absorber AND radiator
- **White:** worst absorber AND radiator
- **Shiny:** reflects radiation like mirror

### Real-World Application

Hot climates: wear white to reflect heat

Cold nights: black asphalt radiates heat faster than grass

Space blankets: shiny surface reflects body heat back

# Attempt: Heating Water

## The Challenge (4 min, silent)

You heat 2.0 kg of water from 20 C to 80 C.

### Given:

- $m = 2.0 \text{ kg}$
- $c_{\text{water}} = 4186 \text{ J/kg}\cdot\text{C}$
- $T_i = 20 \text{ C}, T_f = 80 \text{ C}$

**Find:** Heat energy required (Q)

*Can you calculate energy needed? Work silently.*

# Compare: Heat Calculation

**Turn and talk (2 min):**

- 1 What equation did you use?
- 2 How did you calculate  $\Delta T$ ?
- 3 What units did you get for  $Q$ ?

# Compare: Heat Calculation

**Turn and talk (2 min):**

- 1 What equation did you use?
- 2 How did you calculate  $\Delta T$ ?
- 3 What units did you get for  $Q$ ?

**Name wheel:** One pair share your approach (not your answer).

# Reveal: Energy to Heat Water

**Self-correct in a different color:**

**Step 1:** Calculate temperature change

$$\Delta T = T_f - T_i = 80 - 20 = 60 \text{ C}$$

# Reveal: Energy to Heat Water

**Self-correct in a different color:**

**Step 1:** Calculate temperature change

$$\Delta T = T_f - T_i = 80 - 20 = 60 \text{ C}$$

**Step 2:** Use  $Q = mc\Delta T$



# Reveal: Energy to Heat Water

**Self-correct in a different color:**

**Step 1:** Calculate temperature change

$$\Delta T = T_f - T_i = 80 - 20 = 60 \text{ C}$$

**Step 2:** Use  $Q = mc\Delta T$

**Substitute:**  $Q = (2.0 \text{ kg})(4186 \text{ J/kg}\cdot\text{C})(60 \text{ C})$

# Reveal: Energy to Heat Water

**Self-correct in a different color:**

**Step 1:** Calculate temperature change

$$\Delta T = T_f - T_i = 80 - 20 = 60 \text{ C}$$

**Step 2:** Use  $Q = mc\Delta T$

**Substitute:**  $Q = (2.0 \text{ kg})(4186 \text{ J/kg}\cdot\text{C})(60 \text{ C})$

$$Q = 502,320 \text{ J} = \boxed{502 \text{ kJ}}$$

# Reveal: Energy to Heat Water

**Self-correct in a different color:**

**Step 1:** Calculate temperature change

$$\Delta T = T_f - T_i = 80 - 20 = 60 \text{ C}$$

**Step 2:** Use  $Q = mc\Delta T$

**Substitute:**  $Q = (2.0 \text{ kg})(4186 \text{ J/kg}\cdot\text{C})(60 \text{ C})$

$$Q = 502,320 \text{ J} = \boxed{502 \text{ kJ}}$$

**Check:** About 500 kJ to heat 2 L water by 60 degrees. Reasonable!

# Learning Objectives

By the end of this section, you will be able to:

- **11.3:** Explain changes in heat during changes of state

# Learning Objectives

By the end of this section, you will be able to:


- **11.3:** Explain changes in heat during changes of state
- **11.3:** Describe latent heats of fusion and vaporization

# Learning Objectives

By the end of this section, you will be able to:


- **11.3:** Explain changes in heat during changes of state
- **11.3:** Describe latent heats of fusion and vaporization
- **11.3:** Solve problems involving phase changes

## 11.3 The Four Phases of Matter



**Figure:** Solid, liquid, gas, and plasma

# 11.3 The Four Phases of Matter



**Figure:** Solid, liquid, gas, and plasma

**Energy ranking:** Solid | Liquid | Gas | Plasma



## 11.3 Phase Changes: Energy In/Out

### **Adding energy (heating):**

- Melting: solid  $\rightarrow$  liquid

## 11.3 Phase Changes: Energy In/Out

### **Adding energy (heating):**

- Melting: solid  $\rightarrow$  liquid
- Vaporization: liquid  $\rightarrow$  gas

## 11.3 Phase Changes: Energy In/Out

### **Adding energy (heating):**

- Melting: solid  $\rightarrow$  liquid
- Vaporization: liquid  $\rightarrow$  gas
- Sublimation: solid  $\rightarrow$  gas (skips liquid!)

## 11.3 Phase Changes: Energy In/Out

### **Adding energy (heating):**

- Melting: solid  $\rightarrow$  liquid
- Vaporization: liquid  $\rightarrow$  gas
- Sublimation: solid  $\rightarrow$  gas (skips liquid!)

### **Removing energy (cooling):**

- Freezing: liquid  $\rightarrow$  solid

# 11.3 Phase Changes: Energy In/Out

## Adding energy (heating):

- Melting: solid  $\rightarrow$  liquid
- Vaporization: liquid  $\rightarrow$  gas
- Sublimation: solid  $\rightarrow$  gas (skips liquid!)

## Removing energy (cooling):

- Freezing: liquid  $\rightarrow$  solid
- Condensation: gas  $\rightarrow$  liquid

# 11.3 Phase Changes: Energy In/Out

## Adding energy (heating):

- Melting: solid  $\rightarrow$  liquid
- Vaporization: liquid  $\rightarrow$  gas
- Sublimation: solid  $\rightarrow$  gas (skips liquid!)

## Removing energy (cooling):

- Freezing: liquid  $\rightarrow$  solid
- Condensation: gas  $\rightarrow$  liquid
- Deposition: gas  $\rightarrow$  solid

## 11.3 Latent Heat: The Hidden Energy

### The Paradox

You heat ice at 0 C. Temperature stays at 0 C even as you add energy.

**Where does the energy go?**

## 11.3 Latent Heat: The Hidden Energy

### The Paradox

You heat ice at 0 C. Temperature stays at 0 C even as you add energy.

**Where does the energy go?**

### The Revelation

Energy breaks bonds between molecules, not increase speed.

No temperature change during phase transition!



## 11.3 Latent Heat: The Hidden Energy

### The Paradox

You heat ice at 0 C. Temperature stays at 0 C even as you add energy.

**Where does the energy go?**

### The Revelation

Energy breaks bonds between molecules, not increase speed.

No temperature change during phase transition!

**Latent heat** = hidden energy used to change phase without changing temperature

## 11.3 Phase Diagram: Ice to Steam




Figure: Temperature vs energy for water

## 11.3 Phase Diagram: Ice to Steam




Figure: Temperature vs energy for water

**Flat regions** = phase changes (temp constant)

**Sloped regions** = temperature increasing

## 11.3 Latent Heat Equations

### Nature's Rules for Phase Change

$$Q = mL_f \quad (\text{melting/freezing})$$

$$Q = mL_v \quad (\text{vaporization/condensation})$$

## 11.3 Latent Heat Equations

### Nature's Rules for Phase Change

$$Q = mL_f \quad (\text{melting/freezing})$$

$$Q = mL_v \quad (\text{vaporization/condensation})$$

#### For water:

- $L_f = 334 \text{ kJ/kg}$  (latent heat of fusion)
- $L_v = 2256 \text{ kJ/kg}$  (latent heat of vaporization)

## 11.3 Latent Heat Equations

### Nature's Rules for Phase Change

$$Q = mL_f \quad (\text{melting/freezing})$$

$$Q = mL_v \quad (\text{vaporization/condensation})$$

#### For water:

- $L_f = 334 \text{ kJ/kg}$  (latent heat of fusion)
- $L_v = 2256 \text{ kJ/kg}$  (latent heat of vaporization)

Note: No  $\Delta T$  in these equations - temperature doesn't change!

# Attempt: Melting Ice

## The Challenge (3 min, silent)

How much energy is needed to melt 0.50 kg of ice at 0 C?

### Given:

- $m = 0.50 \text{ kg}$
- $L_f = 334 \text{ kJ/kg}$
- Ice already at 0 C (melting point)

**Find:** Heat energy  $Q$  required

*Can you calculate the hidden energy? Work silently.*

# Compare: Latent Heat

**Turn and talk (2 min):**

- 1 Which equation did you choose?
- 2 Why didn't you use  $Q = mc\Delta T$ ?
- 3 What value did you use for latent heat?



# Compare: Latent Heat

**Turn and talk (2 min):**

- 1 Which equation did you choose?
- 2 Why didn't you use  $Q = mc\Delta T$ ?
- 3 What value did you use for latent heat?

**Name wheel:** One pair share your reasoning.

# Reveal: Energy to Melt Ice

**Self-correct in a different color:**

**Key insight:** Phase change, so use  $Q = mL_f$

# Reveal: Energy to Melt Ice

**Self-correct in a different color:**

**Key insight:** Phase change, so use  $Q = mL_f$

**Substitute:**  $Q = (0.50 \text{ kg})(334 \text{ kJ/kg})$

# Reveal: Energy to Melt Ice

**Self-correct in a different color:**

**Key insight:** Phase change, so use  $Q = mL_f$

**Substitute:**  $Q = (0.50 \text{ kg})(334 \text{ kJ/kg})$

$$Q = \boxed{167 \text{ kJ}}$$

# Reveal: Energy to Melt Ice

**Self-correct in a different color:**

**Key insight:** Phase change, so use  $Q = mL_f$

**Substitute:**  $Q = (0.50 \text{ kg})(334 \text{ kJ/kg})$

$$Q = \boxed{167 \text{ kJ}}$$

**Check:** That's enough energy to raise 1 kg of water by 40 C!

Phase changes require enormous energy.

## 11.3 Why Sweating Cools You

### Real-World Application

When sweat evaporates:

- 1 Water absorbs  $2256 \text{ kJ/kg}$  from your skin

## 11.3 Why Sweating Cools You

### Real-World Application

When sweat evaporates:

- 1 Water absorbs  $2256 \text{ kJ/kg}$  from your skin
- 2 Undergoes phase change: liquid  $\rightarrow$  gas

## 11.3 Why Sweating Cools You

### Real-World Application

When sweat evaporates:

- 1 Water absorbs  $2256 \text{ kJ/kg}$  from your skin
- 2 Undergoes phase change: liquid  $\rightarrow$  gas
- 3 Your skin temperature drops



## 11.3 Why Sweating Cools You

### Real-World Application

When sweat evaporates:

- 1 Water absorbs  $2256 \text{ kJ/kg}$  from your skin
- 2 Undergoes phase change: liquid  $\rightarrow$  gas
- 3 Your skin temperature drops

Evaporation is most effective cooling method for your body!

## 11.3 Why Sweating Cools You

### Real-World Application

When sweat evaporates:

- 1 Water absorbs  $2256 \text{ kJ/kg}$  from your skin
- 2 Undergoes phase change: liquid  $\rightarrow$  gas
- 3 Your skin temperature drops

Evaporation is most effective cooling method for your body!

**On humid days:** Less evaporation  $\rightarrow$  less cooling  $\rightarrow$  you feel hotter

## 11.3 Condensation Releases Heat

### The Reverse Process

Condensation releases same energy that vaporization absorbed.

Gas  $\rightarrow$  liquid releases 2256 kJ/kg for water.

## 11.3 Condensation Releases Heat

### The Reverse Process

Condensation releases same energy that vaporization absorbed.

Gas  $\rightarrow$  liquid releases 2256 kJ/kg for water.

### Why hurricanes are so powerful:

- Water vapor condenses in storm

## 11.3 Condensation Releases Heat

### The Reverse Process

Condensation releases same energy that vaporization absorbed.

Gas  $\rightarrow$  liquid releases 2256 kJ/kg for water.

### Why hurricanes are so powerful:

- Water vapor condenses in storm
- Releases enormous latent heat

## 11.3 Condensation Releases Heat

### The Reverse Process

Condensation releases same energy that vaporization absorbed.

Gas  $\rightarrow$  liquid releases 2256 kJ/kg for water.

### Why hurricanes are so powerful:

- Water vapor condenses in storm
- Releases enormous latent heat
- Heats surrounding air

## 11.3 Condensation Releases Heat

### The Reverse Process

Condensation releases same energy that vaporization absorbed.

Gas  $\rightarrow$  liquid releases 2256 kJ/kg for water.

### Why hurricanes are so powerful:

- Water vapor condenses in storm
- Releases enormous latent heat
- Heats surrounding air
- Creates powerful updrafts and winds

# The Revelations of Thermal Physics

## What You Now Know

- 1 Temperature = average molecular kinetic energy



# The Revelations of Thermal Physics

## What You Now Know

- 1 Temperature = average molecular kinetic energy
- 2 Heat = energy transfer from hot to cold

# The Revelations of Thermal Physics

## What You Now Know

- 1 Temperature = average molecular kinetic energy
- 2 Heat = energy transfer from hot to cold
- 3  $Q = mc\Delta T$  for temperature changes

# The Revelations of Thermal Physics

## What You Now Know

- 1 Temperature = average molecular kinetic energy
- 2 Heat = energy transfer from hot to cold
- 3  $Q = mc\Delta T$  for temperature changes
- 4 Three transfer modes: conduction, convection, radiation

# The Revelations of Thermal Physics

## What You Now Know

- 1 Temperature = average molecular kinetic energy
- 2 Heat = energy transfer from hot to cold
- 3  $Q = mc\Delta T$  for temperature changes
- 4 Three transfer modes: conduction, convection, radiation
- 5  $Q = mL$  for phase changes (no temp change!)

# The Revelations of Thermal Physics

## What You Now Know

- 1 Temperature = average molecular kinetic energy
- 2 Heat = energy transfer from hot to cold
- 3  $Q = mc\Delta T$  for temperature changes
- 4 Three transfer modes: conduction, convection, radiation
- 5  $Q = mL$  for phase changes (no temp change!)
- 6 Water's high specific heat stabilizes climate

# The Revelations of Thermal Physics

## What You Now Know

- 1 Temperature = average molecular kinetic energy
- 2 Heat = energy transfer from hot to cold
- 3  $Q = mc\Delta T$  for temperature changes
- 4 Three transfer modes: conduction, convection, radiation
- 5  $Q = mL$  for phase changes (no temp change!)
- 6 Water's high specific heat stabilizes climate
- 7 Phase changes require enormous energy

# Key Equations

## Temperature Conversions:

$$T_{oF} = \frac{9}{5} T_{oC} + 32 \quad (1)$$

$$T_{oC} = \frac{5}{9} (T_{oF} - 32) \quad (2)$$

$$T_K = T_{oC} + 273.15 \quad (3)$$

## Heat Transfer:

$$Q = mc\Delta T \quad (\text{temperature change}) \quad (4)$$

$$Q = mL_f \quad (\text{melting/freezing}) \quad (5)$$

$$Q = mL_v \quad (\text{vaporization/condensation}) \quad (6)$$

## Water Constants:

- $c_{\text{water}} = 4186 \text{ J/kg}\cdot\text{C}$
- $L_f = 334 \text{ kJ/kg}$
- $L_v = 2256 \text{ kJ/kg}$

Complete the assigned problems  
posted on the LMS