

PHYS11 CH:18 The Force That Moves Everything

Electric Charge and Conservation

Mr. Gullo

December 2025

Outline

Why Does Hair Stand on End?



Why Does Hair Stand on End?



What invisible force makes strands of hair repel each other?

What if everything you touch
is held together by invisible forces?

The Invisible World

What if everything you touch
is held together by invisible forces?

The atoms in your fingertips never actually touch the atoms in this desk...

The Invisible World

What if everything you touch
is held together by invisible forces?

The atoms in your fingertips never actually touch the atoms in this desk...

Electric forces keep them apart.

Learning Objectives

By the end of this lesson, you will be able to:

- **18.1:** Describe positive and negative electric charges

Learning Objectives

By the end of this lesson, you will be able to:

- **18.1:** Describe positive and negative electric charges
- **18.1:** Use conservation of charge to calculate charge transfers

Learning Objectives

By the end of this lesson, you will be able to:

- **18.1:** Describe positive and negative electric charges
- **18.1:** Use conservation of charge to calculate charge transfers
- **18.1:** Characterize conductors vs insulators

Learning Objectives

By the end of this lesson, you will be able to:

- **18.1:** Describe positive and negative electric charges
- **18.1:** Use conservation of charge to calculate charge transfers
- **18.1:** Characterize conductors vs insulators
- **18.1:** Describe electric polarization and charging by induction

18.1 Two Types of Charge

Nature's Binary Code

Electric charge is a property of matter that causes objects to attract or repel each other.

18.1 Two Types of Charge

Nature's Binary Code

Electric charge is a property of matter that causes objects to attract or repel each other.

The discovery:

- Glass rubbed with silk: glass becomes positive, silk becomes negative

18.1 Two Types of Charge

Nature's Binary Code

Electric charge is a property of matter that causes objects to attract or repel each other.

The discovery:

- Glass rubbed with silk: glass becomes positive, silk becomes negative
- Like charges **repel** each other

18.1 Two Types of Charge

Nature's Binary Code

Electric charge is a property of matter that causes objects to attract or repel each other.

The discovery:

- Glass rubbed with silk: glass becomes positive, silk becomes negative
- Like charges **repel** each other
- Unlike charges **attract** each other

18.1 Two Types of Charge

Nature's Binary Code

Electric charge is a property of matter that causes objects to attract or repel each other.


The discovery:

- Glass rubbed with silk: glass becomes positive, silk becomes negative
- Like charges **repel** each other
- Unlike charges **attract** each other


The Mental Model

Charge is nature's binary system: positive or negative. No neutral charge exists.

18.1 Experimental Evidence



(a)




(b)



(c)

18.1 Experimental Evidence



Pattern: Glass rods repel each other. Silk cloths repel each other. Glass attracts silk.

18.1 The Discovery of the Electron

The mystery (1897): What carries negative charge?

18.1 The Discovery of the Electron

The mystery (1897): What carries negative charge?

The experiment: J.J. Thomson studied cathode rays

18.1 The Discovery of the Electron

The mystery (1897): What carries negative charge?

The experiment: J.J. Thomson studied cathode rays

The revelation: Cathode rays are particles carrying negative charge

18.1 The Discovery of the Electron

The mystery (1897): What carries negative charge?



The experiment: J.J. Thomson studied cathode rays

The revelation: Cathode rays are particles carrying negative charge



Universal Law: The Electron

The electron carries the fundamental unit of *negative* electric charge.

18.1 Inside the Atom



18.1 Inside the Atom



Rutherford's model: electrons orbit a tiny, dense nucleus of protons.

18.1 The Fundamental Unit of Charge

Nature's Quantum

$$e = 1.602 \times 10^{-19} \text{ C}$$

The fundamental unit of electric **charge** (magnitude).

18.1 The Fundamental Unit of Charge

Nature's Quantum

$$e = 1.602 \times 10^{-19} \text{ C}$$

The fundamental unit of electric **charge** (magnitude).

The carriers:

- Proton **charge**: $+e = +1.602 \times 10^{-19} \text{ C}$

18.1 The Fundamental Unit of Charge

Nature's Quantum

$$e = 1.602 \times 10^{-19} \text{ C}$$

The fundamental unit of electric **charge** (magnitude).

The carriers:

- Proton **charge**: $+e = +1.602 \times 10^{-19} \text{ C}$
- Electron **charge**: $-e = -1.602 \times 10^{-19} \text{ C}$

18.1 The Fundamental Unit of Charge

Nature's Quantum

$$e = 1.602 \times 10^{-19} \text{ C}$$

The fundamental unit of electric **charge** (magnitude).

The carriers:


- Proton **charge**: $+e = +1.602 \times 10^{-19} \text{ C}$
- Electron **charge**: $-e = -1.602 \times 10^{-19} \text{ C}$

The Paradox


Civilian: "Why is charge so tiny?"

Physicist: "It takes 6.25×10^{18} protons to make just 1 coulomb!"

18.1 Measuring the Electron Charge



18.1 Measuring the Electron Charge



Millikan Oil-Drop Experiment (1909):

- Spray oil droplets between charged plates
- Balance electric force against gravity
- Measure charge on individual drops

18.1 Charge Quantization

Discovery: Charge always comes in multiples of e

18.1 Charge Quantization

Discovery: Charge always comes in multiples of e

Universal Law: Charge Quantization

$$Q = ne$$

where n is an integer ($\pm 1, \pm 2, \pm 3, \dots$)

18.1 Charge Quantization

Discovery: Charge always comes in multiples of e

Universal Law: Charge Quantization

$$Q = ne$$

where n is an integer ($\pm 1, \pm 2, \pm 3, \dots$)

Meaning: You can have 5 electrons or 5 million, but never 5.5 electrons.

18.1 Conservation of Charge

Nature's Accounting System

$$q_{\text{initial}} = q_{\text{final}}$$

Electric **charge** cannot be created or destroyed.

18.1 Conservation of Charge

Nature's Accounting System

$$q_{\text{initial}} = q_{\text{final}}$$

Electric **charge** cannot be created or destroyed.

What this means:

- Total **charge** before interaction = total **charge** after

18.1 Conservation of Charge

Nature's Accounting System

$$q_{\text{initial}} = q_{\text{final}}$$

Electric **charge** cannot be created or destroyed.

What this means:

- Total **charge** before interaction = total **charge** after
- **Charge** can move, but net **charge** stays constant

18.1 Conservation of Charge

Nature's Accounting System

$$q_{\text{initial}} = q_{\text{final}}$$

Electric **charge** cannot be created or destroyed.

What this means:

- Total **charge** before interaction = total **charge** after
- **Charge** can move, but net **charge** stays constant
- Most fundamental conservation law in physics

18.1 Conductors vs Insulators

Conductors

Materials that allow **charge** to move freely

Examples:

- Metals (copper, silver, aluminum)
- Electrons loosely bound

18.1 Conductors vs Insulators

Conductors

Materials that allow **charge** to move freely

Examples:

- Metals (copper, silver, aluminum)
- Electrons loosely bound

Insulators

Materials that prevent **charge** from moving

Examples:

- Rubber, plastic, glass, wood
- Electrons tightly bound

18.1 Conductors vs Insulators

Conductors

Materials that allow **charge** to move freely

Examples:

- Metals (copper, silver, aluminum)
- Electrons loosely bound

Insulators

Materials that prevent **charge** from moving


Examples:

- Rubber, plastic, glass, wood
- Electrons tightly bound


The Mental Model

Conductor = highway for electrons. Insulator = roadblock.

18.1 The Conductivity Spectrum





18.1 The Conductivity Spectrum




Semiconductors: Between conductors and insulators (silicon, germanium)

18.1 Charge Distribution



18.1 Charge Distribution



Conductor: Charge spreads to outer surface (repulsion wins).

Insulator: Charge stays in place (can't move).

18.1 Transferring Charge: Contact

Charging by contact: Surfaces touch and share electrons

18.1 Transferring Charge: Contact

Charging by contact: Surfaces touch and share electrons

Example: Walking across carpet

- Shoes rub carpet

18.1 Transferring Charge: Contact

Charging by contact: Surfaces touch and share electrons

Example: Walking across carpet

- Shoes rub carpet
- Electrons transfer from carpet to shoes

18.1 Transferring Charge: Contact

Charging by contact: Surfaces touch and share electrons

Example: Walking across carpet

- Shoes rub carpet
- Electrons transfer from carpet to shoes
- You accumulate excess negative charge

18.1 Transferring Charge: Contact

Charging by contact: Surfaces touch and share electrons

Example: Walking across carpet

- Shoes rub carpet
- Electrons transfer from carpet to shoes
- You accumulate excess negative charge
- Touch doorknob → ZAP!

18.1 Transferring Charge: Contact

Charging by contact: Surfaces touch and share electrons



Example: Walking across carpet

- Shoes rub carpet
- Electrons transfer from carpet to shoes
- You accumulate excess negative charge
- Touch doorknob → ZAP!


Real-World: Static Shock

Rubbing increases contact between materials, transferring more electrons.

18.1 Transferring Charge: Conduction




18.1 Transferring Charge: Conduction



Charging by conduction: Touching charged object to neutral object

18.1 Transferring Charge: Conduction



Charging by conduction: Touching charged object to neutral object
Charges redistribute to equalize - like water finding same level.

18.1 Polarization

What is polarization? Separation of positive and negative charges within an object

18.1 Polarization

What is polarization? Separation of positive and negative charges within an object

Example: Child on slide (Figure 18.1)

- Excess positive charge on body

18.1 Polarization

What is polarization? Separation of positive and negative charges within an object

Example: Child on slide (Figure 18.1)

- Excess positive charge on body
- Positive charges repel each other

18.1 Polarization

What is polarization? Separation of positive and negative charges within an object

Example: Child on slide (Figure 18.1)

- Excess positive charge on body
- Positive charges repel each other
- Spread to extremities (hair)

18.1 Polarization

What is polarization? Separation of positive and negative charges within an object

Example: Child on slide (Figure 18.1)

- Excess positive charge on body
- Positive charges repel each other
- Spread to extremities (hair)
- Hair strands repel → stand on end!

18.1 Polarization

What is polarization? Separation of positive and negative charges within an object


Example: Child on slide (Figure 18.1)

- Excess positive charge on body
- Positive charges repel each other
- Spread to extremities (hair)
- Hair strands repel → stand on end!


The Mental Model

Like dominos - push on one end, disturbance travels to other end.

18.1 Charging by Induction




18.1 Charging by Induction




Induction: Creating charge separation by approaching a charged object (no touching!)

18.1 Van de Graaff Generator



18.1 Van de Graaff Generator



How it works:

- Belt transfers electrons to metal globe
- Electrons spread over outer surface (repulsion)
- Can accumulate millions of volts!

18.1 Hair-Raising Physics



18.1 Hair-Raising Physics



Why does hair stand up?

- Each hair strand gets excess charge (same sign)
- Like charges repel
- Strands push away from each other as far as possible

Attempt: Conservation Challenge

The Challenge (3 min, silent)

Two metal spheres initially have charges of $+4\text{ C}$ and $+8\text{ C}$. After touching each other, one sphere has $+10\text{ C}$.

Given:

- Blue sphere initial: $q_1 = +4\text{ C}$
- Red sphere initial: $q_2 = +8\text{ C}$
- Blue sphere final: $q'_1 = +10\text{ C}$

Find: Final charge on red sphere q'_2

Can you use conservation of charge? Work silently.

Compare: Conservation Strategy

Turn and talk (2 min):

- 1 What law did you use?
- 2 What is the total initial charge?
- 3 How did you find the final charge on the red sphere?

Compare: Conservation Strategy

Turn and talk (2 min):

- 1 What law did you use?
- 2 What is the total initial charge?
- 3 How did you find the final charge on the red sphere?

Name wheel: One pair share your approach (not your answer).

Reveal: Conservation in Action

Self-correct in a different color:

Law: $q_{\text{initial}} = q_{\text{final}}$

Reveal: Conservation in Action

Self-correct in a different color:

Law: $q_{\text{initial}} = q_{\text{final}}$

Step 1: Find total initial charge

$$q_{\text{initial}} = +4 \text{ C} + 8 \text{ C} = +12 \text{ C}$$

Reveal: Conservation in Action

Self-correct in a different color:

Law: $q_{\text{initial}} = q_{\text{final}}$

Step 1: Find total initial charge

$$q_{\text{initial}} = +4 \text{ C} + 8 \text{ C} = +12 \text{ C}$$

Step 2: Apply conservation law

$$q_{\text{final}} = +10 \text{ C} + q_2'$$

Reveal: Conservation in Action

Self-correct in a different color:

Law: $q_{\text{initial}} = q_{\text{final}}$

Step 1: Find total initial charge

$$q_{\text{initial}} = +4 \text{ C} + 8 \text{ C} = +12 \text{ C}$$

Step 2: Apply conservation law

$$q_{\text{final}} = +10 \text{ C} + q'_2$$

Step 3: Solve for q'_2

$$+12 \text{ C} = +10 \text{ C} + q'_2$$

$$q'_2 = +2 \text{ C}$$

Reveal: Conservation in Action

Self-correct in a different color:

Law: $q_{\text{initial}} = q_{\text{final}}$

Step 1: Find total initial **charge**

$$q_{\text{initial}} = +4 \text{ C} + 8 \text{ C} = +12 \text{ C}$$

Step 2: Apply conservation law

$$q_{\text{final}} = +10 \text{ C} + q'_2$$

Step 3: Solve for q'_2

$$+12 \text{ C} = +10 \text{ C} + q'_2$$

$$q'_2 = +2 \text{ C}$$

Check: $12 = 10 + 2$ ✓ **Charge** is conserved!

Attempt: Counting Electrons

The Challenge (3 min, silent)

An ink droplet in a printer has net **charge** $q = -1.0 \times 10^{-10} \text{ C}$ after passing through an electron beam.

Given:

- Droplet **charge**: $q = -1.0 \times 10^{-10} \text{ C}$
- Electron **charge**: $e = -1.602 \times 10^{-19} \text{ C}$

Find: Number of electrons captured by droplet

How many electrons does it take? Work silently.

Compare: Quantization Strategy

Turn and talk (2 min):

- 1 What equation relates total **charge** to number of electrons?
- 2 Did you multiply or divide?
- 3 What happens with the negative signs?

Compare: Quantization Strategy

Turn and talk (2 min):

- 1 What equation relates total **charge** to number of electrons?
- 2 Did you multiply or divide?
- 3 What happens with the negative signs?

Name wheel: One pair share your approach (not your answer).

Reveal: Quantization at Work

Self-correct in a different color:

Equation: $Q = ne$, so $n = \frac{Q}{e}$

Reveal: Quantization at Work

Self-correct in a different color:

Equation: $Q = ne$, so $n = \frac{Q}{e}$

Substitute:

$$n = \frac{q}{q_{e^-}} = \frac{-1.0 \times 10^{-10} \text{ C}}{-1.602 \times 10^{-19} \text{ C}}$$

Reveal: Quantization at Work

Self-correct in a different color:

Equation: $Q = ne$, so $n = \frac{Q}{e}$

Substitute:

$$n = \frac{q}{q_{e^-}} = \frac{-1.0 \times 10^{-10} \text{ C}}{-1.602 \times 10^{-19} \text{ C}}$$

Simplify:

$$n = \frac{1.0 \times 10^{-10}}{1.602 \times 10^{-19}} = 6.24 \times 10^8$$

Reveal: Quantization at Work

Self-correct in a different color:

Equation: $Q = ne$, so $n = \frac{Q}{e}$

Substitute:

$$n = \frac{q}{q_{e^-}} = \frac{-1.0 \times 10^{-10} \text{ C}}{-1.602 \times 10^{-19} \text{ C}}$$

Simplify:

$$n = \frac{1.0 \times 10^{-10}}{1.602 \times 10^{-19}} = 6.24 \times 10^8$$

$n = 6 \times 10^8 \text{ electrons}$

Reveal: Quantization at Work

Self-correct in a different color:

Equation: $Q = ne$, so $n = \frac{Q}{e}$

Substitute:

$$n = \frac{q}{q_{e^-}} = \frac{-1.0 \times 10^{-10} \text{ C}}{-1.602 \times 10^{-19} \text{ C}}$$

Simplify:

$$n = \frac{1.0 \times 10^{-10}}{1.602 \times 10^{-19}} = 6.24 \times 10^8$$

$n = 6 \times 10^8 \text{ electrons}$

Check: About 600 million electrons - seems large but atoms have 10^{16} atoms!

What You Now Know

The Revelations

- 1 Two types of **charge**: positive (protons) and negative (electrons)

What You Now Know

The Revelations

- 1 Two types of **charge**: positive (protons) and negative (electrons)
- 2 Fundamental **charge**: $e = 1.602 \times 10^{-19} \text{ C}$

What You Now Know

The Revelations

- 1 Two types of **charge**: positive (protons) and negative (electrons)
- 2 Fundamental **charge**: $e = 1.602 \times 10^{-19} \text{ C}$
- 3 **Charge** quantization: $Q = ne$ (only integer multiples)

What You Now Know

The Revelations

- 1 Two types of **charge**: positive (protons) and negative (electrons)
- 2 Fundamental **charge**: $e = 1.602 \times 10^{-19} \text{ C}$
- 3 **Charge** quantization: $Q = ne$ (only integer multiples)
- 4 Conservation: $q_{\text{initial}} = q_{\text{final}}$ (never violated)

What You Now Know

The Revelations

- 1 Two types of **charge**: positive (protons) and negative (electrons)
- 2 Fundamental **charge**: $e = 1.602 \times 10^{-19} \text{ C}$
- 3 **Charge** quantization: $Q = ne$ (only integer multiples)
- 4 Conservation: $q_{\text{initial}} = q_{\text{final}}$ (never violated)
- 5 Conductors let **charge** move; insulators don't

What You Now Know

The Revelations

- 1 Two types of **charge**: positive (protons) and negative (electrons)
- 2 Fundamental **charge**: $e = 1.602 \times 10^{-19} \text{ C}$
- 3 **Charge** quantization: $Q = ne$ (only integer multiples)
- 4 Conservation: $q_{\text{initial}} = q_{\text{final}}$ (never violated)
- 5 Conductors let **charge** move; insulators don't
- 6 Transfer methods: contact, conduction, induction

Key Equations

$$e = 1.602 \times 10^{-19} \text{ C} \quad \text{(fundamental charge)} \quad (1)$$

$$Q = ne \quad \text{(charge quantization)} \quad (2)$$

$$q_{\text{initial}} = q_{\text{final}} \quad \text{(conservation of charge)} \quad (3)$$

Complete the assigned problems
posted on the LMS

Temporary page!

\LaTeX was unable to guess the total number of pages correctly. There was some unprocessed data that should have been added to the document, so this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will disappear, because \LaTeX now knows how many pages to expect for the document.