

PHYS11 CH:1-3: Introduction to Physics and 1D Kinematics

From Core Concepts to Equations of Motion

Mr. Gullo

July 23, 2025

Learning Objectives I

After this lesson, you will be able to:

- Define physics and its role as a fundamental science.
- Describe the scientific method.
- Differentiate between physical quantities, units, accuracy, and precision.
- Explain the rules for significant figures in calculations.
- Distinguish between distance and displacement, and speed and velocity.
- Analyze motion using position vs. time and velocity vs. time graphs.
- Define acceleration as the rate of change of velocity.
- Apply kinematic equations to solve problems involving one-dimensional motion with constant acceleration.

What is Physics? (Sec 1.1)

- **Physics** is the most fundamental of the sciences, concerning itself with energy, matter, space and time, and their interactions.

What is Physics? (Sec 1.1)

- **Physics** is the most fundamental of the sciences, concerning itself with energy, matter, space and time, and their interactions.
- **Modern Physics** includes two revolutionary theories:
 - **Relativity**: Describes how time, space, and gravity can be different for different observers.
 - **Quantum Mechanics**: Describes the behavior of subatomic particles.

What is Physics? (Sec 1.1)

- **Physics** is the most fundamental of the sciences, concerning itself with energy, matter, space and time, and their interactions.
- **Modern Physics** includes two revolutionary theories:
 - **Relativity:** Describes how time, space, and gravity can be different for different observers.
 - **Quantum Mechanics:** Describes the behavior of subatomic particles.
- Physics provides the fundamental principles that underlie all other sciences, including chemistry, biology, and geology.

The Scientific Method (Sec 1.2)

Science is a process for discovering the order and simplicity in nature.

- **Process:** Observation \rightarrow Hypothesis \rightarrow Experiment \rightarrow Conclusion.

The Scientific Method (Sec 1.2)

Science is a process for discovering the order and simplicity in nature.

- **Process:** Observation \rightarrow Hypothesis \rightarrow Experiment \rightarrow Conclusion.
- **Theory:** A scientific explanation that is supported by a large body of experimental results. A theory is a powerful and well-substantiated explanation.
 - Example: The Theory of General Relativity

The Scientific Method (Sec 1.2)

Science is a process for discovering the order and simplicity in nature.

- **Process:** Observation \rightarrow Hypothesis \rightarrow Experiment \rightarrow Conclusion.
- **Theory:** A scientific explanation that is supported by a large body of experimental results. A theory is a powerful and well-substantiated explanation.
 - Example: The Theory of General Relativity
- **Law:** A concise description of a universally true aspect of the universe. Laws often take the form of mathematical equations.
 - Example: Newton's Second Law of Motion ($F = ma$)

Physical Quantities and Units (Sec 1.3)

- A **physical quantity** is a property of an object that can be measured.
 - Examples: length, mass, time, velocity.

Physical Quantities and Units (Sec 1.3)

- A **physical quantity** is a property of an object that can be measured.
 - Examples: length, mass, time, velocity.
- We will use the **SI (Système International) units**, which are part of the metric system.
- The four fundamental units for this course are:
 - Length: **meter (m)**
 - Mass: **kilogram (kg)**
 - Time: **second (s)**
 - Electric Current: **ampere (A)**

Physical Quantities and Units (Sec 1.3)

- A **physical quantity** is a property of an object that can be measured.
 - Examples: length, mass, time, velocity.
- We will use the **SI (Système International) units**, which are part of the metric system.
- The four fundamental units for this course are:
 - Length: **meter (m)**
 - Mass: **kilogram (kg)**
 - Time: **second (s)**
 - Electric Current: **ampere (A)**
- The metric system uses powers of 10 for easy conversion between different scales (e.g., 1 kilometer = 10^3 meters).

Accuracy and Precision (Context)

In physics, measurements are key. Two terms describe the quality of measurements:

- **Accuracy:** How close a measurement is to the true or accepted value.
 - Are your measurements correct?

Accuracy and Precision (Context)

In physics, measurements are key. Two terms describe the quality of measurements:

- **Accuracy:** How close a measurement is to the true or accepted value.
 - Are your measurements correct?
- **Precision:** How close a series of measurements are to each other.
 - Are your measurements repeatable?

An ideal measurement is both accurate and precise. Let's visualize this idea.

Visualizing Accuracy and Precision

[Image of four targets showing: (a) High accuracy, high precision; (b) Low accuracy, high precision; (c) High accuracy, low precision; (d) Low accuracy, low precision]

The classic target analogy for accuracy and precision.

Significant Figures (Sec 1.3)

Significant figures in a measurement express the precision of the measuring tool. When performing calculations, the result cannot be more precise than the least precise measurement. **Rules for Calculations:**

- **Multiplication/Division:** The final answer must have the same number of significant figures as the measurement with the **fewest significant figures**.

Significant Figures (Sec 1.3)

Significant figures in a measurement express the precision of the measuring tool. When performing calculations, the result cannot be more precise than the least precise measurement. **Rules for Calculations:**

- **Multiplication/Division:** The final answer must have the same number of significant figures as the measurement with the **fewest significant figures**.
- **Addition/Subtraction:** The final answer must have the same number of decimal places as the measurement with the **fewest decimal places**.

Distance vs. Displacement (Context)

Understanding the difference between distance and displacement is crucial for describing motion.

- **Distance (Scalar):** The total length of the path traveled. It has magnitude only.
 - "How much ground an object has covered."

Distance vs. Displacement (Context)

Understanding the difference between distance and displacement is crucial for describing motion.

- **Distance (Scalar):** The total length of the path traveled. It has magnitude only.
 - "How much ground an object has covered."
- **Displacement (Vector):** The change in an object's position. It has both magnitude and direction.
 - "How far out of place an object is."
 - Equation: $\Delta \vec{d} = \vec{d}_f - \vec{d}_0$ (final position - initial position)

Visualizing Distance and Displacement

[Diagram showing a winding path from point A to point B. The path length is labeled as 'distance'. A straight arrow from A to B is labeled as 'displacement'.]

Distance is the path taken; displacement is the straight-line change in position.

Speed vs. Velocity (Sec 2.2)

- **Average Speed (Scalar):** The rate at which distance is covered.
 - $v_{avg} = \frac{\text{distance}}{\text{time}}$

Speed vs. Velocity (Sec 2.2)

- **Average Speed (Scalar):** The rate at which distance is covered.
 - $v_{avg} = \frac{\text{distance}}{\text{time}}$
- **Velocity (Vector):** The rate of change of displacement. It includes speed *and* direction.
 - An object's velocity can change even if its speed is constant (e.g., a car turning a corner).

Speed vs. Velocity (Sec 2.2)

- **Average Speed (Scalar):** The rate at which distance is covered.
 - $v_{avg} = \frac{\text{distance}}{\text{time}}$
- **Velocity (Vector):** The rate of change of displacement. It includes speed *and* direction.
 - An object's velocity can change even if its speed is constant (e.g., a car turning a corner).
- **Average Velocity (Vector):** Displacement divided by the time interval.
 - $\vec{v}_{avg} = \frac{\Delta \vec{d}}{\Delta t} = \frac{\vec{d}_f - \vec{d}_0}{t_f - t_0}$

Position vs. Time Graphs (Context)

We can represent motion visually with graphs. For a position-time graph:

- The vertical axis (y-axis) is position (\vec{d}).
- The horizontal axis (x-axis) is time (t).

Position vs. Time Graphs (Context)

We can represent motion visually with graphs. For a position-time graph:

- The vertical axis (y-axis) is position (\vec{d}).
- The horizontal axis (x-axis) is time (t).
- The **slope** of the line gives the **velocity**.


- Slope = $\frac{\text{rise}}{\text{run}} = \frac{\Delta \vec{d}}{\Delta t} = \vec{v}$

Position vs. Time Graphs (Context)

We can represent motion visually with graphs. For a position-time graph:

- The vertical axis (y-axis) is position (\vec{d}).
- The horizontal axis (x-axis) is time (t).
- The **slope** of the line gives the **velocity**.
 - Slope = $\frac{\text{rise}}{\text{run}} = \frac{\Delta \vec{d}}{\Delta t} = \vec{v}$
- A **straight line** means constant velocity.
- A **horizontal line** means zero velocity (the object is at rest).
- A **curved line** means the velocity is changing (acceleration).

Visualizing Position vs. Time



Velocity vs. Time Graphs (Context)

For a velocity-time graph:

- The vertical axis (y-axis) is velocity (\vec{v}).
- The horizontal axis (x-axis) is time (t).

Velocity vs. Time Graphs (Context)

For a velocity-time graph:


- The vertical axis (y-axis) is velocity (\vec{v}).
- The horizontal axis (x-axis) is time (t).
- The **slope** of the line gives the **acceleration**.
 - Slope = $\frac{\text{rise}}{\text{run}} = \frac{\Delta \vec{v}}{\Delta t} = \vec{a}$

Velocity vs. Time Graphs (Context)

For a velocity-time graph:

- The vertical axis (y-axis) is velocity (\vec{v}).
- The horizontal axis (x-axis) is time (t).
- The **slope** of the line gives the **acceleration**.
 - Slope = $\frac{\text{rise}}{\text{run}} = \frac{\Delta \vec{v}}{\Delta t} = \vec{a}$
- The **area** under the line gives the **displacement**.
 - Area = height \times width = $\vec{v} \times t = \Delta \vec{d}$

Visualizing Velocity vs. Time



Acceleration (Sec 3.1)

- **Acceleration** is the rate at which an object's velocity changes.
- It is a **vector** quantity, so it has both magnitude and direction.

Acceleration (Sec 3.1)

- **Acceleration** is the rate at which an object's velocity changes.
- It is a **vector** quantity, so it has both magnitude and direction.
- An object is accelerating if its:
 - speed is changing (speeding up or slowing down).
 - direction is changing.

Acceleration (Sec 3.1)

- **Acceleration** is the rate at which an object's velocity changes.
- It is a **vector** quantity, so it has both magnitude and direction.
- An object is accelerating if its:
 - speed is changing (speeding up or slowing down).
 - direction is changing.
- Average acceleration is calculated as:

$$\vec{a}_{avg} = \frac{\Delta \vec{v}}{\Delta t} = \frac{\vec{v}_f - \vec{v}_0}{t_f - t_0}$$

- The standard unit for acceleration is **meters per second squared (m/s^2)**.

Essential Equations (for constant acceleration)

These are the kinematic equations that relate displacement (d), time (t), initial velocity (v_0), final velocity (v), and acceleration (a).

① **Velocity from acceleration:**

$$v = v_0 + at$$

Use when you don't know displacement.

Essential Equations (for constant acceleration)

These are the kinematic equations that relate displacement (d), time (t), initial velocity (v_0), final velocity (v), and acceleration (a).

① Velocity from acceleration:

$$v = v_0 + at$$

Use when you don't know displacement.

② Displacement from time and velocity:

$$d = d_0 + v_0 t + \frac{1}{2}at^2$$

Use when you don't know final velocity.

Essential Equations (for constant acceleration)

These are the kinematic equations that relate displacement (d), time (t), initial velocity (v_0), final velocity (v), and acceleration (a).

① Velocity from acceleration:

$$v = v_0 + at$$

Use when you don't know displacement.

② Displacement from time and velocity:

$$d = d_0 + v_0 t + \frac{1}{2}at^2$$

Use when you don't know final velocity.

③ Velocity from displacement:

$$v^2 = v_0^2 + 2a(d - d_0)$$

Use when you don't know time.

Essential Equations (for constant acceleration)

These are the kinematic equations that relate displacement (d), time (t), initial velocity (v_0), final velocity (v), and acceleration (a).

① Velocity from acceleration:

$$v = v_0 + at$$

Use when you don't know displacement.

② Displacement from time and velocity:

$$d = d_0 + v_0 t + \frac{1}{2}at^2$$

Use when you don't know final velocity.

③ Velocity from displacement:

$$v^2 = v_0^2 + 2a(d - d_0)$$

Use when you don't know time.

④ Displacement from average velocity:

$$d = d_0 + \left(\frac{v_0 + v}{2} \right) t$$

Example Problem (I do)

Problem: A sports car accelerates from rest to 27 m/s in 9.0 s. What is its average acceleration?

E - Equation:

$$a = \frac{\Delta v}{\Delta t} = \frac{v - v_0}{\Delta t}$$

G - Givens:

- $v_0 = 0 \text{ m/s}$ ("from rest")
- $v = 27 \text{ m/s}$
- $\Delta t = 9.0 \text{ s}$

S - Substitute:

$$a = \frac{27 \text{ m/s} - 0 \text{ m/s}}{9.0 \text{ s}}$$

U - Unknown:

- $a = ?$

S - Solve:

$$a = 3.0 \text{ m/s}^2$$

Guided Practice (We do)

Problem: A dragster starts from rest and accelerates at a constant 10.0 m/s^2 for 4.0 s . How far does it travel in this time?

G - Givens:

- $v_0 = 0 \text{ m/s}$
- $a = 10.0 \text{ m/s}^2$
- $t = 4.0 \text{ s}$
- $d_0 = 0 \text{ m}$

U - Unknown:

- $d = ?$

Guided Practice (We do)

Problem: A dragster starts from rest and accelerates at a constant 10.0 m/s^2 for 4.0 s . How far does it travel in this time?

G - Givens:

- $v_0 = 0 \text{ m/s}$
- $a = 10.0 \text{ m/s}^2$
- $t = 4.0 \text{ s}$
- $d_0 = 0 \text{ m}$

U - Unknown:

- $d = ?$

E - Equation: Which equation should we use?

- A) $v = v_0 + at$
- B) $d = d_0 + v_0t + \frac{1}{2}at^2$
- C) $v^2 = v_0^2 + 2a(d - d_0)$

Guided Practice (We do)

Problem: A dragster starts from rest and accelerates at a constant 10.0 m/s^2 for 4.0 s . How far does it travel in this time?

G - Givens:

- $v_0 = 0 \text{ m/s}$
- $a = 10.0 \text{ m/s}^2$
- $t = 4.0 \text{ s}$
- $d_0 = 0 \text{ m}$

U - Unknown:

- $d = ?$

E - Equation: Which equation should we use?

- A) $v = v_0 + at$
- B) $d = d_0 + v_0t + \frac{1}{2}at^2$
- C) $v^2 = v_0^2 + 2a(d - d_0)$

Correct Answer: B. It has all our givens and our unknown.

Guided Practice (We do)

Problem: A dragster starts from rest and accelerates at a constant 10.0 m/s^2 for 4.0 s . How far does it travel in this time?

G - Givens:

- $v_0 = 0 \text{ m/s}$
- $a = 10.0 \text{ m/s}^2$
- $t = 4.0 \text{ s}$
- $d_0 = 0 \text{ m}$

U - Unknown:

- $d = ?$

E - Equation: Which equation should we use?

- A) $v = v_0 + at$
- B) $d = d_0 + v_0 t + \frac{1}{2}at^2$
- C) $v^2 = v_0^2 + 2a(d - d_0)$

Correct Answer: B. It has all our givens and our unknown.

S - Substitute:

Guided Practice (We do)

Problem: A dragster starts from rest and accelerates at a constant 10.0 m/s^2 for 4.0 s . How far does it travel in this time?

G - Givens:

- $v_0 = 0 \text{ m/s}$
- $a = 10.0 \text{ m/s}^2$
- $t = 4.0 \text{ s}$
- $d_0 = 0 \text{ m}$

U - Unknown:

- $d = ?$

E - Equation: Which equation should we use?

- A) $v = v_0 + at$
- B) $d = d_0 + v_0t + \frac{1}{2}at^2$
- C) $v^2 = v_0^2 + 2a(d - d_0)$

Correct Answer: B. It has all our givens and our unknown.

S - Substitute:

Independent Practice (You do)

Problem: A cyclist is traveling at 15 m/s . She applies the brakes, causing a constant negative acceleration, and comes to a stop in 3.0 s .

Part A: What was her acceleration?

Part B: How far did she travel while braking?

Use the GUESS method and the kinematic equations to solve.

Independent Practice (You do)

Problem: A cyclist is traveling at 15 m/s. She applies the brakes, causing a constant negative acceleration, and comes to a stop in 3.0 s.

Part A: What was her acceleration?

Part B: How far did she travel while braking?

Use the GUESS method and the kinematic equations to solve.

Answers:

- Part A: $a = -5.0 \text{ m/s}^2$
- Part B: $d = 22.5 \text{ m}$

Summary

- Physics describes the fundamental rules of the universe. We use the scientific method to uncover these rules.

Summary

- Physics describes the fundamental rules of the universe. We use the scientific method to uncover these rules.
- We distinguish between **scalar** quantities (distance, speed) and **vector** quantities (displacement, velocity, acceleration).

Summary

- Physics describes the fundamental rules of the universe. We use the scientific method to uncover these rules.
- We distinguish between **scalar** quantities (distance, speed) and **vector** quantities (displacement, velocity, acceleration).
- Motion graphs are powerful tools:
 - **Position-Time Graph:** Slope is velocity.
 - **Velocity-Time Graph:** Slope is acceleration, area is displacement.

Summary

- Physics describes the fundamental rules of the universe. We use the scientific method to uncover these rules.
- We distinguish between **scalar** quantities (distance, speed) and **vector** quantities (displacement, velocity, acceleration).
- Motion graphs are powerful tools:
 - **Position-Time Graph:** Slope is velocity.
 - **Velocity-Time Graph:** Slope is acceleration, area is displacement.
- For motion with **constant acceleration**, we can use the kinematic equations to predict the future state of an object.