

# PHYS12 CH:28.4-28.6

## Relativistic Mechanics

Mr. Gullo

November 21, 2025

# Learning Objectives

By the end of this lesson, you will be able to:

# Learning Objectives

By the end of this lesson, you will be able to:

- Apply the relativistic velocity addition formula.

# Learning Objectives

By the end of this lesson, you will be able to:

- Apply the relativistic velocity addition formula.
- Calculate relativistic momentum and compare it to classical momentum.

# Learning Objectives

By the end of this lesson, you will be able to:

- Apply the relativistic velocity addition formula.
- Calculate relativistic momentum and compare it to classical momentum.
- Define rest energy, total energy, and kinetic energy in relativistic terms.

# Learning Objectives

By the end of this lesson, you will be able to:

- Apply the relativistic velocity addition formula.
- Calculate relativistic momentum and compare it to classical momentum.
- Define rest energy, total energy, and kinetic energy in relativistic terms.
- Solve problems involving mass-energy equivalence ( $E = mc^2$ ).

# Progression and Recap

## From Part 1

- $\gamma = \frac{1}{\sqrt{1-v^2/c^2}}$

# Progression and Recap

## From Part 1

- $\gamma = \frac{1}{\sqrt{1-v^2/c^2}}$
- Time dilates ( $\Delta t = \gamma \Delta t_0$ )



# Progression and Recap

## From Part 1

- $\gamma = \frac{1}{\sqrt{1-v^2/c^2}}$
- Time dilates ( $\Delta t = \gamma \Delta t_0$ )
- Length contracts ( $L = L_0/\gamma$ )

# Progression and Recap

## From Part 1

- $\gamma = \frac{1}{\sqrt{1-v^2/c^2}}$
- Time dilates ( $\Delta t = \gamma \Delta t_0$ )
- Length contracts ( $L = L_0/\gamma$ )

## Today: Dynamics

- What happens when we push objects near the speed of light?

# Progression and Recap

## From Part 1

- $\gamma = \frac{1}{\sqrt{1-v^2/c^2}}$
- Time dilates ( $\Delta t = \gamma \Delta t_0$ )
- Length contracts ( $L = L_0/\gamma$ )

## Today: Dynamics

- What happens when we push objects near the speed of light?
- Does  $F = ma$  still work? (Spoiler: Not simply)

# Progression and Recap

## From Part 1

- $\gamma = \frac{1}{\sqrt{1-v^2/c^2}}$
- Time dilates ( $\Delta t = \gamma \Delta t_0$ )
- Length contracts ( $L = L_0/\gamma$ )

## Today: Dynamics

- What happens when we push objects near the speed of light?
- Does  $F = ma$  still work? (Spoiler: Not simply)
- How do we add velocities correctly?

## 28.4 The Limit of Galilean Relativity

Classically, if you run at  $u'$  on a train moving at  $v$ , your ground speed is simply:

$$u = v + u'$$

## 28.4 The Limit of Galilean Relativity

Classically, if you run at  $u'$  on a train moving at  $v$ , your ground speed is simply:

$$u = v + u'$$

### Problem

If a spaceship moves at  $0.8c$  and fires a probe forward at  $0.8c$ , classical physics says the probe moves at  $1.6c$ .

## 28.4 The Limit of Galilean Relativity

Classically, if you run at  $u'$  on a train moving at  $v$ , your ground speed is simply:

$$u = v + u'$$

### Problem

If a spaceship moves at  $0.8c$  and fires a probe forward at  $0.8c$ , classical physics says the probe moves at  $1.6c$ . **This violates the second postulate ( $c$  is the limit).**

# Essential Equation: Velocity Addition

## Relativistic Velocity Addition

$$u = \frac{v + u'}{1 + \frac{vu'}{c^2}}$$



# Essential Equation: Velocity Addition

## Relativistic Velocity Addition

$$u = \frac{v + u'}{1 + \frac{vu'}{c^2}}$$

- $u$ : Velocity relative to stationary frame.

# Essential Equation: Velocity Addition

## Relativistic Velocity Addition

$$u = \frac{v + u'}{1 + \frac{vu'}{c^2}}$$

- $u$ : Velocity relative to stationary frame.
- $v$ : Velocity of the moving frame (e.g., the ship).

# Essential Equation: Velocity Addition

## Relativistic Velocity Addition

$$u = \frac{v + u'}{1 + \frac{vu'}{c^2}}$$

- $u$ : Velocity relative to stationary frame.
- $v$ : Velocity of the moving frame (e.g., the ship).
- $u'$ : Velocity relative to the moving frame (e.g., the probe).

# Essential Equation: Velocity Addition

## Relativistic Velocity Addition

$$u = \frac{v + u'}{1 + \frac{vu'}{c^2}}$$

- $u$ : Velocity relative to stationary frame.
- $v$ : Velocity of the moving frame (e.g., the ship).
- $u'$ : Velocity relative to the moving frame (e.g., the probe).

Notice: If  $v \ll c$  and  $u' \ll c$ , the denominator  $\approx 1$ , giving back  $u \approx v + u'$ .

## Example: I Do - Velocity Addition

**Problem:** A spaceship travels away from Earth at  $v = 0.5c$ . It fires a missile forward at speed  $u' = 0.5c$  relative to the ship. What is the missile's speed relative to Earth?

# I Do: Velocity Addition - G & U

## G - Givens

- $v = 0.5c$  (Ship relative to Earth)

# I Do: Velocity Addition - G & U

## G - Givens

- $v = 0.5c$  (Ship relative to Earth)
- $u' = 0.5c$  (Missile relative to Ship)

# I Do: Velocity Addition - G & U

## G - Givens

- $v = 0.5c$  (Ship relative to Earth)
- $u' = 0.5c$  (Missile relative to Ship)

## U - Unknown

- $u = ?$  (Missile relative to Earth)



# I Do: Velocity Addition - Equation

## E - Equation

# I Do: Velocity Addition - Equation

## E - Equation

- Use the relativistic addition formula:

$$u = \frac{v + u'}{1 + \frac{vu'}{c^2}}$$

# I Do: Velocity Addition - Substitute & Solve

## S - Substitute

$$\bullet \quad u = \frac{0.5c + 0.5c}{1 + \frac{(0.5c)(0.5c)}{c^2}}$$

# I Do: Velocity Addition - Substitute & Solve

## S - Substitute

- $u = \frac{0.5c + 0.5c}{1 + \frac{(0.5c)(0.5c)}{c^2}}$
- $u = \frac{1.0c}{1 + 0.25}$

# I Do: Velocity Addition - Substitute & Solve

## S - Substitute

- $u = \frac{0.5c + 0.5c}{1 + \frac{(0.5c)(0.5c)}{c^2}}$

- $u = \frac{1.0c}{1 + 0.25}$

## S - Solve

- $u = \frac{1.0c}{1.25} = 0.8c$

# I Do: Velocity Addition - Substitute & Solve

## S - Substitute

- $$u = \frac{0.5c + 0.5c}{1 + \frac{(0.5c)(0.5c)}{c^2}}$$

- $$u = \frac{1.0c}{1 + 0.25}$$

## S - Solve

- $$u = \frac{1.0c}{1.25} = 0.8c$$

- $$u = 0.8c$$

# I Do: Velocity Addition - Substitute & Solve

## S - Substitute

- $u = \frac{0.5c + 0.5c}{1 + \frac{(0.5c)(0.5c)}{c^2}}$

- $u = \frac{1.0c}{1 + 0.25}$

## S - Solve

- $u = \frac{1.0c}{1.25} = 0.8c$

- $u = 0.8c$

- *Result is less than  $c$ , as required.*

## 28.5 Relativistic Momentum

Newtonian momentum  $p = mv$  is not conserved at high speeds. We must adjust the definition.



## 28.5 Relativistic Momentum

Newtonian momentum  $p = mv$  is not conserved at high speeds. We must adjust the definition.

### Relativistic Momentum

$$p = \gamma mv = \frac{mv}{\sqrt{1 - \frac{v^2}{c^2}}}$$

## 28.5 Relativistic Momentum

Newtonian momentum  $p = mv$  is not conserved at high speeds. We must adjust the definition.

### Relativistic Momentum

$$p = \gamma mv = \frac{mv}{\sqrt{1 - \frac{v^2}{c^2}}}$$

- As  $v \rightarrow c$ ,  $\gamma \rightarrow \infty$ , so  $p \rightarrow \infty$ .

## 28.5 Relativistic Momentum


Newtonian momentum  $p = mv$  is not conserved at high speeds. We must adjust the definition.

### Relativistic Momentum


$$p = \gamma mv = \frac{mv}{\sqrt{1 - \frac{v^2}{c^2}}}$$

- As  $v \rightarrow c$ ,  $\gamma \rightarrow \infty$ , so  $p \rightarrow \infty$ .
- This explains why a massive object cannot reach  $c$ : it would require infinite momentum (and infinite energy).

# Concept Visualization: Momentum Limit




# Concept Visualization: Momentum Limit



- **Classical line:** Straight line ( $p = mv$ ).

# Concept Visualization: Momentum Limit



- **Classical line:** Straight line ( $p = mv$ ).
- **Relativistic curve:** Follows classical line at low speeds, then curves upward sharply, approaching a vertical asymptote at  $v = c$ .

## 28.6 Rest Energy and Total Energy

Einstein's most famous equation relates mass and energy.

## 28.6 Rest Energy and Total Energy

Einstein's most famous equation relates mass and energy.

### Rest Energy ( $E_0$ )

The energy an object has simply because it has mass.

$$E_0 = mc^2$$



## 28.6 Rest Energy and Total Energy

Einstein's most famous equation relates mass and energy.

### Rest Energy ( $E_0$ )

The energy an object has simply because it has mass.

$$E_0 = mc^2$$

### Total Energy ( $E$ )

The sum of rest energy and kinetic energy.

$$E = \gamma mc^2 = \frac{mc^2}{\sqrt{1 - \frac{v^2}{c^2}}}$$

# Relativistic Kinetic Energy

Kinetic Energy is the "extra" energy due to motion.

$$E = E_0 + KE$$

# Relativistic Kinetic Energy

Kinetic Energy is the "extra" energy due to motion.

$$E = E_0 + KE$$

Rearranging for KE:

$$KE = E - E_0 = \gamma mc^2 - mc^2$$

# Relativistic Kinetic Energy

Kinetic Energy is the "extra" energy due to motion.

$$E = E_0 + KE$$

Rearranging for KE:

$$KE = E - E_0 = \gamma mc^2 - mc^2$$

## Relativistic Kinetic Energy

$$KE = (\gamma - 1)mc^2$$

# Relativistic Kinetic Energy

Kinetic Energy is the "extra" energy due to motion.

$$E = E_0 + KE$$

Rearranging for KE:

$$KE = E - E_0 = \gamma mc^2 - mc^2$$

## Relativistic Kinetic Energy

$$KE = (\gamma - 1)mc^2$$

*Note: At low speeds, this simplifies to  $\frac{1}{2}mv^2$ .*

# We Do: Relativistic Energy

**Problem:** An electron ( $m = 9.11 \times 10^{-31}$  kg) is accelerated to  $0.999c$ . What is its total energy?

# We Do: Relativistic Energy

**Problem:** An electron ( $m = 9.11 \times 10^{-31}$  kg) is accelerated to  $0.999c$ . What is its total energy? **Steps:**

- 1 Calculate  $\gamma$  for  $v = 0.999c$ .

# We Do: Relativistic Energy

**Problem:** An electron ( $m = 9.11 \times 10^{-31}$  kg) is accelerated to  $0.999c$ . What is its total energy? **Steps:**

- 1 Calculate  $\gamma$  for  $v = 0.999c$ .
- 2 Use  $E = \gamma mc^2$ .



# We Do: Relativistic Energy

**Problem:** An electron ( $m = 9.11 \times 10^{-31}$  kg) is accelerated to  $0.999c$ . What is its total energy? **Steps:**

- 1 Calculate  $\gamma$  for  $v = 0.999c$ .
- 2 Use  $E = \gamma mc^2$ .
- 3 (Optional) Convert Joules to eV or MeV.

# You Do: Practice

**Problem:** A proton has a rest energy of 938 MeV. It is moving such that its total energy is 2000 MeV.

# You Do: Practice

**Problem:** A proton has a rest energy of 938 MeV. It is moving such that its total energy is 2000 MeV.

- 1 What is its kinetic energy? ( $KE = E - E_0$ )

# You Do: Practice

**Problem:** A proton has a rest energy of 938 MeV. It is moving such that its total energy is 2000 MeV.

- 1 What is its kinetic energy? ( $KE = E - E_0$ )
- 2 What is the value of  $\gamma$ ?

# You Do: Practice

**Problem:** A proton has a rest energy of 938 MeV. It is moving such that its total energy is 2000 MeV.

- 1 What is its kinetic energy? ( $KE = E - E_0$ )
- 2 What is the value of  $\gamma$ ?
- 3 How fast is it moving? (Solve  $\gamma$  for  $v$ ).

# Summary

- **Velocity Addition:** Velocities do not add linearly;  $c$  is the max speed.

# Summary

- **Velocity Addition:** Velocities do not add linearly;  $c$  is the max speed.
- **Momentum:**  $p = \gamma mv$ . Infinite momentum is required to reach  $c$ .



# Summary

- **Velocity Addition:** Velocities do not add linearly;  $c$  is the max speed.
- **Momentum:**  $p = \gamma mv$ . Infinite momentum is required to reach  $c$ .
- **Mass-Energy:** Mass is a form of energy ( $E_0 = mc^2$ ).

# Summary

- **Velocity Addition:** Velocities do not add linearly;  $c$  is the max speed.
- **Momentum:**  $p = \gamma mv$ . Infinite momentum is required to reach  $c$ .
- **Mass-Energy:** Mass is a form of energy ( $E_0 = mc^2$ ).
- **Total Energy:**  $E = \gamma mc^2$ .

# Summary

- **Velocity Addition:** Velocities do not add linearly;  $c$  is the max speed.
- **Momentum:**  $p = \gamma mv$ . Infinite momentum is required to reach  $c$ .
- **Mass-Energy:** Mass is a form of energy ( $E_0 = mc^2$ ).
- **Total Energy:**  $E = \gamma mc^2$ .
- **Kinetic Energy:**  $KE = (\gamma - 1)mc^2$ .