



MRC Cognition
and Brain
Sciences Unit



UNIVERSITY OF
CAMBRIDGE

Diffusion MRI: Introduction to Tractography


Marta M. Correia
MRC Cognition and Brain Sciences Unit

Overview


- Introduction to Tractography
- Probabilistic Modelling
- Multiple-fibre models

Diffusion ellipsoid: Eigenvalues and eigenvectors

Isotropic Diffusion




Anisotropic Diffusion



Colour coded FA maps

- Let ε_1 designate the longest axis of the diffusion ellipsoid.
- ε_1 can be identified with the main direction of diffusion.
- This directional information can be added to the FA map using a colour code:



Red indicates directions in the x axis: right to left or left to right.

Green indicates directions in the y axis: front to back or back to front.


Blue indicates directions in the z axis: foot-to-head direction or vice versa.

Colour coded FA map.




Introduction to Tractography

Tractography: basic principles

- After model fitting, the principal direction ϵ_1 can be calculated for all voxels.
- The trajectories of water molecules can be reconstructed using a method similar to the children's activity "connect the dots".
- We connect each voxel to the adjacent one toward which the fibre direction, ϵ_1 , is pointing.



Tractography in the Brain



Fibre tracks obtained for a dataset of a healthy volunteer using simple streamlining (FACT).


Probabilistic Modeling of Diffusion MRI Signal

Probabilistic Modelling

- The information provided by DTI can be very useful for the characterisation of brain white matter.
- However, the estimated tensor can be highly dependent on noise.
- Probabilistic modelling can be used to estimate a probability distribution function for the DTI model parameters.
- For the fibre orientation, this distribution is called orientation distribution function (ODF).


Orientation Distribution Function (ODF)

Anisotropic Diffusion



Orientation Distribution Function (ODF)

Anisotropic Diffusion




ODF calculated from the
displacement probability
distribution



For DTI:


$$ODF(\mathbf{n}) = \frac{1}{4\pi |\mathbf{D}|^{1/2}} \left(\frac{1}{\mathbf{n}^T \mathbf{D}^{-1} \mathbf{n}} \right)^{3/2}$$



Rafael Neto Henriques
Pybrain workshop 2020

Orientation Distribution Function (ODF)

Anisotropic Diffusion




ODF calculated from the
displacement probability
distribution



For DTI:


$$ODF(\mathbf{n}) = \frac{1}{4\pi |\mathbf{D}|^{1/2}} \left(\frac{1}{\mathbf{n}^T \mathbf{D}^{-1} \mathbf{n}} \right)^{3/2}$$



Rafael Neto Henriques
Pybrain workshop 2020

Orientation Distribution Function (ODF)


- Example: 10 samples extracted from the ODF of each voxel





- Regions of one-fibre populations have very narrow distributions, while regions of crossing fibres show greater variability.

Probabilistic Tractography

- For each sample of the directional PDF we can produce a track (or streamline).
- Repeat this for a large number of samples.
- The probability of voxels A and B being connected can be calculated by dividing the number of streamlines that reach B, by the total number of streamlines generated from A.



Probabilistic Tractography in the Brain



Probabilistic tractography dataset obtained for a healthy volunteer.

Multiple fibres

Beyond the Diffusion Tensor

Multiple fibre approaches

Model-Based Approaches

- The multi-tensor model

Non-Parametric Approaches

- Diffusion Spectrum Imaging (DSI)
- Q-ball Imaging
- Constrained Spherical Deconvolution (CSD)
- Persistent Angular Structure (PAS)

Multiple fibre approaches

Model-Based Approaches

- The multi-tensor model

Non-Parametric Approaches

- Diffusion Spectrum Imaging (DSI)
- Q-ball Imaging
- Constrained Spherical Deconvolution (CSD)
- Persistent Angular Structure (PAS)

Constrained Spherical Deconvolution (CSD)

- CSD is able to estimate the distribution of fibre orientations within a voxel without making assumptions about the number of fibres present.
- The original method (Tournier et al. 2007) requires a single-shell high angular resolution diffusion data (>60 directions).
- Ideally, the b -value used should be in the region of 2,500 – 3,000 s/mm² (for *in vivo* human brains).
- Multi-shell multi-tissue CSD (MSMT-CSD) was introduced by Jeurissen et al. in 2014.
- MSMT-CSD requires multi-shell high angular resolution diffusion data, containing multiple b -values.
- To resolve WM, GM & CSF the acquisition should contain at least 2 shells plus the $b=0$ volumes (i.e. 3 unique b -values).

Constrained Spherical Deconvolution (CSD)

Tournier et al. 2007

$$f_1 S_1(\theta, \phi) + f_2 S_2(\theta, \phi) = S(\theta, \phi) = R(\theta) \otimes F(\theta, \phi)$$



- Two distinct fibre orientations
- $S_1(\theta, \phi)$, $S_2(\theta, \phi)$ – diffusion weighted signal attenuation
- f_1, f_2 – volume fractions

- Overall diffusion weighted signal attenuation


- Axially symmetric response function describing the signal attenuation measured for a single fibre population.

- Fibre orientation density function describing the fibre orientations present in the voxel

Constrained Spherical Deconvolution (CSD)




Constrained Spherical Deconvolution (CSD)



**DTI
Tractography**



**CSD Deterministic
Tractography**



**CSD Probabilistic
Tractography**

From Gabriel Girard, DIPY tutorial 2021


Creating the Connectome

- Parcellate the brain using Freesurfer
- Apply parcellation to the tractogram to generate a connectivity matrix, representing the number of streamlines connecting each pair of ROIs



Interpretation of results

- Streamlines generated with diffusion MRI tractography lack polarity
- Even the best fiber orientation reconstruction algorithm cannot resolve all crossing fibers
- Structural connectomes are dominated by false positives
- Choice of fitting model, tracking algorithm, stopping criteria, etc, can also influence the connectome





MRC Cognition
and Brain
Sciences Unit



UNIVERSITY OF
CAMBRIDGE

Questions?