



Mission: Iconic Reefs and National Coral Reef Monitoring Program Benthic Analyses

Nicole Krampitz, Dione Swanson, Julia Mateski, Lexie Sturm, and Shay Viehman

2025-09-26

Table of Contents

1	Introduction	3
2	Benthic Survey Effort	5
2.1	Coral Demography	5
2.2	Benthic Composition	6
3	Benthic Community Composition	7
4	Coral Demography	11
4.1	Focal Species	12
4.2	Coral Density	22
4.3	Old Mortality	23
4.4	Coral Size	24
4.5	Bleaching Estimates	25
4.6	Disease Estimates	26
5	Point of Contact	27
6	References	27

1 Introduction

Mission: Iconic Reefs (M:IR) is a multi-institutional initiative designed to restore ecological function and biodiversity across key reef sites in the Florida Keys. Led by NOAA in collaboration with federal, state, institutional, and non-profit partners, M:IR targets nearly three million square feet of reef at seven key coral reefs in the Florida Keys National Marine Sanctuary (FKNMS): Carysfort (North and South), Cheeca Rocks, Eastern Dry Rocks, Horseshoe Reef, Looe Key, Newfound Harbor, and Sombrero Reef (Figure 1). Across these reefs, M:IR aims to restore coral cover to an average of 25% across the seven sites – an initiative unparalleled in scope and scale that will require hundreds of thousands of coral restoration outplants and over a decade of phased intervention.




Figure 1: Map of the Florida Keys showing the sites of Mission: Iconic Reefs

These Iconic reefs were selected for restoration due to their diversity in reef types, levels of anthropogenic disturbances, initial coral cover, geographic distribution, and cultural significance. Restoration progress at these reefs is informed by ongoing monitoring of coral reef benthic communities and coral populations at M:IR sites using a range of metrics, such as coral cover, size structure, mortality prevalence, and diversity. Such multi-metric monitoring is essential to evaluating whether current restoration practices are effective in establishing and maintaining a healthy coral reef, as restoration ‘success’ encompasses a range of ecological

and management metrics rather than a single measurement (e.g., coral cover) ([Goergen et al., 2020](#)).

In addition to monitoring within the M:IR reefs, it is also important to assess the surrounding reefs in the Florida Keys to interpret restoration progress. The National Coral Reef Monitoring Program (NCRMP), which conducts standardized reef assessments across the Florida Keys on a biennial basis, provides a solid framework for comparison. To align with this program, the M:IR team applied NCRMP survey methodology within its Iconic Sites in 2022 and 2024 to complement NCRMP surveys. These surveys focused on shallow (<30 m), hard-bottom reef habitats, using a stratified-random, one-stage design within 50 m × 50 m grid cells to ensure representative sampling across depth and rugosity strata ([Ault et al., 2021](#)). The survey design ensures that survey sites are allocated by cross-shelf zone, rugosity type, and depth strata, with sites distributed around the Florida Keys from nearshore to offshore to a maximum depth of 30 m. The NCRMP sample grid was spatially joined with the seven distinct M:IR zones (Figure 1). Survey sites were proportionally allocated within NCRMP strata to maintain consistent representation across habitats.

Corals and benthic communities were monitored using two different NCRMP surveys: the Benthic Community Assessment survey and the Coral Demographics survey (NOAA CRCP 2022a, 2022b, 2024a, 2024b). The Benthic Community Assessment survey includes: 1) benthic cover (%) estimates along a 15-m transect with a line point intercept method, 2) presence/absence of Endangered Species Act (ESA)-listed coral species ([NOAA National Marine Fisheries Service, 2014](#)), 3) abundance of key macroinvertebrates, and 4) reef rugosity measurements within a 15 m × 2 m belt-transect area (NOAA CRCP 2022a, 2024b). At the same site, coral demographics were surveyed within a 10 m × 1 m belt-transect area (NOAA CRCP 2022b, 2024b). NCRMP coral demographic survey data were combined with complementarily allocated survey data from Florida Fish and Wildlife Conservation Commission's Disturbance Response Monitoring ([DRM](#)). In all coral demographics surveys, all live coral colonies □ 4 cm were counted, identified to species, measured to the nearest centimeter, and estimates were made of the proportion per colony of any present mortality (recent or old), disease (absent, present: slow, fast), and/or bleaching (none, total, partial, paling). Only live coral colonies were included in the survey; dead colonies with 100% mortality were not surveyed (e.g., colonies killed by coral disease). Juvenile corals (< 4 cm) were reported for species richness only and were not included in counts, size measurements, or estimates of condition (NOAA CRCP 2022b); in 2024, juveniles of select coral species were counted (NOAA CRCP 2024b). In 2024, NCRMP benthic surveys also included large area imaging (LAI) collected at most survey sites; analyses of these LAI data are in progress and not presented here.

To allow for direct comparisons of benthic communities inside the M:IR restoration areas with control, non-restored areas across the Florida Keys reefs, the NCRMP dataset was restricted to strata types and depth zones (0-12 m) found within M:IR sites. Statistical comparisons were conducted using a two-tailed t-test between M:IR and NCRMP estimates for each survey year (2022 or 2024) and between years for M:IR and NCRMP. This report focuses on coral

populations and benthic community metrics; a separate [report](#) can be found for fish population assessments ([Blondeau et al., 2025](#)).

NCRMP analyses scripts for corals and benthic communities are open source and available at [NCRMP Benthic R package](#) ([Groves et al., 2025](#)).

2 Benthic Survey Effort

In 2022, a total of 584 surveys occurred (Table 1) across 403 sites ([Viehman et al., 2023](#)). In 2024, 771 surveys occurred across 542 sites ([Grove et al., in prep](#)).

Table 1. Number of demographic (Demo) and benthic community assessment (BCA) surveys completed by M:IR, NCRMP, and DRM in 2022 and 2024.

	M:IR		NCRMP		DRM
Year	Demo	BCA	Demo	BCA	Demo
2022	90	89	95	92	218
2024	100	100	129	129	313
Total	379		445		531

2.1 Coral Demography

For the purpose of this comparison, surveys were divided into two groups: inside M:IR and outside. Almost all of the inside (M:IR) surveys were completed by the M:IR team; however, if an NCRMP or DRM sample happened to fall within the boundaries of an M:IR iconic reef, it was added to the inside group. The inverse occurred if an M:IR survey happened to fall outside the M:IR boundary.

Table 2. The number of coral demographic sites sampled inside and outside M:IR areas in each stratum (includes DRM data).

Study Area	Strata	Strata Description	2022	2024
Inside (M:IR)	CFK01	Inshore reef, all relief types and depths	12	10
Inside (M:IR)	CFK02	Mid-channel patch reef, all relief types and depths	16	12
Inside (M:IR)	CFK03	Offshore patch reef, all relief types and depths	7	11
Inside (M:IR)	CFK04	Forereef, low relief, shallow (0-6 m)	—	4

(continued)

Study Area	Strata	Strata Description	2022	2024
Inside (M:IR)	CFK05	Forereef, high relief, shallow (0-6 m)	25	38
Inside (M:IR)	CFK06	Forereef, low relief, mid-shallow (6-12 m)	4	12
Inside (M:IR)	CFK07	Forereef, high relief, mid-shallow (6-12 m)	34	34
Outside (NCRMP + DRM)	CFK01	Inshore reef, all relief types and depths	27	24
Outside (NCRMP + DRM)	CFK02	Mid-channel patch reef, all relief types and depths	79	109
Outside (NCRMP + DRM)	CFK03	Offshore patch reef, all relief types and depths	37	42
Outside (NCRMP + DRM)	CFK04	Forereef, low relief, shallow (0-6 m)	4	31
Outside (NCRMP + DRM)	CFK05	Forereef, high relief, shallow (0-6 m)	21	45
Outside (NCRMP + DRM)	CFK06	Forereef, low relief, mid-shallow (6-12 m)	19	33
Outside (NCRMP + DRM)	CFK07	Forereef, high relief, mid-shallow (6-12 m)	60	61

2.2 Benthic Composition

Disturbance Response Monitoring (DRM) only performs coral demographic surveys (Table 1) and therefore DRM surveys are not included in the summary of Benthic Community Assessment surveys (Table 3).

Table 3. Number of Benthic Community Assessment (BCA) sites sampled inside and outside M:IR areas in each stratum.

Study Area	Strat	Strat Description	2022	2024
Inside (M:IR)	CFK01	Inshore reef, all relief types and depths	8	10
Inside (M:IR)	CFK02	Mid-channel patch reef, all relief types and depths	12	12
Inside (M:IR)	CFK03	Offshore patch reef, all relief types and depths	7	8
Inside (M:IR)	CFK04	Forereef, low relief, shallow (0-6 m)	—	3
Inside (M:IR)	CFK05	Forereef, high relief, shallow (0-6 m)	24	33
Inside (M:IR)	CFK06	Forereef, low relief, mid-shallow (6-12 m)	4	9
Inside (M:IR)	CFK07	Forereef, high relief, mid-shallow (6-12 m)	32	28
Outside (NCRMP)	CFK01	Inshore reef, all relief types and depths	10	9
Outside (NCRMP)	CFK02	Mid-channel patch reef, all relief types and depths	21	13
Outside (NCRMP)	CFK03	Offshore patch reef, all relief types and depths	9	8
Outside (NCRMP)	CFK04	Forereef, low relief, shallow (0-6 m)	—	6
Outside (NCRMP)	CFK05	Forereef, high relief, shallow (0-6 m)	7	9
Outside (NCRMP)	CFK06	Forereef, low relief, mid-shallow (6-12 m)	9	7
Outside (NCRMP)	CFK07	Forereef, high relief, mid-shallow (6-12 m)	10	28

3 Benthic Community Composition

Temporal trends in benthic cover of functional groups such as coral and macroalgae cover provide information on changes to the benthic community composition over time.




Figure 2. Cover estimates for main benthic groups inside and outside M:IR areas for 2022 and 2024. A black plus (+) indicates a significant difference between groups in that survey year (i.e., 2022 or 2024). A colored asterisk (*) indicates significance difference between years for the given group (i.e., M:IR or NCRMP).




Figure 2. Cover estimates for main benthic groups inside and outside M:IR areas for 2022 and 2024. A black plus (+) indicates a significant difference between groups in that survey year (i.e., 2022 or 2024). A colored asterisk (*) indicates significance difference between years for the given group (i.e., M:IR or NCRMP).




Figure 2. Cover estimates for main benthic groups inside and outside M:IR areas for 2022 and 2024. A black plus (+) indicates a significant difference between groups in that survey year (i.e., 2022 or 2024). A colored asterisk (*) indicates significance difference between years for the given group (i.e., M:IR or NCRMP).




Figure 2. Cover estimates for main benthic groups inside and outside M:IR areas for 2022 and 2024. A black plus (+) indicates a significant difference between groups in that survey year (i.e., 2022 or 2024). A colored asterisk (*) indicates significance difference between years for the given group (i.e., M:IR or NCRMP).

4 Coral Demography

Although several metrics reported here encompass all observed coral species, ten focal species are highlighted as particularly important to M:IR's goal: *Acropora palmata*, *Acropora cervicornis*, *Colpophyllia natans*, *Dendrogyra cylindrus*, *Dichocoenia stokesii*, *Diploria labyrinthiformis*, *Meandrina meandrites*, *Montastraea cavernosa*, *Pseudodiploria clivosa*, and *Pseudodiploria strigosa*. These species have experienced drastic population declines and are targeted for restoration at the M:IR sites. Specifically, most were severely affected by White Band Disease (*Acropora* spp.) or Stony Coral Tissue Loss Disease (SCTLD).

The following table summarizes the number of observations across all strata and survey years of each focal species and identifies the most abundant species encountered.

Table 4. Number of colonies observed for focal M:IR species and the most commonly encountered non-focal species in 2022 and 2024.

Species	Colonies in 2022	Colonies in 2024
<i>Siderastrea siderea</i>	3,868	5,618
<i>Porites astreoides</i>	2,657	2,132
<i>Stephanocoenia intersepta</i>	1,345	1,534
<i>Agaricia agaricites</i>	884	237
<i>Porites porites</i>	776	558
* <i>Montastraea cavernosa</i>	437	645
<i>Orbicella faveolata</i>	266	237
<i>Orbicella annularis</i>	211	156
<i>Siderastrea radians</i>	182	328
<i>Porites furcata</i>	160	77
<i>Porites divaricata</i>	118	100
* <i>Dichocoenia stokesii</i>	106	137
* <i>Colpophyllia natans</i>	103	76
<i>Solenastrea bournoni</i>	79	113
* <i>Acropora cervicornis</i>	74	13
* <i>Diploria labyrinthiformis</i>	50	65
* <i>Pseudodiploria strigosa</i>	50	46
<i>Eusmilia fastigiata</i>	45	75
* <i>Pseudodiploria clivosa</i>	23	31
* <i>Acropora palmata</i>	14	0
* <i>Meandrina meandrites</i>	5	11

4.1 Focal Species

The stratified random surveys are designed to optimize sampling efficiency and provide high precision estimates of coral population metrics at relatively low sample sizes ([Smith et al., 2011](#)). The effectiveness of restoration can be quantified by comparison of these metrics between M:IR sites and similar habitats outside these restoration areas.

The following figures for these metrics are grouped by each focal species. Significant statistical differences using a two-tailed t-test are indicated for differences between survey years (2022 vs. 2024) and between areas (M:IR vs. NCRMP) for density, old mortality, and mean colony size.

Acropora cervicornis

Colony density, mortality, size, and size class distribution (2022 vs. 2024)




Figure 3. Coral density, old mortality, colony size, and size frequency distribution for focal M:IR species across sample years (2022 and 2024) and survey types (M:IR and NCRMP + DRM). A black plus (+) indicates a significant difference in cover estimates between groups in that survey year (i.e., 2022 or 2024). A colored asterisk (*) indicates significance between years for the given group (i.e., M:IR or NCRMP). Note: Axis bounds vary by species.

Acropora palmata

Colony density, mortality, size, and size class distribution (2022 vs. 2024)




Figure 3. Coral density, old mortality, colony size, and size frequency distribution for focal M:IR species across sample years (2022 and 2024) and survey types (M:IR and NCRMP + DRM). A black plus (+) indicates a significant difference in cover estimates between groups in that survey year (i.e., 2022 or 2024). A colored asterisk (*) indicates significance between years for the given group (i.e., M:IR or NCRMP). Note: Axis bounds vary by species.

Colpophyllia natans

Colony density, mortality, size, and size class distribution (2022 vs. 2024)




Figure 3. Coral density, old mortality, colony size, and size frequency distribution for focal M:IR species across sample years (2022 and 2024) and survey types (M:IR and NCRMP + DRM). A black plus (+) indicates a significant difference in cover estimates between groups in that survey year (i.e., 2022 or 2024). A colored asterisk (*) indicates significance between years for the given group (i.e., M:IR or NCRMP). Note: Axis bounds vary by species.

Dichocoenia stokesii

Colony density, mortality, size, and size class distribution (2022 vs. 2024)





Figure 3. Coral density, old mortality, colony size, and size frequency distribution for focal M:IR species across sample years (2022 and 2024) and survey types (M:IR and NCRMP + DRM). A black plus (+) indicates a significant difference in cover estimates between groups in that survey year (i.e., 2022 or 2024). A colored asterisk (*) indicates significance between years for the given group (i.e., M:IR or NCRMP). Note: Axis bounds vary by species.

Diploria labyrinthiformis

Colony density, mortality, size, and size class distribution (2022 vs. 2024)



Size Frequency Distributions




Figure 3. Coral density, old mortality, colony size, and size frequency distribution for focal M:IR species across sample years (2022 and 2024) and survey types (M:IR and NCRMP + DRM). A black plus (+) indicates a significant difference in cover estimates between groups in that survey year (i.e., 2022 or 2024). A colored asterisk (*) indicates significance between years for the given group (i.e., M:IR or NCRMP). Note: Axis bounds vary by species.

Meandrina meandrites

Colony density, mortality, size, and size class distribution (2022 vs. 2024)




Figure 3. Coral density, old mortality, colony size, and size frequency distribution for focal M:IR species across sample years (2022 and 2024) and survey types (M:IR and NCRMP + DRM). A black plus (+) indicates a significant difference in cover estimates between groups in that survey year (i.e., 2022 or 2024). A colored asterisk (*) indicates significance between years for the given group (i.e., M:IR or NCRMP). Note: Axis bounds vary by species.

Pseudodiploria clivosa

Colony density, mortality, size, and size class distribution (2022 vs. 2024)





Figure 3. Coral density, old mortality, colony size, and size frequency distribution for focal M:IR species across sample years (2022 and 2024) and survey types (M:IR and NCRMP + DRM). A black plus (+) indicates a significant difference in cover estimates between groups in that survey year (i.e., 2022 or 2024). A colored asterisk (*) indicates significance between years for the given group (i.e., M:IR or NCRMP). Note: Axis bounds vary by species.

Pseudodiploria strigosa

Colony density, mortality, size, and size class distribution (2022 vs. 2024)



Size Frequency Distributions




Figure 3. Coral density, old mortality, colony size, and size frequency distribution for focal M:IR species across sample years (2022 and 2024) and survey types (M:IR and NCRMP + DRM). A black plus (+) indicates a significant difference in cover estimates between groups in that survey year (i.e., 2022 or 2024). A colored asterisk (*) indicates significance between years for the given group (i.e., M:IR or NCRMP). Note: Axis bounds vary by species.

Montastraea cavernosa

Colony density, mortality, size, and size class distribution (2022 vs. 2024)




Figure 3. Coral density, old mortality, colony size, and size frequency distribution for focal M:IR species across sample years (2022 and 2024) and survey types (M:IR and NCRMP + DRM). A black plus (+) indicates a significant difference in cover estimates between groups in that survey year (i.e., 2022 or 2024). A colored asterisk (*) indicates significance between years for the given group (i.e., M:IR or NCRMP). Note: Axis bounds vary by species.

4.2 Coral Density




Figure 4. Species-specific coral density (colonies/m²) outside (NCRMP and DRM) and inside M:IR areas in 2022 and 2024. Focal M:IR species are denoted with a (*).

4.3 Old Mortality




Figure 5. Percent old mortality (\pm SEM) by coral species outside (NCRMP and DRM) and inside M:IR areas in 2022 and 2024. Focal M:IR species are denoted with a (*).

4.4 Coral Size




Figure 6. Maximum diameter (cm) \pm SEM by coral species outside (NCRMP and DRM) and inside M:IR areas in 2022 and 2024. Focal M:IR species are denoted with a (*).

4.5 Bleaching Estimates




Figure 7. Species-specific bleaching frequency observed outside (NCRMP + DRM) and inside M:IR areas. Focal M:IR species are denoted with a (*).

4.6 Disease Estimates




Figure 8. Species-specific disease frequency observed outside (NCRMP + DRM) and inside M:IR areas. Focal M:IR species are denoted with a (*).

5 Point of Contact

NCRMP Atlantic Benthic Team Lead: **Shay Viehman** (shay.viehman@noaa.gov)

6 References

- Ault, J. S., Smith, S. G., Luo, J., Grove, L. J., Johnson, M. W., & Blondeau, J. (2021). *Refinement of the Southern Florida Reef Tract Benthic Habitat Map with Habitat Use Patterns of Reef Fish Species* [Dataset]. NOAA National Centers for Environmental Information. <https://www.ncei.noaa.gov/archive/accession/0224176>
- Blondeau, J. B., Grove, J., & Harper, R. (2025). *Mission: Iconic Reefs RVC Analysis*. NOAA Coral Reef Conservation Program.
- Goergen, E. A., Schopmeyer, S., Moulding, A. L., & Moura, A. (2020). *Coral Reef Restoration Monitoring Guide: Methods to evaluate restoration success from local to ecosystem scales* (Technical Report NOS NCCOS 279). National Oceanic; Atmospheric Administration. <https://doi.org/10.25923/xndz-h538>
- Grove, L. J. W., Viehman, T. S., Blondeau, J., Cain, E., Edwards, K. F., Krampitz, N., Langwiser, C., Mateski, J., Sturm, A. B., Swanson, D., & Towle, E. K. (in prep). *National Coral Reef Monitoring Program, Biological Monitoring Summary – Florida: 2024* [Technical Memorandum NOS CRCP, in preparation]. NOAA.
- Groves, S., Mateski, J., Krampitz, N., Sturm, A. M., & Viehman, T. S. (2025). *NCRMP benthics statistical package in R* (website). R package version 1.0.0. https://github.com/MSE-NCCOS-NOAA/NCRMP_benthics
- NOAA Coral Reef Conservation Program (CRCP). (2024a). *National Coral Reef Monitoring Program (NCRMP) Benthic Community Assessment Survey Field Protocols for U.S. Atlantic: Florida, Flower Garden Banks, Puerto Rico, and U.S. Virgin Islands* (p. 29). NOAA Coral Reef Conservation Program. <https://doi.org/10.25923/e3gf-vk84>
- NOAA Coral Reef Conservation Program (CRCP). (2024b). *National Coral Reef Monitoring Program (NCRMP) Coral Demographics Survey Field Protocols for U.S. Atlantic: Florida, Flower Garden Banks, Puerto Rico, and U.S. Virgin Islands* (p. 28). NOAA Coral Reef Conservation Program. <https://doi.org/10.25923/hgpr-jm80>
- NOAA Coral Reef Conservation Program (NOAA CRCP). (2022a). *National Coral Reef Monitoring Program (NCRMP) Benthic Community Assessment Survey Field Protocols for U.S. Atlantic: Florida, Flower Garden Banks, Puerto Rico, and U.S. Virgin Islands–2022*. NOAA Coral Reef Conservation Program. <https://doi.org/10.25923/0708-8333>
- NOAA Coral Reef Conservation Program [NOAA]. (2022b). *National Coral Reef Monitoring Program (NCRMP) Coral Demographics Survey Field Protocols for U.S. Atlantic: Florida, Flower Garden Banks, Puerto Rico, and U.S. Virgin Islands–2022*. NOAA Coral Reef Conservation Program. <https://doi.org/10.25923/9a1r-m911>

- NOAA National Marine Fisheries Service. (2014). Endangered and Threatened Wildlife and Plants: Final Listing Determinations on Proposal to List 66 Reef-Building Coral Species and to Reclassify Elkhorn and Staghorn Corals. *Federal Register*, 79(175), 53851–54123.
- Smith, S. G., Swanson, D. W., Chiappone, M., Miller, S. L., & Ault, J. S. (2011). Probability sampling of stony coral populations in the Florida Keys. *Environmental Monitoring and Assessment*, 183(1–4), 121–138. <https://doi.org/10.1007/s10661-011-1912-2>
- Viehman, T. S., Grove, J. G., Blondeau, J., Cain, E., Edwards, K. F., Groves, S. H., Krampitz, N., Langwiser, C., Siceloff, L., Swanson, D., Towle, E. K., & Williams, B. (2023). *National Coral Reef Monitoring Program, Biological Monitoring Summary – Florida: 2022* [Technical Memorandum NOS CRCP 48]. NOAA. <https://doi.org/10.25923/e6xg-1989>

Data Sources:

NOAA National Centers for Coastal Ocean Science; NOAA Southeast Fisheries Science Center (2024). National Coral Reef Monitoring Program: Assessment of coral reef benthic communities in Florida [2022, 2024]. NOAA National Centers for Environmental Information. Dataset. doi: [10.7289/v5xw4h4z](https://doi.org/10.7289/v5xw4h4z)

Florida Fish and Wildlife Conservation Commission's Disturbance Response Monitoring (DRM)