



The low-latitude boundary layer (LLBL)

Takuma (T.K.M.) Nakamura

Space Research Institute/
Austrian Academy of Sciences



A note of my talk:

My speech is awkward by stuttering disability ☹

But, enjoy my talk!

- the content is not awkward ☺
- science has no borders ☺

Outline

1. Observational features of LLBL

- Basic structures
- IMF dependences
- Local time dependences
- Coupling with ionosphere

2. Candidate formation mechanisms

- Magnetic reconnection
- Kelvin-Helmholtz (KH) instability
- Kinetic Alfvén waves (KAWs)
- Cross-process couplings

Outline

1. Observational features of LLBL

- Basic structures
- IMF dependences
- Local time dependences
- Coupling with ionosphere

2. Candidate formation mechanisms

- Magnetic reconnection
- Kelvin-Helmholtz (KH) instability
- Kinetic Alfvén waves (KAWs)
- Cross-process couplings

Basic features

"The Plasma Boundary Layer and Magnetopause Layer of the Earth's Magnetosphere"

PhD thesis by Eastman [1979]




Figure 1.1 A sketch of the earth's magnetosphere showing the plasma boundary layer (PBL) and magnetopause layer (MPL).




Figure 2.2.2 Satellite coverage of the sunward magnetopause and plasma boundary layer (PBL) for IMP 6, HEOS 2 and the ISEE 1 and ISEE 2 satellite pair. ISEE coverage is given only for October–December, 1977.

- In the 1970's, in-situ observations initially explored various boundary layers in the Earth's magnetosphere.






Figure 2.3.6.1 22 May 1972 plasma data.

Basic features

- In the last 50 years, the development of observational techniques/resources enabled to know detailed structure of the LLBL.
- Magnetopause has a multi-layered structure.
 - PDL(1), MSBL(2), LLBL(4-6)
- LLBL has a distinct (at least) two-layered structure.
 - Inner LLBL: closed
 - Outer LLBL: newly closed



- LLBL tends to be thicker during northward IMF.
- Multi-spacecraft observations revealed the steady expansion of the LLBL during northward IMF.

IMF dependences

Mitchell+ [1987]

Southward IMF (S)



Dawn 13 passes Average = 13.8 minutes
Median = 5 minutes

Dusk 8 passes Average = 15 minutes
Median = 7 minutes


Northward IMF (N)

Dawn 13 passes Average = 90 minutes
Median = 60 minutes

Dusk 6 passes Average = 46 minutes
Median = 38 minutes



Øieroset+ [2008]




IMF dependences

- When IMF is southward, LLBL tends to be highly-structured, and frequently shows transients such as FTEs.
→ by low-lat. reconnection.
- When IMF is northward, LLBL tends to be weakly-structured.

LLBL is thicker and steady during the northward IMF.

Vaisberg+ [2001]



Local time dependences

- The LLBL thickness tends to increase with increasing downtail distance.
 - The dense & stagnant LLBL often appears in the flank-to-tail region especially on the dawnside.
- More solar wind entry occurs in a more tailward region on the dawnside?





Mitchell+ [1987]

Southward IMF (S)

Dawn	13 passes	Average = 13.8 minutes
		Median = 5 minutes
Dusk	8 passes	Average = 15 minutes
		Median = 7 minutes

Northward IMF (N)


Dawn	13 passes	Average = 90 minutes
		Median = 60 minutes
Dusk	6 passes	Average = 46 minutes
		Median = 38 minutes




Hasegawa+ [2004a]

Local time dependences

- $T_{e\perp} < T_{e\parallel}$: whole LLBL
- $T_{i\perp} > T_{i\parallel}$: dayside
- $T_{i\perp} < T_{i\parallel}$: flank-to-tail
- $T_{i\parallel}$ is increasing with increasing distance from the subsolar point.
- A clear dawn-dusk asymmetry is seen in the ion energy spectrum in the LLBL.
 - Duskside: two distinct components (M'sphere & M'sheath components)
 - Dawnside: one-component (M'sheath-like low energy component is heated)





Nishino+ [2007]



Coupling with ionosphere



- Velocity shear across the LLBL can drive the field-aligned currents (FACs).
- Theoretical model of the FACs are well consistent with the observed currents.

Sonnerup [1980]



Johnson & Wing [2015]

Wing & Johnson [2015]



Outline

1. Observational features of LLBL

- Basic structures
- IMF dependences
- Local time dependences
- Coupling with ionosphere




2. Candidate formation mechanisms

- Magnetic reconnection
- Kelvin-Helmholtz (KH) instability
- Kinetic Alfvén waves (KAWs)
- Cross-process couplings

Reconnection

Southward IMF


- Reconnection is expected to occur at low-latitude, leading to transient formation of LLBL.
- LLBL can also be peeled away by low-latitude reconnection.
→ may explain why LLBL is thinner and highly-structured during the southward IMF.




Northward IMF

- Reconnection is expected to occur behind the cusps.
- Double high-latitude reconnection lead to the formation of the dayside LLBL.
- Bi-directional heated electrons confirmed the occurrence of double high-lat. Reconnection.


Reconnection



Song & Russell [1992]





Li+ [2005]



Northward IMF




- Considering 3D space, high-latitude reconnection occurs sequentially, and forms the closed LLBL more or less away from the Y=0 plane.
- High-latitude reconnection would form the open/closed LLBL in the dayside-to flank region.

Reconnection





Kelvin-Helmholtz instability

- The velocity shear across the magnetopause can drive the KHI at low-latitudes.
- The KH waves/vortices tends to be observed much more frequently during northward IMF.





Kelvin-Helmholtz instability

- Theories/simulations predicted that the non-linear vortex flow induces 2ndary KH/RT instabilities and 2ndary reconnection.
 - The 2ndary reconnection causes a rapid expansion of the mixing layer, while the 2ndary KH/RT causes a gradual transport deeper into the magnetosphere.
- KHI can form the flank-to-tail LLBL.




Kinetic Alfvén waves

- KAWs can be induced at the magnetopause by coupled with compressional waves in the magnetosheath, and cause diffusive transport.
- KAWs are frequently observed near the flank-to-tail magnetopause, and cause perpendicular ion heating.
→ may explain $T_{i\perp} > T_{i\parallel}$ observed in the dayside LLBL.




Cross-process couplings


- Local processes often drive other processes, and such a local cross-process coupling can enhance the mixing/transport rate.
- e.g., KH-induced reconnection, KH/RT and other waves.



KAWs within KH
Chaston+ [2007]



LHs within KH
Tang+ [2012]




Ion acoustic waves within KH
Wilder+ [2016]

Cross-process couplings

- High-latitude reconnection & low-latitude KHI can be “globally” coupled.
- High-latitude reconnection can form the LLBL and enhance density along the magnetopause, which can excite stronger KHI.

$$\gamma_{\text{KH}}^2 = \frac{\rho_1 \rho_2}{(\rho_1 + \rho_2)^2} [\mathbf{k} \cdot (\mathbf{U}_1 - \mathbf{U}_2)]^2 - \frac{1}{\mu_0(\rho_1 + \rho_2)} [(\mathbf{k} \cdot \mathbf{B}_1)^2 + (\mathbf{k} \cdot \mathbf{B}_2)^2] > 0$$



$$M_A^2 = \frac{A}{B} = \frac{\rho_1 \rho_2}{\rho_1 + \rho_2} \cdot \frac{[\mathbf{k} \cdot (\mathbf{U}_1 - \mathbf{U}_2)]^2}{[(\mathbf{k} \cdot \mathbf{B}_1)^2 + (\mathbf{k} \cdot \mathbf{B}_2)^2]/\mu_0} > 1$$




Hasegawa+ [2009]

Cross-process couplings

- MMS observed strong KHWs between pre-existing LLBL and magnetosheath.
- 3D Fully kinetic simulation of this event showed efficient mixing/transport.
- Simulation data quantitatively agrees with observations.




Nakamura+ [2017a, b]



Cross-process couplings

- Parallel electron heating starts just after the onset of reconnection within KHs.
→ may explain $T_{e\perp} < T_{e\parallel}$ observed in the whole LLBL.



Summary

1. Observational features of LLBL


- Basic structures
→ plasma mixing, multi-layered structures
- IMF dependences
→ Thicker & more weakly-structured during northward IMF
- Local time dependences
→ Thicker & larger $T_{i\parallel}$ in flank-to-tail, one-(dawn) or two-(dusk) components
- Coupling with ionosphere
→ via FACs by velocity shear

2. Candidate formation mechanisms

- Magnetic reconnection
→ low- or high-latitude reconnection, dayside-to-flank
- Kelvin-Helmholtz (KH) instability
→ 2ndary instabilities, flank-to-tail
- Kinetic Alfvén waves (KAWs)
→ widely induced in the dayside region
- Cross-process couplings
→ local coupling with KAWs/LHDWs, global coupling of reconnection + KH (and KAWs?)

More recent works


- Global MHD simulation for pure northward IMF + test particles (ions).
- Test particle trajectories show (i) more particles enter from the dawnside, and (ii) entered particles being more strongly heated on the dawnside.
→ consistent with observed dawn-dusk asymmetry in ion-energy spectrum.



Sorathia+ [2019]


More recent works

- Magnetosheath turbulence can excite the KHI more strongly and enhance the mixing (solar wind entry) rate.
- Simulated turbulent spectrum, whose amplitude is enhanced by the KHI, is well consistent with the observations.




Future perspectives

- Global hybrid simulation of the Earth's magnetosphere is also becoming feasible.
→ would support quantitative understanding of cross-process/region couplings.
- Future missions
 - MMS and beyond: multi-points, conjunction
 - BepiColombo: comparison with Mercury



[Palmroth+, 2018]



[Liljeblad+, 2015]

References-1

- Chaston, C. C., et al. (2007), Mode conversion and anomalous transport in Kelvin-Helmholtz vortices and kinetic Alfvén waves at the Earth's magnetopause, *Phys. Rev. Lett.*, 99, 175004.
- Chaston, C. C., Y. Yao, N. Lin, C. Salem, and G. Ueno (2013), Ion heating by broadband electromagnetic waves in the magnetosheath and across the magnetopause, *J. Geophys. Res. Space Physics*, 118, 5579–5591, doi: 10.1002/jgra.50506.
- Cowee, M. M., D. Winske, and S. P. Gary (2009), Two-dimensional hybrid simulations of superdiffusion at the magnetopause driven by Kelvin-Helmholtz instability, *J. Geophys. Res.*, 114, A10209, doi:10.1029/2009JA014222.
- Eastman, T. E., *The Plasma Boundary Layer and Magnetopause Layer of the Earth's Magnetosphere*, Ph.D. Thesis, University of Alaska, 1979 [Los Alamos Scientific Laboratory Report LA-7842-T, June, 1979]
- Eriksson, S., et al. (2016), Magnetospheric Multiscale observations of magnetic reconnection associated with Kelvin-Helmholtz waves, *Geophys. Res. Lett.*, 43, 5606-5615, doi:10.1002/2016GL068783
- Fuselier, S. A., K. J. Trattner, S. M. Petrinec, and B. Lavraud (2012), Dayside magnetic topology at the Earth's magnetopause for northward IMF, *J. Geophys. Res.*, 117, A08235, doi:10.1029/2012JA017852.
- Fujimoto, M., T. Terasawa, T. Mukai, Y. Saito, T. Yamamoto, and S. Kokubun (1998), Plasma entry from the flanks of the near-Earth magnetotail: Geotail observations, *J. Geophys. Res.*, 103, 4391–4408.
- Gosling, J. T., M. F. Thomsen, S. J. Bame, T. G. Onsager, and C. T. Russell (1990), The electron edge of the low latitude boundary layer during accelerated flow events, *Geophys. Res. Lett.*, 17, 1833–1836.
- Hasegawa, H., M. Fujimoto, Y. Saito, T. Mukai (2004a), Dense and stagnant ions in the low-latitude boundary region under northward interplanetary magnetic field. *Geophys. Res. Lett.* **31**, L06802, doi:10.1029/2003GL19120
- Hasegawa, H., M. Fujimoto, T.-D. Phan, H. Rème, A. Balogh, M.W. Dunlop, C. Hashimoto, R. TanDokoro (2004b), Transport of solar wind into Earth's magnetosphere through rolled-up Kelvin-Helmholtz vortices. *Nature* **430**, 755–758
- Hasegawa, H. (2012). Structure and dynamics of the magnetopause and its boundary layers. *Monographs on Environment, Earth and Planets*, 1(2), 71–119. <https://doi.org/10.5047/meep.2012.00102.0071>
- Johnson, J. R., and S. Wing (2015), The dependence of the strength and thickness of field-aligned currents on solar wind and ionospheric parameters, *J. Geophys. Res. Space Physics*, 120, 3987–4008, doi:10.1002/2014JA020312.
- Kavosi, S., & Raeder, J. (2015). Ubiquity of Kelvin–Helmholtz waves at Earth's magnetopause. *Nature Communications*, 6, 7019. <https://doi.org/10.1038/ncomms8019>
- Øieroset, M., et al. (2008), THEMIS multi-spacecraft observations of magnetosheath plasma penetration deep into the dayside low-latitude magnetosphere for northward and strong By IMF, *Geophys. Res. Lett.*, 35, L17S11, doi:10.1029/2008GL033661.

References-2

- Lavraud, B., et al. (2006), Evidence for newly closed magnetosheath field lines at the dayside magnetopause under northward IMF, *J. Geophys. Res.*, 111, A05211, doi:10.1029/2005JA011266.
- Li, W., J. Raeder, J. Dorelli, M. Oieroset, and T. D. Phan (2005), Plasma sheet formation during long period of northward IMF, *Geophys. Res. Lett.*, 32, L12S08, doi:10.1029/2004GL021524.
- Liljeblad, E., T. Karlsson, J. M. Raines, J. A. Slavin, A. Kullen, T. Sundberg, and T. H. Zurbuchen (2015), MESSENGER observations of the dayside low-latitude boundary layer in Mercury's magnetosphere, *J. Geophys. Res. Space Physics*, 120, 8387–8400, doi:10.1002/2015JA021662.
- Mitchell, D. G., F. Kutchko, D. J. Williams, T. E. Eastman, L. A. Frank, and C. T. Russell (1987), An extended study of the low-latitude boundary layer on the dawn and dusk flanks of the magnetopause, *J. Geophys. Res.*, 92, 7394–7404, doi:10.1029/JA092iA07p07394.
- Nakamura, T. K. M., & Daughton, W. (2014). Turbulent plasma transport across the Earth's low-latitude boundary layer. *Geophysical Research Letters*, 41, 8704–8712. <https://doi.org/10.1002/2014GL061952>
- Nakamura, T. K. M., Hasegawa, H., Daughton, W., Eriksson, S., Li, W. Y., & Nakamura, R. (2017a). Turbulent mass transfer caused by vortex induced reconnection in collisionless magnetospheric plasmas. *Nature Communications*, 8, 1582. <https://doi.org/10.1038/s41467-017-01579-0>
- Nakamura, T. K. M., et al. (2017b), Mass and energy transfer across the Earth's magnetopause caused by vortex-induced reconnection. *Journal of Geophysical Research: Space Physics*, 122, 11,505–11,522. <https://doi.org/10.1002/2017JA024346>
- Nakamura, T. K. M. (2020). The Earth's low-latitude boundary layer. In R. Maggiolo et al. (Eds.), *Magnetospheres in the Solar System*. Hoboken, NJ: John Wiley & Sons.
- Nakamura, T. K. M., Stawarz, J. E., Hasegawa, H., Narita, Y., Franci, L., Wilder, F. D. et al. (2020). Effects of fluctuating magnetic field on the growth of the Kelvin-Helmholtz instability at the Earth's magnetopause. *Journal of Geophysical Research: Space Physics*, 125, e2019JA027515. <https://doi.org/10.1029/2019JA027515>
- Němeček, Z., J. Šafránková, O. Kruparova, L. Přech, K. Jelínek, Š. Dušík, J. Šimůnek, K. Grygorov, and J.-H. Shue (2015), Analysis of temperature versus density plots and their relation to the LLBL formation under southward and northward IMF orientations. *J. Geophys. Res. Space Physics*, 120, 3475–3488, doi:10.1002/2014JA020308.
- Nishino, M. N., M. Fujimoto, T. Terasawa, G. Ueno, K. Maezawa, T. Mukai, and Y. Saito (2007), Geotail observations of temperature anisotropy of the two-component protons in the dusk plasma sheet, *Ann. Geophys.*, 25, 769–777.
- Nykyri, K., and A. Otto (2001), Plasma transport at the magnetospheric boundary due to reconnection in Kelvin-Helmholtz vortices, *Geophys. Res. Lett.*, 28(18), 3565–3568
- Palmroth, M., U., Ganse, Y., Pfau-Kempf et al. (2018), Vlasov methods in space physics and astrophysics, *Living Rev. Comput. Astrophys.* 4, 1, <https://doi.org/10.1007/s41115-018-0003-2>

References-3

- Song, P., and C. T. Russell (1992), Model of the formation of the low latitude boundary layer for strongly northward interplanetary magnetic field, *J. Geophys. Res.*, 97, 1411.
- Sonnerup, B. U. Ö, (1980), Theory of the low-latitude boundary layer, *J. Geophys. Res.*, 85(A5), 2017–2026.
- Sorathia, K., Merkin, V. G., Ukhorskiy, A. Y., Allen, R. C., Nykyri, K., & Wing, S. (2019). Solar wind ion entry into the magnetosphere during northward IMF. *Journal of Geophysical Research: Space Physics*, 124, 5461–5481. <https://doi.org/10.1029/2019JA026728>
- Tang, B., Li, W., Wang, C., Dai, L., Khotyaintsev, Y., Lindqvist, P.-A., et al. (2018). Magnetic depression and electron transport in an ion-scale flux rope associated with Kelvin–Helmholtz waves. *Annales Geophysicae*, 36, 879–889. <https://doi.org/10.5194/angeo-36-879-2018>
- Vaisberg, O. L., V. sN. Smirnov, L. A. Avanov, J. H. Waite Jr., J. L. Burch, D. L. Gallagher, and N. L. Borodkova (2001), Different types of low-latitude boundary layer as observed by Interball Tail probe, *J. Geophys. Res.*, 106, 13,067.
- Wing, S., J., et al. (2014), Review of Solar Wind Entry into and Transport Within the Plasma Sheet, *Space Science Reviews*, 10.1007/s11214-014-0108-9, **184**, 1-4, (33-86).
- Wing, S., and J. R. Johnson (2015), Theory and observations of upward field-aligned currents at the magnetopause boundary layer, *Geophys. Res. Lett.*, 42, 9149–9155, doi:10.1002/2015GL065464.
- Wilder, F. D., et al. (2016), Observations of large-amplitude, parallel, electrostatic waves associated with the Kelvin-Helmholtz instability by the magnetospheric multiscale mission, *Geophys. Res. Lett.*, 43, 8859–8866, doi:10.1002/2016GL070404.
- Yao, Y., C. C. Chaston, K.-H. Glassmeier, and V. Angelopoulos (2011), Electromagnetic waves on ion gyro-radii scales across the magnetopause, *Geophys. Res. Lett.*, 38, L09102, doi:10.1029/2011GL047328.