

Diagnosing Geospace Dynamics with Magnetometer Arrays

Prof. Ian R. Mann

Department of Physics
University of Alberta

ian.mann@ualberta.ca




Institute for Space Science, Exploration and Technology
University of Alberta

"This presentation is being recorded"



(Image Courtesy of NASA)



Gespace Dynamics & Space weather

"This presentation is being recorded"

The Challenge: Characterise Global Geospace Dynamics



Sparse In-situ Satellite Data



Institute for Space Science, Exploration and Technology
University of Alberta

One of the major scientific opportunities for ground-based magnetometer data is in support of satellite missions.
Examples from THEMIS and Van Allen Probes.


"This presentation is being recorded"



Future Magnetospheric Constellation Mission (MagCon)?



Cubesats?



MagCon Mission?




Institute for Space Science, Exploration and Technology
University of Alberta

"This presentation is being recorded"



Low-Earth Orbit Constellation: AMPERE

28 Aug 2014 13:46:00 - 13:56:00 UT



- Using ADCS magnetometer measurements from Iridium constellation (resolution ~ 30 nT).
- E.g., Anderson et al. JGR (2000).




Institute for Space Science, Exploration and Technology
University of Alberta



"This presentation is being recorded"

Dungey Cycle and Field-Aligned Current Closure

ULF waves transport energy in the global system, and Alfvén waves are required to establish or change FAC.



Recent work with Swarm:
Pakhotin et al. (2018, 2020),
Mann et al. (2019).

From Coxon et al., JGR, (2014).



Institute for Space Science, Exploration and Technology
University of Alberta



"This presentation is being recorded"

Global Scale View Possible with Ground-Magnetometer Arrays



SuperMAG: <http://supermag.jhuapl.edu/>
(PI: Jesper Gjerloev; Jesper.Gjerloev@jhuapl.edu)



Institute for Space Science, Exploration and Technology
University of Alberta

Global Coordination promoted by Ultra-Large Terrestrial
International Magnetometer Array (ULTIMA)
Current Chair: Peter Chi (pchi@igpp.ucla.edu)

"This presentation is being recorded"



Meso-Scale View Possible with Dense Arrays



Fluxgate
magnetometers
(CGSM and FGM)
Resolution < 0.1 nT
Raw 8 sps;
standard 1 or 2 sps.

Induction Coil
Magnetometers
(ICM).
Resolution in pT.
100 sps.

Canadian Array for Real-time Investigations of
Magnetic Activity (PI: Mann; PM: Milling)

www.carisma.ca



Institute for Space Science, Exploration and Technology
University of Alberta



Diagnosing Geospace Dynamics With Magnetometer Arrays

- Energy input into the geospace system – focus on ULF wave/disturbances here.
- Characterising magnetospheric substorms - especially onset and early expansion phase
- Magnetic Storm dynamics: Radiation Belt and Ring Current Wave-Particle Interactions
- Magnetospheric magneto-seismology.




Institute for Space Science, Exploration and Technology
University of Alberta




"This presentation is being recorded"

Kelvin-Helmholtz Instability at the Magnetopause

- Observed at flanks of magnetised planetary magnetospheres.
 - Earth (e.g., Fairfield and Otto, 2000; Hasegawa et al., Nature, 2004; Taylor et al., 2008)
 - Saturn (e.g, Masters et al., 2009), etc




Institute for Space Science, Exploration and Technology
University of Alberta




"This presentation is being recorded"

Excitation of Waveguide Modes



SLOW SOLAR WIND



FAST SOLAR WIND



Institute for Space Science, Exploration and Technology
University of Alberta



Cavity/waveguide mode excitation by solar wind impulses or by magnetopause shear flow/KHI. (Mann and Wright, 1999)


"This presentation is being recorded"

Transport of Energy via Ultra-Low Frequency (ULF) Waves

Rae et al. 2007 estimate total ULF wave energy into ionosphere via Joule heating may actually be >30% of that in a substorm cycle.



Institute for Space Science, Exploration and Technology
University of Alberta





From Mann (2019); in Geomagnetism, Aeronomy and Space Weather: A Journey from the Earth's Core to the Sun, Mandea, Korte, Yau, and Petrovsky (eds.). CUP, (2019).

"This presentation is being recorded"



Driven Field Line Resonance

25 Nov 2001: CANOPUS magnetometers



Ozeke et al., JGR, 2009



Institute for Space Science, Exploration and Technology
University of Alberta



"This presentation is being recorded"

Flank ULF Interactions?

- Large scale dB/dt is also related to the risk of geomagnetically induced currents (GICs) in the electric power grid.
- Recent statistical analysis of U.K. ground-based magnetometer data show the largest dB/dt occur on the flanks due to convection and not at local midnight due to substorms (Freeman et al., Space Weather, 2019).
- New research (Dimitrakoudis et al., in prep., 2020) reveal some new and unexpected flank magnetic dynamics.




Institute for Space Science, Exploration and Technology
University of Alberta

"This presentation is being recorded"



Flank dB/dt: March 2015





Courtesy of Stavros
Dimitrakoudis.



Institute for Space Science, Exploration and Technology
University of Alberta


"This presentation is being recorded"





"This presentation is being recorded"

MARCH 2015 STORM



Courtesy of
SuperMAG.




Institute for Space Science, Exploration and Technology
University of Alberta

"This presentation is being recorded"



Magnetospheric Substorm



Angelopoulos et al. (2008), *Science*

Substorm growth, expansion and recovery phases
(e.g., McPherron et al. 1973).




Institute for Space Science, Exploration and Technology
University of Alberta



"This presentation is being recorded"


Magnetospheric Substorm

Outside
To
In



Inside
To
Out

What is the physics of
near-Earth Substorm
processes?



Current Disruption Paradigm: $A \rightarrow B \rightarrow C$

"This presentation is being recorded"



Institute for Space Science, Exploration and Technology
University of Alberta



What triggers the fast release of stored magnetic energy in reconnection driven magnetotails?




Courtesy of NASA
THEMIS team.

Forecast GIC in
electrical power
grids?



Institute for Space Science, Exploration and Technology
University of Alberta



"This presentation is being recorded"




Ground-based Magnetic Substorm Diagnostics

- Expansion phase onset.
- Substorm onset magnetosphere-ionosphere coupling
- Development of substorm expansion phase: westward travelling surge, westward and poleward expansion, etc.



Institute for Space Science, Exploration and Technology
University of Alberta




"This presentation is being recorded"



Pi2s: Timing Expansion Phase Onset

- Pi2s are irregular waves in the 40s – 150s period band (e.g., Jacobs et al., 1964).
- Can be used to time, and to some degree locate, expansion phase currents.
- Pi2 polarisation ellipses point to longitude of centre of the substorm current wedge (e.g., Lester et al., 1983).
- Pi2 timing, with ~2 minute uncertainty, can time substorm onset. Especially clear in mid-latitude stations in the sub-auroral zone.



Institute for Space Science, Exploration and Technology
University of Alberta


"This presentation is being recorded"



Pi1, and Pi1-2s: Timing and Location of Expansion Phase Onset

24 - 96s
period
wavelet
band.

Rae,
Mann, et
al., JGR,
2009.






Institute for Space Science, Exploration and Technology
University of Alberta



- Actually, the wave response is spatio-temporally inhomogeneous across the Pi1, Pi1-2, and Pi2 spectra.

"This presentation is being recorded"



Rae, Mann et al. JGR (2009)

Keograms


Ground
Substorm bays

Filtered dB

24-96s DWT



Institute for Space Science, Exploration and Technology
University of Alberta



"This presentation is being recorded"





Relation between substorm onset auroral beads and Pi1-2



Institute for Space Science, Exploration and Technology
University of Alberta

Supplementary
Material from Rae,
Mann, et al., JGR,
2009.



"This presentation is being recorded"





Movie courtesy of
Jonathan Rae;
(Rae, Mann et al.
JGR, 2009).

"This presentation is being recorded"



Institute for Space Science, Exploration and Technology
University of Alberta







2007-03-07 05:51:18

Propagation is Waveband Dependent

- Small substorm on 5th March 2008.
- From Mann et al. (2019); adapted from Figures 1 and 4 from Rae et al. (2017).



Substorm Magnetosphere-Ionosphere Coupling



Institute for Space Science, Exploration and Technology
University of Alberta

Pi1-2 activity also at GEO.




"This presentation is being recorded"

Delays THEMIS to Ground

From Smith et al., (2020b)

- Smith et al., (2020a, 2020b) examine THEMIS satellite to ground ULF wave delays, and relative powers.
- Examine period dependence of patro-temporal characteristics.
- Relative to GILL at 06:01:54 in the 93.5s band.



Wave paths connecting onset magnetosphere and ground are key to understanding.

Clear connection to onset processes in the near-tail.



Institute for Space Science, Exploration and Technology
University of Alberta

"This presentation is being recorded"



ULF Wave Impacts on Radiation Belt Electron Dynamics

- Coherent ULF wave interactions (e.g., Mann et al., 2013; Claudepierre et al., 2013; Hartinger et al., 2020.)
- Data driven ULF wave radial diffusion (e.g., Mann et al., 2016; Ozeke et al., 2020)
- Electromagnetic ion cyclotron (EMIC) waves and related loss to the atmosphere (e.g., Usanova et al. 2015 and references therein).



Institute for Space Science, Exploration and Technology
University of Alberta




"This presentation is being recorded"

Power of the Combination of Ground and Satellite Data



Institute for Space Science, Exploration,
University of Alberta



"This presentation is being recorded"

Ultra-Low Frequency (ULF) Wave-Particle Interactions




ULF Wave-MeV Electron Energy Exchange

- Rate of energy change due to ULF interactions:

$$\frac{dW}{dt} = q\mathbf{E} \bullet \mathbf{V}_d + \frac{M}{\gamma} \frac{\partial b}{\partial t}$$



- Effect from electric field and compressional magnetic field; dominated by electric component (Ozeke et al., 2012).
- Can transport particles along phase space density gradients: inwards (energisation) or outwards (e.g., magnetopause loss; Loto'aniu et al., 2010; Turner et al., 2012).




"This presentation is being recorded"



ULF Wave-Particle Drift Resonance




Images courtesy of Scot Elkington, LASP.
"This presentation is being recorded"



Strong Correlation Between Solar Wind, ULF waves, and Radiation Belt Electron Dynamics.

ULF Waves, Fast Solar Wind Streams and MeV Electrons at GEO


(From Mathie and Mann, GRL, 2000).




Coherent ULF Wave Interactions

From Mann et al.,
Nature Comm.,
2013.

See also
Hartinger et al.
2020.



Institute for Space Science, Exploration and Technology
University of Alberta



Thanks to LANL and CDAWeb for LANL and CRRES Data.

"This presentation is being recorded"



Van Allen Probes


- Mann et al., Nature Comms., 2013;
Claudepierre et al., GRL, 2013.




Institute for Space Science, Exploration and Technology
University of Alberta

"This presentation is being recorded"

(a) Van Allen Probes A/ECT/MagEIS & REPT
Spin-Averaged Differential Electron Flux




(b)



ULF Wave Radial Diffusive Transport

Model (Brizard & Chan, Phys. Plasmas, 2004)



*These two terms can be derived in space empirically.
But electric dominates – allows DLL characterization
from ground.*



Institute for Space Science, Exploration and Technology
University of Alberta



Ozeke et al., 2012, 2013, 2014


"This presentation is being recorded"

Third Radiation Belt

(this also required L* mapping)

Driven by observed ULF waves.


Mann et al., Nature Physics, 2016.



3.4 MeV

2012 September

"This presentation is being recorded"




5.2 MeV

2012 September

ULF Wave Radial Diffusion: Data Driven Transport

March 2013 Storm
(Ozeke et al. 2020)

Enhanced very short
additional DLL based on
Olifer et al. (2019)




"This presentation is being recorded"



ULF Wave Radial Diffusion: Data Driven Transport

March 2015
Storm
(Ozeke et al.
2020)



Additional Loss at
 $L \leq 3.5$



"This presentation is being recorded"



Electromagnetic Ion Cyclotron (EMIC) Waves



Institute for Space Science, Exploration and Technology
University of Alberta


Van Allen Probes and Ground -
Mann et al. GRL (2014)

"This presentation is being recorded"



Impact of EMIC Waves on Radiation Belt Electrons?

From Usanova et al.
(2014). See also recent
modelling from Drozdov
et al., (2020).



Magnetoseismology: The cross phase technique

- Cross phase allows the determination of the field line resonant frequency between two latitudinally spaced ground magnetometers.
- The technique relies on the phase difference across the resonant field line.
- Works even when the signal at the resonance is not the dominant one.
- Can also be used with a magnetic field and field-aligned density model to invert into equatorial plasma mass density
- E.g., Baransky et al., 1985; Waters et al., 1991, 1994, 1995; Mathie et al., 1999a, 1999b; Chi et al., 2000, 2005, 2013; Dent et al., 2006...




Institute for Space Science, Exploration and Technology
University of Alberta



"This presentation is being recorded"

Storm-time Mass Density Dynamics

- Dynamic evolution of plasmasphere proposed as a result of distortion of plasmasphere in time dependent convection plus corotation field (e.g., Grebowsky, 1970).



Institute for Space Science, Exploration and Technology
University of Alberta


"This presentation is being recorded"



ISEE Electron Density Evolution

- Electron density dynamically eroded in plasmasphere and plasmapause dynamics.

(From Carpenter and Anderson, 1992)





Institute for Space Science, Exploration and Technology
University of Alberta

"This presentation is being recorded"

IMAGE Mission Observations

- Plasmaspheric evolution from IMAGE EUV (Goldstein et al., 2004)
- “Difference images show plasmaspheric structure inside plasmasphere (Adrian et al., 2004)

(From Goldstein et al, 2004)




Cross-phase and IMAGE RPI plasma mass density profiles.

- Equatorial plane cross-phase densities have been calculated assuming a dipolar field geometry and a r^3 radial density distribution.
- In-situ RPI electron number densities are shown.
- The heavy ion population may be inferred from the difference between the two profiles. For example the O⁺ ion torus (cf. Fraser et al., 2005).



Institute for Space Science, Exploration and Technology
University of Alberta



Dent et al., JGR, 2006.

"This presentation is being recorded"

Summary and Conclusions

- Ground magnetometer networks can provide powerful global and dense meso-scale monitoring of geospace environment.
- Requires physics interpretation to map into geospace, but provides power augment to in-situ measurements from satellites.
- The field-guided nature of Alfvénic disturbances means ground-based magnetometry is a very powerful tool for discovery on a network basis.
- Comments or questions welcome: imann@ualberta.ca



Institute for Space Science, Exploration and Technology
University of Alberta



"This presentation is being recorded"

Acknowledgements

- IRM thanks many collaborators over many years, and for research materials used in this talk.
- The work is supported by Canadian NSERC, the Canadian Space Agency, Canada Foundation for Innovation, Compute Canada, and DND.



Institute for Space Science, Exploration and Technology
University of Alberta



"This presentation is being recorded"