

Notes of Ch6: Estimation I, "introductory Statistics" by Wonnacott

Mahdi Sadeghzadeh Ghamsary

June 2019

1 Introduction

because population is fixed so the population parameters like mean μ and variance σ^2 are fixed (though generally unknown).

the sample mean \bar{X} and sample variance s^2 are random variables, varying from sample to sample, with certain probability distribution. the random variables calculated from the observations in a sample are called sample statistic.

the point estimate of μ is the an estimation by computing one sample mean, but it is distributed around μ . so we must estimate that μ is bracketed by some interval –known as confidence interval– of the following form:

$$\mu = \bar{X} \pm \text{an error allowance} \quad (1)$$

as we can be specific about distribution of \bar{X} , we can be specific about this error allowance.

How confident do we wish to be that our estimate is right about the μ ? it is common to choose 95%100 confidence. to get this confidence level the smallest range under normal distribution of \bar{X} that we just enclose a 95%100 probability. since the distribution is symmetric around the middle chunk, we can leave 2.5%100 probability in each side. we note that this involves going above(or below) the mean by 1.96 standard deviation of \bar{X} . we there for write:

$$\begin{aligned} p\left(\mu - 1.96 \frac{\sigma}{\sqrt{n}} < \bar{X} < \mu + 1.96 \frac{\sigma}{\sqrt{n}}\right) &= 95\% \\ \Rightarrow p\left(-1.96 < \frac{\bar{X} - \mu}{\frac{\sigma}{\sqrt{n}}} < 1.96\right) &= 95\% \end{aligned}$$