

Practice Questions

Sunday, 4 October 2026 7:13 pm

EXAMPLE 3 Find the local extreme values of the function

$$f(x, y) = xy - x^2 - y^2 - 2x - 2y + 4.$$

Solution The function is defined and differentiable for all x and y , and its domain has no boundary points. The function therefore has extreme values only at the points where f_x and f_y are simultaneously zero. This leads to

$$f_x = y - 2x - 2 = 0, \quad f_y = x - 2y - 2 = 0,$$

or

$$x = y = -2.$$

Therefore, the point $(-2, -2)$ is the only point where f may take on an extreme value. To see if it does so, we calculate

$$f_{xx} = -2, \quad f_{yy} = -2, \quad f_{xy} = 1.$$



The discriminant of f at $(a, b) = (-2, -2)$ is

$$f_{xx}f_{yy} - f_{xy}^2 = (-2)(-2) - (1)^2 = 4 - 1 = 3.$$

The combination

$$f_{xx} < 0 \quad \text{and} \quad f_{xx}f_{yy} - f_{xy}^2 > 0$$

tells us that f has a local maximum at $(-2, -2)$. The value of f at this point is $f(-2, -2) = 8$. ■



EXAMPLE 6 Find the absolute maximum and minimum values of

$$f(x, y) = 2 + 2x + 4y - x^2 - y^2$$

on the triangular region in the first quadrant bounded by the lines $x = 0$, $y = 0$, and $y = 9 - x$.

Solution Since f is differentiable, the only places where f can assume these values are points inside the triangle (Figure 14.50a) where $f_x = f_y = 0$ and points on the boundary.

(a) **Interior points.** For these we have

$$f_x = 2 - 2x = 0, \quad f_y = 4 - 2y = 0,$$

yielding the single point $(x, y) = (1, 2)$. The value of f there is

$$f(1, 2) = 7.$$

(b) **Boundary points.** We take the triangle one side at a time:

i) On the segment OA , $y = 0$. The function

$$f(x, y) = f(x, 0) = 2 + 2x - x^2$$

may now be regarded as a function of x defined on the closed interval $0 \leq x \leq 9$. Its extreme values (we know from Chapter 4) may occur at the endpoints

$$\begin{array}{ll} x = 0 & \text{where } f(0, 0) = 2 \\ x = 9 & \text{where } f(9, 0) = 2 + 18 - 81 = -61 \end{array}$$

and at the interior points where $f'(x, 0) = 2 - 2x = 0$. The only interior point where $f'(x, 0) = 0$ is $x = 1$, where

$$f(x, 0) = f(1, 0) = 3.$$

ii) On the segment OB , $x = 0$ and

$$f(x, y) = f(0, y) = 2 + 4y - y^2.$$

As in part i), we consider $f(0, y)$ as a function of y defined on the closed interval $[0, 9]$. Its extreme values can occur at the endpoints or at interior points where $f'(0, y) = 0$. Since $f'(0, y) = 4 - 2y$, the only interior point where $f'(0, y) = 0$ occurs at $(0, 2)$, with $f(0, 2) = 6$. So the candidates for this segment are

$$f(0, 0) = 2, \quad f(0, 9) = -43, \quad f(0, 2) = 6.$$

iii) We have already accounted for the values of f at the endpoints of AB , so we need only look at the interior points of the line segment AB . With $y = 9 - x$, we have

$$f(x, y) = 2 + 2x + 4(9 - x) - x^2 - (9 - x)^2 = -43 + 16x - 2x^2.$$



Setting $f'(x, 9 - x) = 16 - 4x = 0$ gives

$$x = 4.$$

At this value of x ,

$$y = 9 - 4 = 5 \quad \text{and} \quad f(x, y) = f(4, 5) = -11.$$

Summary We list all the function value candidates: 7, 2, -61, 3, -43, 6, -11. The maximum is 7, which f assumes at $(1, 2)$. The minimum is -61, which f assumes at $(9, 0)$. See Figure 14.50b. ■



Screen clipping taken: 04/10/2026 7:21 pm




FIGURE 14.51 The box in Example 7.

EXAMPLE 7 A delivery company accepts only rectangular boxes the sum of whose length and girth (perimeter of a cross-section) does not exceed 108 in. Find the dimensions of an acceptable box of largest volume.

Solution Let x , y , and z represent the length, width, and height of the rectangular box, respectively. Then the girth is $2y + 2z$. We want to maximize the volume $V = xyz$ of the box (Figure 14.51) satisfying $x + 2y + 2z = 108$ (the largest box accepted by the delivery company). Thus, we can write the volume of the box as a function of two variables:

$$\begin{aligned} V(y, z) &= (108 - 2y - 2z)yz & \begin{array}{l} V = xyz \text{ and} \\ x = 108 - 2y - 2z \end{array} \\ &= 108yz - 2y^2z - 2yz^2. \end{aligned}$$

Setting the first partial derivatives equal to zero,

$$V_y(y, z) = 108z - 4yz - 2z^2 = (108 - 4y - 2z)z = 0$$

$$V_z(y, z) = 108y - 2y^2 - 4yz = (108 - 2y - 4z)y = 0,$$

gives the critical points $(0, 0)$, $(0, 54)$, $(54, 0)$, and $(18, 18)$. The volume is zero at $(0, 0)$, $(0, 54)$, and $(54, 0)$, which are not maximum values. At the point $(18, 18)$, we apply the Second Derivative Test (Theorem 11):

$$V_{yy} = -4z, \quad V_{zz} = -4y, \quad V_{yz} = 108 - 4y - 4z.$$

Then

$$V_{yy}V_{zz} - V_{yz}^2 = 16yz - 16(27 - y - z)^2.$$

Thus,

$$V_{yy}(18, 18) = -4(18) < 0$$

and

$$[V_{yy}V_{zz} - V_{yz}^2]_{(18,18)} = 16(18)(18) - 16(-9)^2 > 0$$

imply that $(18, 18)$ gives a maximum volume. The dimensions of the package are $x = 108 - 2(18) - 2(18) = 36$ in., $y = 18$ in., and $z = 18$ in. The maximum volume is $V = (36)(18)(18) = 11,664$ in³, or 6.75 ft³. ■

33. $f(x, y) = x^2 + y^2$ on the closed triangular plate bounded by the lines $x = 0$, $y = 0$, $y + 2x = 2$ in the first quadrant
34. $T(x, y) = x^2 + xy + y^2 - 6x$ on the rectangular plate $0 \leq x \leq 5$, $-3 \leq y \leq 3$
54. Find three positive numbers whose sum is 3 and whose product is a maximum.
55. Find the maximum value of $s = xy + yz + xz$ where $x + y + z = 6$.
56. Find the minimum distance from the cone $z = \sqrt{x^2 + y^2}$ to the point $(-6, 4, 0)$.

31. A trough with triangular cross-section lies along the x -axis for $0 \leq x \leq 10$. The slanted sides are given by $z = y$ and $z = -y$ for $0 \leq z \leq 1$ and the ends by $x = 0$ and $x = 10$, where x, y, z are in meters. The trough contains a sludge whose density at the point (x, y, z) is $\delta = e^{-3x}$ kg per m^3 .

- (a) Express the total mass of sludge in the trough in terms of triple integrals.
- (b) Find the mass.

In Exercises 2–7, find an equation for the surface.

2. The vertical plane $y = x$ in cylindrical coordinates.
3. The top half of the sphere $x^2 + y^2 + z^2 = 1$ in cylindrical coordinates.
4. The cone $z = \sqrt{x^2 + y^2}$ in cylindrical coordinates.
5. The cone $z = \sqrt{x^2 + y^2}$ in spherical coordinates.
8. $f(x, y, z) = \sin(x^2 + y^2)$, W is the solid cylinder with height 4 and with base of radius 1 centered on the z axis at $z = -1$.

In Exercises 10–11, evaluate the triple integrals in spherical coordinates.

10. $f(\rho, \theta, \phi) = \sin \phi$, over the region $0 \leq \theta \leq 2\pi$, $0 \leq \phi \leq \pi/4$, $1 \leq \rho \leq 2$.
11. $f(x, y, z) = 1/(x^2 + y^2 + z^2)^{1/2}$ over the bottom half of the sphere of radius 5 centered at the origin.
39. Suppose W is the region outside the cylinder $x^2 + y^2 = 1$ and inside the sphere $x^2 + y^2 + z^2 = 2$. Calculate

$$\int_W (x^2 + y^2) dV.$$

32. Between the cone $z = \sqrt{x^2 + y^2}$ and the first quadrant of the xy -plane, with $x^2 + y^2 \leq 7$.

In Problems 68–70, explain what is wrong with the statement.

70. The volume of a cylinder of height and radius 1 is


$$\int_0^{2\pi} \int_0^1 \int_0^1 1 dz dr d\theta.$$

■ In Problems 71–72, give an example of:

71. An integral in spherical coordinates that gives the volume of a hemisphere.

A bead is made by drilling a cylindrical hole of radius 1 mm through a sphere of radius 5 mm. See Figure 16.55.

- Set up a triple integral in cylindrical coordinates representing the volume of the bead.
- Evaluate the integral.



Derive the general solution of $\frac{dy}{dx} = 2x + 3$

Exercise 5:

Derive the general solution of $\frac{d^2y}{dx^2} = -\sin(x)$

Exercise 6:

Derive the general solution of $\frac{d^2y}{dt^2} = a$, where $a = \text{constant}$

Derive the general solution of $\frac{d^3y}{dt^2} = 3x^2$

Exercise 8:

Derive the general solution of $e^{-x} \frac{d^2y}{dx^2} = 3$

Solve the ODE

$$\frac{d^2y}{dx^2} = x^2 + 2$$

given that $y = 1$ when $x = 0$ and $\frac{dy}{dx} = 2$ when $x = 0$.

Solve the PDE

$$\frac{\partial^2 u}{\partial x \partial y} = \sin x \cos y$$

subject to the conditions

$$\frac{\partial u}{\partial x} = 2x \text{ at } y = \frac{\pi}{2}, \quad u = 2 \sin y \text{ at } x = \pi.$$

Questions for PDEs are included in the lectures already!