

Derivatives and Rates of Change

Tangents

If a curve C has equation y = f(x) and we want to find the tangent line to C at the point P(a, f(a)), then we consider a nearby point Q(x, f(x)), where $x \neq a$, and compute the slope of the secant line PQ:

$$m_{PQ} = \frac{f(x) - f(a)}{x - a}$$





1 Definition The **tangent line** to the curve y = f(x) at the point P(a, f(a)) is the line through P with slope

$$m = \lim_{x \to a} \frac{f(x) - f(a)}{x - a}$$

provided that this limit exists.


EXAMPLE 1 Find an equation of the tangent line to the parabola $y = x^2$ at the point P(1, 1).

SOLUTION Here we have a = 1 and $f(x) = x^2$, so the slope is

$$m = \lim_{x \to 1} \frac{f(x) - f(1)}{x - 1} = \lim_{x \to 1} \frac{x^2 - 1}{x - 1}$$
$$= \lim_{x \to 1} \frac{(x - 1)(x + 1)}{x - 1}$$
$$= \lim_{x \to 1} (x + 1) = 1 + 1 = 2$$

Using the point-slope form of the equation of a line, we find that an equation of the tangent line at (1, 1) is

$$y - 1 = 2(x - 1)$$
 or $y = 2x - 1$








FIGURE 2 Zooming in toward the point (1, 1) on the parabola $y = x^2$



There is another expression for the slope of a tangent line that is sometimes easier to use. If h = x - a, then x = a + h and so the slope of the secant line PQ is

$$m_{PQ} = \frac{f(a+h) - f(a)}{h}$$

$$m = \lim_{h \to 0} \frac{f(a+h) - f(a)}{h}$$



EXAMPLE 2 Find an equation of the tangent line to the hyperbola y = 3/x at the point (3, 1).


Screen clipping taken: 29/10/2024 8:00 pm

SOLUTION Let f(x) = 3/x. Then the slope of the tangent at (3, 1) is


$$m = \lim_{h \to 0} \frac{f(3+h) - f(3)}{h} = \lim_{h \to 0} \frac{\frac{3}{3+h} - 1}{h} = \lim_{h \to 0} \frac{\frac{3 - (3+h)}{3+h}}{h}$$
$$= \lim_{h \to 0} \frac{-h}{h(3+h)} = \lim_{h \to 0} -\frac{1}{3+h} = -\frac{1}{3}$$

Therefore an equation of the tangent at the point (3, 1) is

$$y - 1 = -\frac{1}{3}(x - 3)$$



Velocities 4 1



average velocity =
$$\frac{\text{displacement}}{\text{time}} = \frac{f(a+h) - f(a)}{h}$$

$$v(a) = \lim_{h \to 0} \frac{f(a+h) - f(a)}{h}$$

Derivatives

Definition The **derivative of a function** f at a number a, denoted by f'(a), is

$$f'(a) = \lim_{h \to 0} \frac{f(a+h) - f(a)}{h}$$

if this limit exists.

$$f'(a) = \lim_{x \to a} \frac{f(x) - f(a)}{x - a}$$

EXAMPLE 4 Find the derivative of the function $f(x) = x^2 - 8x + 9$ at the number a.

SOLUTION From Definition 4 we have

$$f'(a) = \lim_{h \to 0} \frac{f(a+h) - f(a)}{h}$$

$$= \lim_{h \to 0} \frac{\left[(a+h)^2 - 8(a+h) + 9 \right] - \left[a^2 - 8a + 9 \right]}{h}$$

$$= \lim_{h \to 0} \frac{a^2 + 2ah + h^2 - 8a - 8h + 9 - a^2 + 8a - 9}{h}$$

$$= \lim_{h \to 0} \frac{2ah + h^2 - 8h}{h} = \lim_{h \to 0} (2a + h - 8)$$

$$= 2a - 8$$

The tangent line to y = f(x) at (a, f(a)) is the line through (a, f(a)) whose slope is equal to f'(a), the derivative of f at a.


If we use the point-slope form of the equation of a line, we can write an equation of the tangent line to the curve y = f(x) at the point (a, f(a)):

$$y - f(a) = f'(a)(x - a)$$

EXAMPLE 5 Find an equation of the tangent line to the parabola $y = x^2 - 8x + 9$ at the point (3, -6).

SOLUTION From Example 4 we know that the derivative of $f(x) = x^2 - 8x + 9$ at the number a is f'(a) = 2a - 8. Therefore the slope of the tangent line at (3, -6) is f'(3) = 2(3) - 8 = -2. Thus an equation of the tangent line, shown in Figure 7, is

$$y - (-6) = (-2)(x - 3)$$
 or $y = -2x$



Rates of Change

$$\frac{\Delta y}{\Delta x} = \frac{f(x_2) - f(x_1)}{x_2 - x_1}$$

instantaneous rate of change =
$$\lim_{\Delta x \to 0} \frac{\Delta y}{\Delta x} = \lim_{x_2 \to x_1} \frac{f(x_2) - f(x_1)}{x_2 - x_1}$$

The derivative f'(a) is the instantaneous rate of change of y = f(x) with respect to x when x = a.