

Linear Algebra and Geometry 1


Systems of equations, matrices, vectors, and geometry

Pythagorean Theorem and distance between points


Hania Uscka-Wehlou, Ph.D. (2009, Uppsala University: Mathematics)

University teacher in mathematics (Associate Professor / Senior Lecturer) at Mälardalen University, Sweden







$$a^2 + b^2 = c^2$$



$$a^2 + b^2 = c^2$$




$$a^2 + b^2 = c^2$$



$$a^2 + b^2 = c^2$$


$$\frac{x}{a} = \frac{a}{c} \Rightarrow a^2 = xc$$



$$a^2 + b^2 = c^2$$

$$\frac{x}{a} = \frac{a}{c} \Rightarrow a^2 = xc$$

$$\frac{c-x}{b} = \frac{b}{c} \Rightarrow b^2 = c(c-x)$$




$$a^2 + b^2 = c^2$$

$$\frac{x}{a} = \frac{a}{c} \Rightarrow a^2 = xc$$

$$\frac{c-x}{b} = \frac{b}{c} \Rightarrow b^2 = c(c-x)$$

$$a^2 + b^2 =$$




$$a^2 + b^2 = c^2$$

$$\frac{x}{a} = \frac{a}{c} \Rightarrow a^2 = xc$$

$$\frac{c-x}{b} = \frac{b}{c} \Rightarrow b^2 = c(c-x)$$

$$a^2 + b^2 = xc + c(c-x)$$




$$a^2 + b^2 = c^2$$

$$\frac{x}{a} = \frac{a}{c} \Rightarrow a^2 = xc$$

$$\frac{c-x}{b} = \frac{b}{c} \Rightarrow b^2 = c(c-x)$$

$$a^2 + b^2 = xc + c(c-x) = xc + c^2 - xc$$




$$a^2 + b^2 = c^2$$

$$\frac{x}{a} = \frac{a}{c} \Rightarrow a^2 = xc$$

$$\frac{c-x}{b} = \frac{b}{c} \Rightarrow b^2 = c(c-x)$$


$$a^2 + b^2 = xc + c(c-x) = xc + c^2 - xc = c^2$$

Distance between points = length of a vector





Pythagorean theorem

Distance between points = length of a vector



$$d(P, Q) = \|\overrightarrow{PQ}\| = \|\overrightarrow{OQ} - \overrightarrow{OP}\|$$


Cartesian coordinate system



Distance between points = length of a vector

$$P = (x_1, y_1, z_1), \quad Q = (x_2, y_2, z_2)$$

$$d(P, Q) = | \overrightarrow{PQ} | = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$$



Length / Norm

$$|\vec{v}|$$

$$|\mathbf{v}|$$

$$\|\vec{v}\|$$

$$\|\mathbf{v}\|$$

$$\mathbb{R}^n$$

$$\mathbf{x} = (x_1, x_2, \dots, x_n)$$

$$\mathbf{y} = (y_1, y_2, \dots, y_n)$$

$$d(\mathbf{x}, \mathbf{y}) = \sqrt{(y_1 - x_1)^2 + (y_2 - x_2)^2 + \dots + (y_n - x_n)^2}$$