


Projection onto the subspace


We learned earlier how to find the projection of a vector onto a line, but we can also project a vector onto a subspace. So let's start by generalizing the definition of a projection onto a line, so that we can see how it might apply to a subspace.

If you project a vector \vec{x} onto a line L , then the projection vector that sits on L is $\text{Proj}_L \vec{x}$, and the vector from L to the terminal point of \vec{x} , which is orthogonal to L , is $\vec{x} - \text{Proj}_L \vec{x}$.



But let's say we call $\text{Proj}_L \vec{x}$ the vector \vec{v} , and we call $\vec{x} - \text{Proj}_L \vec{x}$ the vector \vec{w} . Then we could say that $\vec{x} - \vec{v} = \vec{w}$.





Then we can rewrite $\vec{x} - \vec{v} = \vec{w}$ as

$$\vec{x} = \vec{v} + \vec{w}$$

Because \vec{w} is orthogonal to L , in the same way that \vec{v} is a member of L , \vec{w} is a member of the orthogonal complement of L , or L^\perp .

And this doesn't just work for the projection of a vector onto a line. It also works for projecting a vector onto any subspace. And just like with the projection onto a line, if \vec{x} is projected onto a subspace V , and the projection is $\text{Proj}_V \vec{x}$, then the vector \vec{x} is closer to the vector $\text{Proj}_V \vec{x}$ than it is to any other vector in V .

The projection is a linear transformation

If V is a subspace of \mathbb{R}^n , then the projection of \vec{x} onto a subspace V is a linear transformation that can be written as the matrix-vector product

$$\text{Proj}_V \vec{x} = A(A^T A)^{-1} A^T \vec{x}$$



where A is a matrix of column vectors that form the basis for the subspace V .

There are a couple of important things we want to realize about this formula. First, notice that the product $A(A^T A)^{-1}A^T$ will eventually simplify to one matrix. Second, it's important to say that we cannot distribute the inverse across the matrix product inside the parentheses. In other words, we **cannot** rewrite the formula as

$$\text{Proj}_V \vec{x} = A(A)^{-1}(A^T)^{-1}A^T \vec{x}$$

The only time we'd be allowed to do this is if A is a square, invertible matrix. But if A is a square, invertible matrix, then by definition A defines all of \mathbb{R}^n , and the projection of a vector \vec{x} onto V will just be itself, \vec{x} . So we only need the projection formula when A is not square and not invertible. But if A is not square and not invertible, such that it makes sense to use the projection formula, then we can't split up the $(A^T A)^{-1}$.

Let's do an example so that we can see how to use it to find the projection of a vector \vec{x} onto the subspace V .

Example

If \vec{x} is a vector in \mathbb{R}^3 , find an expression for the projection of any \vec{x} onto the subspace V .

$$V = \text{Span}\left(\begin{bmatrix} 1 \\ -2 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 1 \\ -2 \end{bmatrix}\right)$$



Because the vectors that span V are linearly independent, the matrix A of the basis vectors that define V is

$$A = \begin{bmatrix} 1 & 2 \\ -2 & 1 \\ 0 & -2 \end{bmatrix}$$

The transpose A^T is then

$$A^T = \begin{bmatrix} 1 & -2 & 0 \\ 2 & 1 & -2 \end{bmatrix}$$

Our next step is to find $A^T A$.

$$A^T A = \begin{bmatrix} 1 & -2 & 0 \\ 2 & 1 & -2 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ -2 & 1 \\ 0 & -2 \end{bmatrix}$$

$$A^T A = \begin{bmatrix} 1(1) - 2(-2) + 0(0) & 1(2) - 2(1) + 0(-2) \\ 2(1) + 1(-2) - 2(0) & 2(2) + 1(1) - 2(-2) \end{bmatrix}$$

$$A^T A = \begin{bmatrix} 1 + 4 + 0 & 2 - 2 + 0 \\ 2 - 2 - 0 & 4 + 1 + 4 \end{bmatrix}$$

$$A^T A = \begin{bmatrix} 5 & 0 \\ 0 & 9 \end{bmatrix}$$

Then we need to find the inverse of $A^T A$.

$$[A^T A \mid I_2] = \left[\begin{array}{cc|cc} 5 & 0 & 1 & 0 \\ 0 & 9 & 0 & 1 \end{array} \right]$$

$$[A^T A \mid I_2] = \left[\begin{array}{cc|cc} 1 & 0 & \frac{1}{5} & 0 \\ 0 & 9 & 0 & 1 \end{array} \right]$$



$$[A^T A \mid I_2] = \left[\begin{array}{cc|cc} 1 & 0 & \frac{1}{5} & 0 \\ 0 & 1 & 0 & \frac{1}{9} \end{array} \right]$$

So $(A^T A)^{-1}$ is

$$(A^T A)^{-1} = \begin{bmatrix} \frac{1}{5} & 0 \\ 0 & \frac{1}{9} \end{bmatrix}$$

Now the projection of \vec{x} onto the subspace V will be

$$\text{Proj}_V \vec{x} = A(A^T A)^{-1} A^T \vec{x}$$

$$\text{Proj}_V \vec{x} = \begin{bmatrix} 1 & 2 \\ -2 & 1 \\ 0 & -2 \end{bmatrix} \begin{bmatrix} \frac{1}{5} & 0 \\ 0 & \frac{1}{9} \end{bmatrix} \begin{bmatrix} 1 & -2 & 0 \\ 2 & 1 & -2 \end{bmatrix} \vec{x}$$

First, simplify $(A^T A)^{-1} A^T$.

$$\text{Proj}_V \vec{x} = \begin{bmatrix} 1 & 2 \\ -2 & 1 \\ 0 & -2 \end{bmatrix} \begin{bmatrix} \frac{1}{5}(1) + 0(2) & \frac{1}{5}(-2) + 0(1) & \frac{1}{5}(0) + 0(-2) \\ 0(1) + \frac{1}{9}(2) & 0(-2) + \frac{1}{9}(1) & 0(0) + \frac{1}{9}(-2) \end{bmatrix} \vec{x}$$

$$\text{Proj}_V \vec{x} = \begin{bmatrix} 1 & 2 \\ -2 & 1 \\ 0 & -2 \end{bmatrix} \begin{bmatrix} \frac{1}{5} + 0 & -\frac{2}{5} + 0 & 0 + 0 \\ 0 + \frac{2}{9} & 0 + \frac{1}{9} & 0 - \frac{2}{9} \end{bmatrix} \vec{x}$$

$$\text{Proj}_V \vec{x} = \begin{bmatrix} 1 & 2 \\ -2 & 1 \\ 0 & -2 \end{bmatrix} \begin{bmatrix} \frac{1}{5} & -\frac{2}{5} & 0 \\ \frac{2}{9} & \frac{1}{9} & -\frac{2}{9} \end{bmatrix} \vec{x}$$



Next, simplify $A(A^T A)^{-1}A^T$.

$$\text{Proj}_V \vec{x} = \begin{bmatrix} 1\left(\frac{1}{5}\right) + 2\left(\frac{2}{9}\right) & 1\left(-\frac{2}{5}\right) + 2\left(\frac{1}{9}\right) & 1(0) + 2\left(-\frac{2}{9}\right) \\ -2\left(\frac{1}{5}\right) + 1\left(\frac{2}{9}\right) & -2\left(-\frac{2}{5}\right) + 1\left(\frac{1}{9}\right) & -2(0) + 1\left(-\frac{2}{9}\right) \\ 0\left(\frac{1}{5}\right) - 2\left(\frac{2}{9}\right) & 0\left(-\frac{2}{5}\right) - 2\left(\frac{1}{9}\right) & 0(0) - 2\left(-\frac{2}{9}\right) \end{bmatrix} \vec{x}$$

$$\text{Proj}_V \vec{x} = \begin{bmatrix} \frac{1}{5} + \frac{4}{9} & -\frac{2}{5} + \frac{2}{9} & 0 - \frac{4}{9} \\ -\frac{2}{5} + \frac{2}{9} & \frac{4}{5} + \frac{1}{9} & 0 - \frac{2}{9} \\ 0 - \frac{4}{9} & 0 - \frac{2}{9} & 0 + \frac{4}{9} \end{bmatrix} \vec{x}$$

$$\text{Proj}_V \vec{x} = \begin{bmatrix} \frac{9}{45} + \frac{20}{45} & -\frac{18}{45} + \frac{10}{45} & -\frac{4}{9} \\ -\frac{18}{45} + \frac{10}{45} & \frac{36}{45} + \frac{5}{45} & -\frac{2}{9} \\ -\frac{4}{9} & -\frac{2}{9} & \frac{4}{9} \end{bmatrix} \vec{x}$$

$$\text{Proj}_V \vec{x} = \begin{bmatrix} \frac{29}{45} & -\frac{8}{45} & -\frac{4}{9} \\ -\frac{8}{45} & \frac{41}{45} & -\frac{2}{9} \\ -\frac{4}{9} & -\frac{2}{9} & \frac{4}{9} \end{bmatrix} \vec{x}$$

To simplify the matrix, let's factor out a $1/9$.

$$\text{Proj}_V \vec{x} = \frac{1}{9} \begin{bmatrix} \frac{29}{5} & -\frac{8}{5} & -4 \\ -\frac{8}{5} & \frac{41}{5} & -2 \\ -4 & -2 & 4 \end{bmatrix} \vec{x}$$



This result tells us that we can find the projection of any \vec{x} onto the subspace V by multiplying the transformation matrix

$$\frac{1}{9} \begin{bmatrix} \frac{29}{5} & -\frac{8}{5} & -4 \\ -\frac{8}{5} & \frac{41}{5} & -2 \\ -4 & -2 & 4 \end{bmatrix}$$

by the vector \vec{x} .

