

Linear Algebra and Geometry 1

Systems of equations, matrices, vectors, and geometry

Linear transformations

Hania Uscka-Wehlou, Ph.D. (2009, Uppsala University: Mathematics)

University teacher in mathematics (Associate Professor / Senior Lecturer) at Mälardalen University, Sweden



An introduction to linear operations

Linear operations preserve linear combinations

Differentiation is a linear operation

$$(f + g)' = f' + g'$$

$$(\alpha f)' = \alpha f'$$

$$(\alpha f + \beta g)' = \alpha f' + \beta g'$$

Integration is linear

$$\int_a^b (f(x) + g(x))dx = \int_a^b f(x)dx + \int_a^b g(x)dx$$

$$\int_a^b \alpha f(x)dx = \alpha \int_a^b f(x)dx$$

$$\int_a^b (\alpha f(x) + \beta g(x))dx = \alpha \int_a^b f(x)dx + \beta \int_a^b g(x)dx$$


Sine is not a linear function

$$\sin(x + y) \neq \sin x + \sin y$$

$$\sin(x + y) = \sin x \cos y + \sin y \cos x$$

Sine is not a linear function

$$\sin(x + y) \neq \sin x + \sin y$$



$$\sin(x + y) = \sin x \cos y + \sin y \cos x$$


Logarithm is not a linear function

$$\ln(x + y) \neq \ln x + \ln y$$

$$\ln(xy) = \ln x + \ln y$$

Logarithm is not a linear function

$$\ln(x + y) \neq \ln x + \ln y$$



$$\ln(xy) = \ln x + \ln y$$


Exponential function is not linear

$$a^{x+y} \neq a^x + a^y$$

$$a^{x+y} = a^x \cdot a^y$$

Exponential function is not linear

$$a^{x+y} \neq a^x + a^y$$



$$a^{x+y} = a^x \cdot a^y$$


Power functions are not linear

$$(x + y)^2 \neq x^2 + y^2$$

$$(x + y)^2 = x^2 + 2xy + y^2$$

Power functions are not linear

$$(x + y)^2 \neq x^2 + y^2$$




$$(x + y)^2 = x^2 + 2xy + y^2$$

Square root is not a linear function

$$\sqrt{x+y} \neq \sqrt{x} + \sqrt{y}$$

Square root is not a linear function

$$\sqrt{x+y} \neq \sqrt{x} + \sqrt{y}$$



Functions, generally

Domain, codomain, range

$$f : X \rightarrow Y$$

$D_f \subset X$ all the possible arguments

Y

$V_f \subset Y$ $V_f = \{ y; y = f(x) \text{ for some } x \in D_f \}$

Domain, codomain, range

$$f : X \rightarrow Y$$

$D_f \subset X$ all the possible arguments

Y

$V_f \subset Y$ $V_f = \{ y; y = f(x) \text{ for some } x \in D_f \}$

$$f : \mathbb{R} \rightarrow \mathbb{R}, \quad f(x) = \frac{1}{x}$$

Domain, codomain, range

$$f : X \rightarrow Y$$

$D_f \subset X$ all the possible arguments

Y

$V_f \subset Y$ $V_f = \{ y; y = f(x) \text{ for some } x \in D_f \}$

$$f : \mathbb{R} \rightarrow \mathbb{R}, \quad f(x) = \frac{1}{x}$$


$$D_f = \mathbb{R} \setminus \{0\} = V_f$$

Functions in Calculus 1

Continuous functions

$$\lim_{x \rightarrow a} f(x) = L$$

$$\forall \varepsilon > 0 \quad \exists \delta > 0 \quad \forall x \in D_f \quad 0 < |x - a| < \delta \Rightarrow |f(x) - L| < \varepsilon$$




The values $f(x)$ can get arbitrarily close (ε -close) to L if only the arguments x are close enough (δ -close) the point a

$$x \rightarrow a \Rightarrow |f(x) - L| \rightarrow 0$$

$$\lim_{x \rightarrow a} f(x) = f(a)$$

$$\forall \varepsilon > 0 \quad \exists \delta > 0 \quad \forall x \in D_f \quad |x - a| < \delta \Rightarrow |f(x) - f(a)| < \varepsilon$$



The values $f(x)$ can get arbitrarily close (ε -close) to $f(a)$ if only the arguments x are close enough (δ -close) the point a

$$x \rightarrow a \Rightarrow |f(x) - f(a)| \rightarrow 0$$

Functions in Linear Algebra

Linear transformations

$$T: \mathbb{R}^n \rightarrow \mathbb{R}^m$$

$$T : \mathbb{R}^n \rightarrow \mathbb{R}^m$$

$$T(\mathbf{x} + \mathbf{y}) = T(\mathbf{x}) + T(\mathbf{y})$$

$$T(\alpha \mathbf{x}) = \alpha T(\mathbf{x})$$

$$T : \mathbb{R}^n \rightarrow \mathbb{R}^m$$

$$T(\mathbf{x} + \mathbf{y}) = T(\mathbf{x}) + T(\mathbf{y})$$

$$T(\alpha \mathbf{x}) = \alpha T(\mathbf{x})$$

$$T(\alpha \mathbf{x} + \beta \mathbf{y}) = \alpha T(\mathbf{x}) + \beta T(\mathbf{y})$$

The image of a linear combination of two vectors
is the same linear combination of the images of the vectors.


$$T : \mathbb{R} \rightarrow \mathbb{R}$$

Linear in Calculus (its graph is a straight line)

Not linear in Linear Algebra

$$f(x_1 + x_2) = x_1 + x_2 + 2$$

$$f(x_1) + f(x_2) = x_1 + 2 + x_2 + 2 = x_1 + x_2 + 4$$



$$T : \mathbb{R} \rightarrow \mathbb{R}$$


Linear in Calculus (its graph is a straight line)

Linear in Linear Algebra

$$f(x_1 + x_2) = x_1 + x_2$$

$$f(x_1) + f(x_2) = x_1 + x_2$$

$$f(\alpha x) = \alpha x = \alpha f(x)$$



$$T : \mathbb{R} \rightarrow \mathbb{R}$$


Linear in Calculus (its graph is a straight line)

Linear in Linear Algebra

$$f(x_1 + x_2) = 3(x_1 + x_2) = 3x_1 + 3x_2$$

$$f(x_1) + f(x_2) = 3x_1 + 3x_2$$

$$f(\alpha x) = 3\alpha x = \alpha 3x = \alpha f(x)$$




$T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$

$T(\alpha \mathbf{x} + \beta \mathbf{y}) = \alpha T(\mathbf{x}) + \beta T(\mathbf{y})$

$T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$

$T(\mathbf{e}_1) = \begin{bmatrix} 2 \\ 4 \end{bmatrix} = \mathbf{f}_1, \quad T(\mathbf{e}_2) = \begin{bmatrix} 3 \\ 1 \end{bmatrix} = \mathbf{f}_2$


$T(\alpha \mathbf{x} + \beta \mathbf{y}) = \alpha T(\mathbf{x}) + \beta T(\mathbf{y})$



$T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$

$T(\mathbf{e}_1) = \begin{bmatrix} 2 \\ 4 \end{bmatrix} = \mathbf{f}_1, \quad T(\mathbf{e}_2) = \begin{bmatrix} 3 \\ 1 \end{bmatrix} = \mathbf{f}_2$

$T(\alpha \mathbf{x} + \beta \mathbf{y}) = \alpha T(\mathbf{x}) + \beta T(\mathbf{y})$




$T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$

$T(\mathbf{e}_1) = \begin{bmatrix} 2 \\ 4 \end{bmatrix} = \mathbf{f}_1, \quad T(\mathbf{e}_2) = \begin{bmatrix} 3 \\ 1 \end{bmatrix} = \mathbf{f}_2$

$T \begin{bmatrix} x \\ y \end{bmatrix} = T(x\mathbf{e}_1 + y\mathbf{e}_2) =$

$T(\alpha \mathbf{x} + \beta \mathbf{y}) = \alpha T(\mathbf{x}) + \beta T(\mathbf{y})$




$$T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

$$T(\mathbf{e}_1) = \begin{bmatrix} 2 \\ 4 \end{bmatrix} = \mathbf{f}_1, \quad T(\mathbf{e}_2) = \begin{bmatrix} 3 \\ 1 \end{bmatrix} = \mathbf{f}_2$$

$$T \begin{bmatrix} x \\ y \end{bmatrix} = T(x\mathbf{e}_1 + y\mathbf{e}_2) =$$

$$= xT(\mathbf{e}_1) + yT(\mathbf{e}_2) = x \begin{bmatrix} 2 \\ 4 \end{bmatrix} + y \begin{bmatrix} 3 \\ 1 \end{bmatrix}$$




$$T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

$$T(\mathbf{e}_1) = \begin{bmatrix} 2 \\ 4 \end{bmatrix} = \mathbf{f}_1, \quad T(\mathbf{e}_2) = \begin{bmatrix} 3 \\ 1 \end{bmatrix} = \mathbf{f}_2$$

$$T \begin{bmatrix} x \\ y \end{bmatrix} = T(x\mathbf{e}_1 + y\mathbf{e}_2) =$$

$$= xT(\mathbf{e}_1) + yT(\mathbf{e}_2) = x \begin{bmatrix} 2 \\ 4 \end{bmatrix} + y \begin{bmatrix} 3 \\ 1 \end{bmatrix}$$




$$T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

$$T(\mathbf{e}_1) = \begin{bmatrix} 2 \\ 4 \end{bmatrix} = \mathbf{f}_1, \quad T(\mathbf{e}_2) = \begin{bmatrix} 3 \\ 1 \end{bmatrix} = \mathbf{f}_2$$

$$T \begin{bmatrix} x \\ y \end{bmatrix} = T(x\mathbf{e}_1 + y\mathbf{e}_2) =$$

$$= xT(\mathbf{e}_1) + yT(\mathbf{e}_2) = x \begin{bmatrix} 2 \\ 4 \end{bmatrix} + y \begin{bmatrix} 3 \\ 1 \end{bmatrix}$$



$T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$


$T(\alpha \mathbf{x} + \beta \mathbf{y}) = \alpha T(\mathbf{x}) + \beta T(\mathbf{y})$

$$T(\mathbf{e}_1) = \begin{bmatrix} 2 \\ 4 \end{bmatrix} = \mathbf{f}_1, \quad T(\mathbf{e}_2) = \begin{bmatrix} 3 \\ 1 \end{bmatrix} = \mathbf{f}_2$$

$$T \begin{bmatrix} x \\ y \end{bmatrix} = T(x\mathbf{e}_1 + y\mathbf{e}_2) =$$

$$= xT(\mathbf{e}_1) + yT(\mathbf{e}_2) = x \begin{bmatrix} 2 \\ 4 \end{bmatrix} + y \begin{bmatrix} 3 \\ 1 \end{bmatrix}$$

The grid lines transform into new grid lines:
parallel and equally spaced.




$$T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

$$T(\mathbf{e}_1) = \begin{bmatrix} 2 \\ 4 \end{bmatrix} = \mathbf{f}_1, \quad T(\mathbf{e}_2) = \begin{bmatrix} 3 \\ 1 \end{bmatrix} = \mathbf{f}_2$$

$$T \begin{bmatrix} x \\ y \end{bmatrix} = T(x\mathbf{e}_1 + y\mathbf{e}_2) =$$

$$= xT(\mathbf{e}_1) + yT(\mathbf{e}_2) = x \begin{bmatrix} 2 \\ 4 \end{bmatrix} + y \begin{bmatrix} 3 \\ 1 \end{bmatrix}$$

The grid lines transform into new grid lines:
parallel and equally spaced.




$$T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

$$T(\mathbf{e}_1) = \begin{bmatrix} 2 \\ 4 \end{bmatrix} = \mathbf{f}_1, \quad T(\mathbf{e}_2) = \begin{bmatrix} 3 \\ 1 \end{bmatrix} = \mathbf{f}_2$$

$$T \begin{bmatrix} x \\ y \end{bmatrix} = T(x\mathbf{e}_1 + y\mathbf{e}_2) =$$

$$= xT(\mathbf{e}_1) + yT(\mathbf{e}_2) = x \begin{bmatrix} 2 \\ 4 \end{bmatrix} + y \begin{bmatrix} 3 \\ 1 \end{bmatrix}$$

The grid lines transform into new grid lines:
parallel and equally spaced.




$$T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

$$T(\mathbf{e}_1) = \begin{bmatrix} 2 \\ 4 \end{bmatrix} = \mathbf{f}_1, \quad T(\mathbf{e}_2) = \begin{bmatrix} 3 \\ 1 \end{bmatrix} = \mathbf{f}_2$$

$$T \begin{bmatrix} x \\ y \end{bmatrix} = T(x\mathbf{e}_1 + y\mathbf{e}_2) =$$

$$= xT(\mathbf{e}_1) + yT(\mathbf{e}_2) = x \begin{bmatrix} 2 \\ 4 \end{bmatrix} + y \begin{bmatrix} 3 \\ 1 \end{bmatrix}$$

The grid lines transform into new grid lines:
parallel and equally spaced.



$$T : \mathbb{R}^n \rightarrow \mathbb{R}^m$$

$$T(\alpha \mathbf{x} + \beta \mathbf{y}) = \alpha T(\mathbf{x}) + \beta T(\mathbf{y})$$

$$T : \mathbb{R}^n \rightarrow \mathbb{R}^m$$

$$T(\alpha \mathbf{x} + \beta \mathbf{y}) = \alpha T(\mathbf{x}) + \beta T(\mathbf{y})$$

$$\mathbf{x} = (x_1, x_2, \dots, x_n) = x_1 \mathbf{e}_1 + x_2 \mathbf{e}_2 + \dots + x_n \mathbf{e}_n$$

$$T : \mathbb{R}^n \rightarrow \mathbb{R}^m$$

$$T(\alpha \mathbf{x} + \beta \mathbf{y}) = \alpha T(\mathbf{x}) + \beta T(\mathbf{y})$$

$$\mathbf{x} = (x_1, x_2, \dots, x_n) = x_1 \mathbf{e}_1 + x_2 \mathbf{e}_2 + \dots + x_n \mathbf{e}_n$$

$$T(\mathbf{e}_1) = \begin{bmatrix} a_{11} \\ a_{21} \\ \vdots \\ a_{m1} \end{bmatrix} = \mathbf{f}_1, \quad T(\mathbf{e}_2) = \begin{bmatrix} a_{12} \\ a_{22} \\ \vdots \\ a_{m2} \end{bmatrix} = \mathbf{f}_2, \quad \dots, \quad T(\mathbf{e}_n) = \begin{bmatrix} a_{1n} \\ a_{2n} \\ \vdots \\ a_{mn} \end{bmatrix} = \mathbf{f}_n$$

$$T : \mathbb{R}^n \rightarrow \mathbb{R}^m$$

$$T(\alpha \mathbf{x} + \beta \mathbf{y}) = \alpha T(\mathbf{x}) + \beta T(\mathbf{y})$$

$$\mathbf{x} = (x_1, x_2, \dots, x_n) = x_1 \mathbf{e}_1 + x_2 \mathbf{e}_2 + \dots + x_n \mathbf{e}_n$$

$$T(\mathbf{e}_1) = \begin{bmatrix} a_{11} \\ a_{21} \\ \vdots \\ a_{m1} \end{bmatrix} = \mathbf{f}_1, \quad T(\mathbf{e}_2) = \begin{bmatrix} a_{12} \\ a_{22} \\ \vdots \\ a_{m2} \end{bmatrix} = \mathbf{f}_2, \quad \dots, \quad T(\mathbf{e}_n) = \begin{bmatrix} a_{1n} \\ a_{2n} \\ \vdots \\ a_{mn} \end{bmatrix} = \mathbf{f}_n$$

$$T \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = T(x_1 \mathbf{e}_1 + x_2 \mathbf{e}_2 + \dots + x_n \mathbf{e}_n) = x_1 T(\mathbf{e}_1) + x_2 T(\mathbf{e}_2) + \dots + x_n T(\mathbf{e}_n) =$$

induction



$$T : \mathbb{R}^n \rightarrow \mathbb{R}^m$$

$$T(\alpha \mathbf{x} + \beta \mathbf{y}) = \alpha T(\mathbf{x}) + \beta T(\mathbf{y})$$

$$\mathbf{x} = (x_1, x_2, \dots, x_n) = x_1 \mathbf{e}_1 + x_2 \mathbf{e}_2 + \dots + x_n \mathbf{e}_n$$

$$T(\mathbf{e}_1) = \begin{bmatrix} a_{11} \\ a_{21} \\ \vdots \\ a_{m1} \end{bmatrix} = \mathbf{f}_1, \quad T(\mathbf{e}_2) = \begin{bmatrix} a_{12} \\ a_{22} \\ \vdots \\ a_{m2} \end{bmatrix} = \mathbf{f}_2, \quad \dots, \quad T(\mathbf{e}_n) = \begin{bmatrix} a_{1n} \\ a_{2n} \\ \vdots \\ a_{mn} \end{bmatrix} = \mathbf{f}_n$$

$$T \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = T(x_1 \mathbf{e}_1 + x_2 \mathbf{e}_2 + \dots + x_n \mathbf{e}_n) = x_1 T(\mathbf{e}_1) + x_2 T(\mathbf{e}_2) + \dots + x_n T(\mathbf{e}_n) =$$

$$T : \mathbb{R}^n \rightarrow \mathbb{R}^m$$

$$T(\alpha \mathbf{x} + \beta \mathbf{y}) = \alpha T(\mathbf{x}) + \beta T(\mathbf{y})$$

$$\mathbf{x} = (x_1, x_2, \dots, x_n) = x_1 \mathbf{e}_1 + x_2 \mathbf{e}_2 + \dots + x_n \mathbf{e}_n$$

$$T(\mathbf{e}_1) = \begin{bmatrix} a_{11} \\ a_{21} \\ \vdots \\ a_{m1} \end{bmatrix} = \mathbf{f}_1, \quad T(\mathbf{e}_2) = \begin{bmatrix} a_{12} \\ a_{22} \\ \vdots \\ a_{m2} \end{bmatrix} = \mathbf{f}_2, \quad \dots, \quad T(\mathbf{e}_n) = \begin{bmatrix} a_{1n} \\ a_{2n} \\ \vdots \\ a_{mn} \end{bmatrix} = \mathbf{f}_n$$

$$T \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = T(x_1 \mathbf{e}_1 + x_2 \mathbf{e}_2 + \dots + x_n \mathbf{e}_n) = x_1 T(\mathbf{e}_1) + x_2 T(\mathbf{e}_2) + \dots + x_n T(\mathbf{e}_n) =$$

$$= x_1 \begin{bmatrix} a_{11} \\ a_{21} \\ \vdots \\ a_{m1} \end{bmatrix} + x_2 \begin{bmatrix} a_{12} \\ a_{22} \\ \vdots \\ a_{m2} \end{bmatrix} + \dots + x_n \begin{bmatrix} a_{1n} \\ a_{2n} \\ \vdots \\ a_{mn} \end{bmatrix}$$