

# A City in Crisis? Analyzing Attended Person in Crisis Calls in Toronto From 2014 to 2024\*

Sindhu Priya Mallavarapu

January 25, 2024

The Covid-19 pandemic wreaked havoc on the mental health of many across the world. This paper uses a dataset about the types of person in crisis calls that were attended between 2014 and 2024 taken from OpenDataToronto to analyze the effects of the pandemic on the type of attended person in crisis calls. This dataset is analyzed to show that despite the decreasing quality of mental health, there has been no significant increase in overdoses or suicide related crisis calls attended post-pandemic. Though some reasons for this have been suggested in the paper, further study is required to understand this phenomenon.

## 1 Introduction

In 2020, the world faced a global catastrophic event that even three years later, it hasn't fully recovered from. The COVID - 19 pandemic caused many numerous challenges, including challenges with mental health. The years following 2020 were filled with talk about burnout, needing a break, or needing help to deal with the mental load of just day to day living. Social media posts, workplace meetings, and in person conversations tended to revolve around sharing what a negative affect the pandemic had on mental health. It would seem logical than to conclude that if there was such a decline in mental health for the general populace, there would be an increase in overdoses and suicide attempts. However, analyzing the data of attended person in crisis calls in Toronto shows no drastic increases in either of these categories in the attended calls.

It could seem logical to hypothesize that the reason there is no significant increase in overdose or suicide related calls attended is because there is not enough personnel to attend them. If that were the case, there there should also not be a significant increase in other types of person in crisis calls attended. However, an analysis of the data, as discussed in the Results sections

---

\*Code and data are available at: [https://github.com/MSindhuPriya/pic\\_calls](https://github.com/MSindhuPriya/pic_calls)

shows that there is a significant increase in general person in crisis calls, but no significant increase in overdose or suicide-related crisis calls. Therefore, a lack of personnel doesn't seem to be the reason for the low overdose or suicide-related calls attended. It is of significance to note that because of social distancing policies, many people were isolated. Thus, if they were overdosing or attempting suicide, it would be hard for others to notice and call in for help. Therefore, it could be that the reason for a lack of significant increase in the overdose and suicide-related category of calls attended is because there was no one around to call.

In order to perform this analysis, this paper uses data from OpenDataToronto. Specifically, it uses the "Person in Crisis Calls for Service Attended" data. The Data section of this paper goes into more detail about this dataset and the cleaning process. A sample of the cleaned data is shown in the table given below. Also in this paper is the Results section. This section analyses the data and shares any interesting findings. The Discussion section provides suggestions for possible avenues of further study and research on the findings brought out in the Results section. Finally, this paper includes a Conclusion section that summarizes the main insights discussed throughout the paper.

	event_type	event_year
1	Suicide-related	2014
2	Person in Crisis	2014
3	Suicide-related	2014
4	Suicide-related	2014
5	Person in Crisis	2014
6	Overdose	2014

## 2 Data

The data for this paper is taken from OpenDataToronto (Gelfand (2022)). Specifically, the dataset used in this paper is called "Person in Crisis Calls for Service Attended" (Service (2024)). The decision to use this dataset was made for the regular updates that are made to it and the trustworthy source that this data comes from. This dataset is updated quarterly and is sourced by the Toronto Police Service. Another dataset that was under consideration to be used in this paper was the "Mental Health Apprehensions" dataset. This was ultimately not selected due to the fact that this dataset excluded all crisis calls that were made but didn't end in an apprehension.

The selected dataset contains variables such as event\_type, event\_year, event\_month, neighbourhood\_158, neighbourhood\_140. For the purposes of this paper, only the event\_type and event\_year variables will be used. The event\_type variable contains the following categories: Person in Crisis, Suicide-related, and Overdose. All person in crisis and elopee calls are grouped into the Person in Crisis category. All attempted suicide, jumper, and threatened suicide calls are grouped into the Suicide-related category. Lastly, all overdose calls are in the

Overdose category. Additionally, event\_year is the variable that records what year the call was made ranging from 2014 to 2024.

Data cleaning and analysis was done using the statistical programming software R (R Core Team (2022)). Along with that, additional packages such as, tidyverse (Wickham et al. (2019)), janitor (Firke (2023)), ggplot2 (Wickham (2016)), knitr (Xie (2014)), dplyr (Wickham et al. (2023)) were also used. The variable names of the data were cleaned first, and then the two required variables, event\_type and event\_year, were selected into the data frame. In order to ensure that there are no problems later on in the analysis, all null values were dropped from the data. A contingency table counting the occurrence of one variable in relationship to the other is given in the table below. This table shows the count of person in crisis, overdose, and suicide-related calls in each year from 2014 to 2024.

	event_year									
event_type	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023
Overdose	1654	1673	1963	2497	2693	3274	3855	5628	3952	4555
Person in Crisis	13967	14397	15926	16361	17433	17267	19519	19210	18288	16378
Suicide-related	6608	7108	7578	8488	9573	10351	9769	10556	10831	10639

### 3 Results

Since the year 2014 there has been a steady increase in person in crisis, suicide-related, and overdose calls attended in Toronto. As is seen in the graph below, there is a spike in person in crisis calls in 2020, but a drop in suicide-related calls. The overdose calls, however, have seen the same steady increase as in the previous years. It is also of note that there is a significant increase in overdose calls in 2021, but it goes back to pre covid levels in 2022 and then continues with the steady increase in those types of calls that are consistent with years before. Also, after 2020, there is a steady decrease in the general person in crisis category of calls. The suicide-related calls are stable after the year 2020.

## Change in Type of Crisis Calls Attended (2014 – 2024)




Figure 1: Visualization of change in type of person in crisis calls attended over time.

A reason for the spike in the general category of person in crisis calls in 2020 could be explained because of the increased anxiety felt during the pandemic. As the pandemic went on, the overdose calls increased as mental health deteriorated. However, they came back to pre pandemic levels by 2022. Other than these two instances, there were no other abnormal increases in any of the categories. This is very surprising as mental health after 2020 seemed to take a turn for the worse. One reason for the lack of significant rises in overdose or suicide related calls beign attended could be because there was no one to make the call for the person that was in crisis. It could be argued that another reason for this is because there is not enough personnel to attend the overdose or suicide-related calls. If this was the case, the general person in crisis calls attended category should also not have any significant increases. However, as discussed earlier, this is not the case.

## 4 Discussion

The benefits of this dataset is that the data was well maintained and required minimal cleaning. However, the drawback is that it does not measure calls made, but only person in crisis calls attended. It would be beneficial for future papers to look at all calls that were made rather than jsut the calls that were attended.

## References

- Firke, Sam. 2023. *Janitor: Simple Tools for Examining and Cleaning Dirty Data*. <https://CRAN.R-project.org/package=janitor>.
- Gelfand, Sharla. 2022. *Opendatatoronto: Access the City of Toronto Open Data Portal*. <https://CRAN.R-project.org/package=opendatatoronto>.
- R Core Team. 2022. *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing. <https://www.R-project.org/>.
- Service, Toronto Police. 2024. *Person in Crisis Calls for Service Attended*. <https://open.toronto.ca/dataset/persons-in-crisis-calls-for-service-attended/>.
- Wickham, Hadley. 2016. *Ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York. <https://ggplot2.tidyverse.org>.
- Wickham, Hadley, Mara Averick, Jennifer Bryan, Winston Chang, Lucy D'Agostino McGowan, Romain François, Garrett Golemund, et al. 2019. "Welcome to the tidyverse." *Journal of Open Source Software* 4 (43): 1686. <https://doi.org/10.21105/joss.01686>.
- Wickham, Hadley, Romain François, Lionel Henry, Kirill Müller, and Davis Vaughan. 2023. *Dplyr: A Grammar of Data Manipulation*. <https://CRAN.R-project.org/package=dplyr>.
- Xie, Yihui. 2014. *Knitr: A Comprehensive Tool for Reproducible Research in r*. Chapman; Hall/CRC.